

Two-generator free Kleinian groups and hyperbolic displacements

İLKER S. YÜCE

The log 3 Theorem, proved by Culler and Shalen, states that every point in the hyperbolic 3-space $\mathbb{H}^3$ is moved a distance at least $\log 3$ by one of the non-commuting isometries ξ or η of $\mathbb{H}^3$ provided that ξ and η generate a torsion-free, discrete group which is not co-compact and contains no parabolic. This theorem lies in the foundation of many techniques that provide lower estimates for the volumes of orientable, closed hyperbolic 3-manifolds whose fundamental group has no 2-generator subgroup of finite index and, as a consequence, gives insights into the topological properties of these manifolds.

Under the hypotheses of the log 3 Theorem, the main result of this paper shows that every point in $\mathbb{H}^3$ is moved a distance at least $\log \sqrt{5 + 3\sqrt{2}}$ by one of the isometries ξ, η or $\xi\eta$.

[58C30](#), [20E05](#); [26B25](#), [26B35](#)

1 Introduction

Let $U^3 = \{(x, y, t) \in \mathbb{R}^3 | t > 0\}$ be the upper half 3-space and $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. The set denoted by $\hat{\mathbb{C}}$ is called the sphere at infinity when regarded as lying on the boundary of U^3 as $t = 0$. Let $\text{PSL}(2, \mathbb{C})$ be the quotient of the group of all 2×2 matrices with complex entries and determinant 1 by its center $\{\pm I\}$. Then, any element γ of $\text{PSL}(2, \mathbb{C})$ has the form

$$\gamma = \left\{ \pm \begin{pmatrix} a & b \\ c & d \end{pmatrix} \right\}$$

for some $a, b, c, d \in \mathbb{C}$ so that $ad - bc = 1$. Each element $\gamma \in \text{PSL}(2, \mathbb{C})$ acts on $\hat{\mathbb{C}}$ when it is identified with the linear fractional transformation $z \mapsto (az + b)/(cz + d)$. The action of each γ on the sphere at infinity extends to an action on U^3 as follows: γ is a product of an even number of inversions in circles and lines in $\mathbb{C}$. Each circle C and line ℓ in $\mathbb{C}$ has a unique hemisphere $\hat{C}$ and a plane $\hat{\ell}$ in U^3 , orthogonal to $\mathbb{C}$ and meeting $\mathbb{C}$ in C or ℓ . The extension of γ is the corresponding product of inversions in $\hat{C}$ and reflections in $\hat{\ell}$. This extension is called the Poincaré extension of γ . A

subgroup of $\mathrm{PSL}(2, \mathbb{C})$ is said to be *non-elementary* if it has no finite orbit in its action on $U^3 \cup \hat{\mathbb{C}}$.

The upper half 3-space U^3 is called the *upper half-space model* for the hyperbolic 3-space $\mathbb{H}^3$ (i.e., the unique three-dimensional connected and simply connected Riemannian manifold with constant sectional curvature -1) if it is equipped with the hyperbolic metric induced from the line element $ds^2 = (dx^2 + dy^2 + dt^2)/t^2$. The line element induces a distance metric on U^3 so that every Cauchy sequence in U^3 converges with respect to this metric. The inversions in hemispheres and reflections in planes are isometries of U^3 with the hyperbolic metric and generate the full group of orientation-preserving isometries denoted by $\mathrm{Isom}^+(U^3)$. Under the Poincaré extension, $\mathrm{PSL}(2, \mathbb{C})$ is isomorphic to $\mathrm{Isom}^+(U^3)$, hence, is isomorphic to the group $\mathrm{Isom}^+(\mathbb{H}^3)$ of orientation-preserving isometries of $\mathbb{H}^3$.

A subgroup Γ of $\mathrm{Isom}^+(\mathbb{H}^3)$ is called a *Kleinian group* if it is isomorphic to a discrete, non-elementary subgroup of $\mathrm{PSL}(2, \mathbb{C})$. A Kleinian group Γ acts discontinuously on $\mathbb{H}^3$ which means that the set $\{\gamma \in \Gamma \mid \gamma K \cap K \neq \emptyset\}$ is finite for any compact subset K of $\mathbb{H}^3$ ([8], Chapter 5). Since Γ acts discontinuously on $\mathbb{H}^3$, it is possible to construct a closed subset F of $\mathbb{H}^3$ for the action of Γ on $\mathbb{H}^3$ so that $\bigcup_{\gamma \in \Gamma} \gamma \cdot F = \mathbb{H}^3$, $\mathrm{int}(F) \cap \gamma \cdot \mathrm{int}(F) = \emptyset$ for every non-identity $\gamma \in \Gamma$ and the boundary of F has measure 0 ([6], Chapter 1). The set F is called a *fundamental domain* for the action of Γ on $\mathbb{H}^3$. The group Γ is said to be *co-compact* if Γ has a compact fundamental domain F ([8], Chapter 6).

Each non-identity element γ of Γ is classified by using the set of fixed points of γ . The isometry γ is said to be *elliptic* if and only if it has a finite orbit in $\mathbb{H}^3$. It is said to be *parabolic* if and only if γ fixes a point of $\hat{\mathbb{C}}$ in its action on $\hat{\mathbb{C}}$ and has no other finite orbits in $U^3 \cup \hat{\mathbb{C}}$. The isometry γ is called *loxodromic* or *hyperbolic* if it is neither elliptic nor parabolic. In this paper, we prove the following statement:

Let ξ and η be non-commuting isometries of $\mathbb{H}^3$. Suppose that ξ and η generate a torsion-free discrete group which is not co-compact and contains no parabolic. Let $\Gamma_{\dagger}$ and $\alpha_{\dagger}$ denote the set of isometries $\{\xi, \eta, \xi\eta\}$ and the real number $5 + 3\sqrt{2}$, respectively. Then, for any $z_0 \in \mathbb{H}^3$, we have

$$e^{(2 \max_{\gamma \in \Gamma_{\dagger}} \{\mathrm{dist}(z_0, \gamma \cdot z_0)\})} \geq \alpha_{\dagger},$$

which is stated as Theorem 5.1 in Section 5. Theorem 5.1 improves the lower bound given in the log 3 Theorem, proved by Culler and Shalen ([9], Theorem 9.1) which states that

Let ξ and η be non-commuting isometries of $\mathbb{H}^3$. Suppose that ξ and η generate a torsion-free, topologically tame, discrete group which is not co-compact and contains no parabolic. Let Γ_1 and α_1 denote the set of isometries $\{\xi, \eta\}$ and the real number 9, respectively. Then, for any $z_0 \in \mathbb{H}^3$, we have

$$e^{(2 \max_{\gamma \in \Gamma_1} \{\text{dist}(z_0, \gamma \cdot z_0)\})} \geq \alpha_1.$$

An orientable hyperbolic 3-manifold may be regarded as the quotient of the hyperbolic 3-space $\mathbb{H}^3$ by a discrete group Γ of orientation-preserving isometries of $\mathbb{H}^3$. If Γ is a torsion free Kleinian group and $M = \mathbb{H}^3/\Gamma$, then Γ is called *topologically tame* if M is homeomorphic to the interior of a compact 3-manifold. In [1] and [3], Agol and Calegari-Gabai prove that every finitely generated Kleinian group is topologically tame. Therefore, we drop the tameness hypothesis from Theorem 5.1.

The proof of Theorem 5.1 requires the use of the same ingredients introduced in [9] to prove the log 3 Theorem. In the following subsections of this section, we review these ingredients briefly and give the necessary definitions. In particular, we summarize the proof of the log 3 Theorem in § 1.2 with an emphasis on the calculations required to obtain the number log 3. In § 1.3, we propose an alternative technique to perform these calculations which leads to completion of one of the key steps in the proof of Theorem 5.1.

In the following, the sphere at infinity of $\mathbb{H}^3$ will be denoted by S_∞ and the canonical compactification of $\mathbb{H}^3$ will be denoted by $\overline{\mathbb{H}^3}$. Note that S_∞ is the boundary of $\overline{\mathbb{H}^3}$. Any isometry $\gamma: \mathbb{H}^3 \rightarrow \mathbb{H}^3$ extends to a conformal automorphism $\overline{\gamma}: \overline{\mathbb{H}^3} \rightarrow \overline{\mathbb{H}^3}$. The conformal automorphism of S_∞ obtained by restricting $\overline{\gamma}$ will be denoted by γ_∞ .

1.1 The Patterson-Sullivan measure

Let B^3 be the open unit ball in $\mathbb{R}^3$. The set B^3 is called the *Poincaré model* for the hyperbolic 3-space $\mathbb{H}^3$ when it is equipped with the hyperbolic metric induced from the line element $ds^2 = 4(dx^2 + dy^2 + dz^2) / (1 - r^2)^2$ where r is the Euclidean distance from the base point to the origin. For any given point $z \in \mathbb{H}^3$, there exists an isometry $h: \mathbb{H}^3 \rightarrow B^3$ which is unique modulo composition with orthogonal linear transformations of $\mathbb{R}^3$ such that $h(z) = 0$. The pull back of the metric on $S^2 = \partial B^3$ under $\overline{h}$ gives a metric on S_∞ which is uniquely determined by the point z . This metric is called the *round metric centered at z* .

If $\gamma: \mathbb{H}^3 \rightarrow \mathbb{H}^3$ is an isometry, then γ induces a conformal diffeomorphism γ_∞ of S_∞ . Let z_0 be a given point in $\mathbb{H}^3$. If S_∞ is equipped with the round metric centered

at z_0 , then at each point ζ in S_∞ , the tangent map $d\gamma_\infty: T_\zeta S_\infty \rightarrow T_{\gamma_\infty(\zeta)} S_\infty$ satisfies $|d\gamma_\infty(v)| = \lambda|v|$ for every tangent vector v at ζ where $|\cdot|$ is the length in the round metric at z_0 . The constant λ is called the *conformal expansion factor* of γ_∞ at the point ζ and will be denoted by $\lambda_{\gamma, z_0}(\zeta)$. The expression λ_{γ, z_0} gives a smooth function on S_∞ . In fact, for every $\zeta \in S_\infty$ ([9], Proposition 2.4), it is possible to obtain that

$$(1) \quad \lambda_{\gamma, z_0}(\zeta) = \mathcal{P}(z_0, \gamma^{-1}z_0, \zeta)$$

where $\mathcal{P}$ is defined as follows: let $\Pi: \mathbb{H}^3 \times \mathbb{H}^3 \times \mathbb{H}^3 \rightarrow \mathbb{R}$ be the continuous function given by $\Pi(z, z', w) = e^{\text{dist}(z, w) - \text{dist}(z', w)}$. The map Π has a unique positive continuous extension $\bar{\Pi}: \mathbb{H}^3 \times \mathbb{H}^3 \times \overline{\mathbb{H}^3} \rightarrow \mathbb{R}$. The restriction $\mathcal{P}$ of $\bar{\Pi}$ is a function on $\mathbb{H}^3 \times \mathbb{H}^3 \times S_\infty$ with the formula

$$(2) \quad \mathcal{P}(z, z', \zeta) = 1 / (\cosh \text{dist}(z, z') - \sinh \text{dist}(z, z') \cos \angle z' z \zeta)$$

where $\angle z' z \zeta$ is the angle between the ray from z to z' and the ray from z to ζ . The function $\mathcal{P}$ is called the *Poisson kernel* for $\mathbb{H}^3$. Since the function $\phi_\zeta: \mathbb{H}^3 \rightarrow \mathbb{R}$ defined by $\phi_\zeta(z) = \mathcal{P}(z_0, z, \zeta)^r$ satisfies the Laplace equation

$$(3) \quad \Delta \phi_\zeta(z) = -r(3 - r - 1)\phi_\zeta(z)$$

for any $z_0 \in \mathbb{H}^3$, $\zeta \in \mathbb{H}^3$ and $r \in \mathbb{R}$ ([9], Proposition 2.5), the Poisson kernel $\mathcal{P}$ for $\mathbb{H}^3$ provides a connection between conformal expansion factors and *D-conformal densities*, a more general concept introduced by Sullivan to construct solutions to the equation in (3).

A *D-conformal density* for S_∞ is a family $\mathcal{M} = (\mu_z)_{z \in \mathbb{H}^3}$ of finite Borel measures on S_∞ such that the equality $d\mu_{z'} = \mathcal{P}(z, z', \cdot)^D d\mu_z$ holds for any given two points $z, z' \in \mathbb{H}^3$ where D , called the *degree* of $\mathcal{M}$, is a number in $[0, 2]$. The equation $d\mu_{z'} = \mathcal{P}(z, z', \cdot)^D d\mu_z$ means that $\mu_{z'}(E) = \int_E \mathcal{P}(z, z', \cdot)^D d\mu_z$ for every measurable set $E \subset S_\infty$. A family $\mathcal{M} = (\mu_z)_{z \in \mathbb{H}^3}$ of finite Borel measures on S_∞ will be called a *conformal density* if $\mathcal{M} = (\mu_z)_{z \in \mathbb{H}^3}$ is *D-conformal* for some $D \in [0, 2]$. If $\mathcal{M} = (\mu_z)_{z \in \mathbb{H}^3}$ is a conformal density, then all the measures $\mu_z, z \in \mathbb{H}^3$ have the same support since $\mathcal{P}(z, z', \cdot)$ is a strictly positive function. The common support of the measures $\mu_z, z \in \mathbb{H}^3$ is denoted by $\text{supp } \mathcal{M}$. The family $\mathcal{A} = (A_z)_{z \in \mathbb{H}^3}$, for example, is a 2-conformal density such that $\text{supp } \mathcal{A} = S_\infty$ where A_z denotes a normalized area measure determined by the round metric at z . The density $\mathcal{A}$ is called the *area density* on S_∞ .

Let z_0 be a point in $\mathbb{H}^3$ and γ be a loxodromic isometry. Under certain assumptions, Culler and Shalen find a lower bound for the hyperbolic displacement $\text{dist}(z_0, \gamma \cdot z_0)$ using the measure $A_{z_0} \in \mathcal{A}$ as follows: let $d = \text{dist}(z_0, \gamma \cdot z_0)$, $c = \cosh d$ and

$s = \sinh d$. Let us identify $\overline{\mathbb{H}}^3$ conformally with the unit ball in $\mathbb{R}^3$ so that z_0 is mapped to the origin and $\gamma^{-1} \cdot z_0$ is on the positive vertical axis. Then the equalities in (1) and (2) give that $\lambda_{\gamma, z_0}(\zeta) = 1/(c - s \cos \phi)$ where $\phi = \phi(\zeta)$ is the angle between the positive vertical axis and the ray from the origin through ζ . Note that ϕ is the polar angle of ζ in spherical coordinates. Since S_∞ has the round metric at z_0 , the measure $A_{z_0} \in \mathcal{A}$ is obtained by dividing the area measure on the unit sphere by the area 4π of the sphere. In other words, the derivative dA_{z_0} is equal to $(1/4\pi) \sin \phi d\phi d\theta$ in spherical coordinates ϕ and θ . If C denotes the spherical cap defined by the inequality $\phi \geq \phi_0$ where $\phi_0 = \arccos(1 - 2a) \in [0, \pi]$ and $0 \leq a \leq 1$, the evaluation of an elementary iterated integral shows that $A_{z_0}(C) = a$. Assume that there exists a measure ν on S_∞ such that $\nu \leq A_{z_0}$, $\nu(S_\infty) \leq a$ and $\int_{S_\infty} \lambda_{\gamma, z_0}^2 d\nu \geq b$ for some $0 \leq b \leq 1$. The assumption $\nu \leq A_{z_0}$ gives that $A_{z_0}(C) \geq \nu(S_\infty)$ and

$$\nu(S_\infty - C) = \nu(S_\infty) - \nu(C) \leq A_{z_0}(C) - \nu(C) = (A_{z_0} - \nu)(C)$$

where $A_{z_0} - \nu$ is again a measure on S_∞ . The inequality $\nu(S_\infty - C) \leq (A_{z_0} - \nu)(C)$ combined with the fact that λ_{γ, z_0} is positive and monotone decreasing for $0 < \phi < \pi$ implies that $\inf \lambda_{\gamma, z_0}^2(C) \geq \sup \lambda_{\gamma, z_0}^2(S_\infty - C)$ and consequently

$$\begin{aligned} \int_{S_\infty - C} \lambda_{\gamma, z_0}^2 d\nu &\leq \sup(\lambda_{\gamma, z_0}^2(C)) \nu(S_\infty - C) \leq \inf(\lambda_{\gamma, z_0}^2(C)) (A_{z_0} - \nu)(C) \\ &\leq \int_C \lambda_{\gamma, z_0}^2 d(A_{z_0} - \nu). \end{aligned}$$

It follows from the assumption $\int_{S_\infty} \lambda_{\gamma, z_0}^2 d\nu \geq b$ and the inequalities given above that

$$\begin{aligned} b &\leq \int_{S_\infty} \lambda_{\gamma, z_0}^2 d\nu = \int_C \lambda_{\gamma, z_0}^2 d\nu + \int_{S_\infty - C} \lambda_{\gamma, z_0}^2 d\nu \\ &\leq \int_C \lambda_{\gamma, z_0}^2 d\nu + \int_C \lambda_{\gamma, z_0}^2 d(A_{z_0} - \nu) = \int_C \lambda_{\gamma, z_0}^2 dA_{z_0} \\ &= \int_C (c - s \cos \phi)^{-2} dA_{z_0} \\ &= \frac{1}{4\pi} \int_0^{2\pi} \int_0^{\phi_0} (c - s \cos \phi)^{-2} \sin \phi d\phi d\theta \\ (4) \quad &= \frac{a}{(c - s)(c - s + 2as)}. \end{aligned}$$

Notice that if we have $a = 0$, then we get $b \leq 0$ and if we have $b = 1$, then we get $a \geq 1$. Rewriting c and s as $(1/2)(e^d + e^{-d})$ and $(1/2)(e^d - e^{-d})$ and solving the inequality in (4) for $d = \text{dist}(z_0, \gamma \cdot z_0)$ provide a lower bound expressed explicitly in the conclusion of the following lemma:

Lemma 1.1 ([10], Lemma 2.1) *Let a and b be numbers in $[0, 1]$ which are not both equal to 0 and are not both equal to 1. Let γ be a loxodromic isometry of $\mathbb{H}^3$ and let z_0 be a point in $\mathbb{H}^3$. Suppose that ν is a measure on S_∞ such that (i) $\nu \leq A_{z_0}$, (ii) $\nu(S_\infty) \leq a$, (iii) $\int_{S_\infty} \lambda_{\gamma, z_0}^2 d\nu \geq b$. Then we have $a > 0$, $b < 1$, and*

$$\text{dist}(z_0, \gamma \cdot z_0) \geq \frac{1}{2} \log \frac{b(1-a)}{a(1-b)}.$$

The existence of such a measure ν used in Lemma 1.1 requires the investigation of the decompositions of the Patterson density and the Patterson-Sullivan measure.

Let Γ be a non-elementary discrete subgroup of $\text{Isom}^+(\mathbb{H}^3)$ and $\mathcal{M} = (\mu_z)_{z \in \mathbb{H}^3}$ be a D -conformal density and $\gamma: \mathbb{H}^3 \rightarrow \mathbb{H}^3$ be an isometry in Γ . The family of Borel measures $(\gamma_\infty^* \mu_{\gamma z})_{z \in \mathbb{H}^3}$ gives a D -conformal density where $\gamma_\infty^* \mu_{\gamma z}$ is the pull-back of μ_z defined by $(\gamma_\infty^* \mu_{\gamma z})(E) = \mu_z(\gamma_\infty(E))$ for every Borel set $E \subset S_\infty$. The family of measures $(\gamma_\infty^* \mu_{\gamma z})$, denoted by $\gamma^* \mathcal{M}$, is called the *pull-back* of $\mathcal{M} = (\mu_z)_{z \in \mathbb{H}^3}$. A conformal density $\mathcal{M}$ is said to be Γ -invariant if it satisfies that $\gamma^* \mathcal{M} = \mathcal{M}$ for every $\gamma \in \Gamma$.

A point a of S_∞ is called a *limit point* of the group Γ if there is a point x in $\mathbb{H}^3$ and a sequence $\{\gamma_j\}_{j=1}^\infty \subset \Gamma$ such that $\{\gamma_j \cdot x\}_{j=1}^\infty$ converges to a . If Λ_Γ denotes the set of all limit points of Γ , Patterson [7] proves that

Theorem 1.2 ([9], Corollary 4.2) *There is a Γ -invariant conformal density for $\mathbb{H}^3$, denoted by $\mathcal{M}_{\Gamma, z_0} = (\mu_z)_{z \in \mathbb{H}^3}$, with support Λ_Γ .*

In the following, $\mathcal{M}_{\Gamma, z_0}$ will be called the *Patterson density*. The member μ_{z_0} of the Patterson density will be referred as the *Patterson-Sullivan measure at z_0* on S_∞ . Note that if $\mathcal{M} = (\mu_z)_{z \in \mathbb{H}^3}$ is a D -conformal Γ -invariant density, then for any $z \in \mathbb{H}^3$ and $\gamma \in \Gamma$, the equalities $d(\gamma^* \mu_z) = d\mu_{\gamma^{-1}z}$ and $d\mu_{\gamma^{-1}z} = \mathcal{P}(z, \gamma^{-1}z, \cdot)^D d\mu_z$ follow so that

$$(5) \quad d(\gamma^* \mu_z) = \lambda_{\gamma, z}^D d\mu_z$$

The last equation is obtained from the expression in (1).

A subset W of $\mathbb{H}^3$ is called *uniformly discrete* if there exists a positive number ϵ such that $\text{dist}(z, w) > \epsilon$ for any two distinct points z, w of $\mathbb{H}^3$. The closure of W in $\mathbb{H}^3$ is a union of the form $W \cup \Lambda_W$ where Λ_W , the limit set of W , is a closed subset of S_∞ . Note that the orbit $W = \Gamma \cdot z$ is a uniformly discrete set for any $z \in \mathbb{H}^3$ since Γ is discrete and z has compact stabilizer in $\text{Isom}^+(\mathbb{H}^3)$. Let W_0 denote the orbit $\Gamma \cdot z_0$. Theorem 1.2 is a corollary of the following proposition

Proposition 1.3 ([9], Proposition 4.2) *Let W be an infinite, uniformly discrete subset of $\mathbb{H}^3$, and let $\mathcal{V}$ be a countable collection of subsets of W . Suppose $W \in \mathcal{V}$. Then there exists a number $D \in [0, 2]$ and a family $(\mathcal{M}_V)_{V \in \mathcal{V}}$ of D -conformal densities for $\mathbb{H}^3$, indexed by the collection $\mathcal{V}$, satisfying the following conditions:*

- (i) $\mathcal{M}_W \neq 0$.
- (ii) *For any finite family $(V_i)_{1 \leq i \leq m}$ of disjoint sets in $\mathcal{V}$ such that $V = \bigcup_{i=1}^m V_i \in \mathcal{V}$, we have $\mathcal{M}_V = \sum_{i=1}^m \mathcal{M}_{V_i}$.*
- (iii) *For any $V \in \mathcal{V}$ and any isometry $\gamma: \mathbb{H}^3 \rightarrow \mathbb{H}^3$ such that $\gamma V \in \mathcal{V}$, we have $\gamma_\infty^*(\mathcal{M}_{\gamma V}) = \mathcal{M}_V$.*
- (iv) *For any $V \in \mathcal{V}$, the support of $\mathcal{M}_V$ is contained in the limit set Λ_V of V . In particular, for any finite set $V \in \mathcal{V}$ we have $\mathcal{M}_V = 0$,*

when W is set as W_0 and $\mathcal{V}$ is chosen as $\{W_0\}$, i.e., $\mathcal{M}_{W_0} = \mathcal{M}_{\Gamma, z_0}$. In the proposition above, $\sum_{i=1}^m \mathcal{M}_{V_i}$ denotes the sum $(\sum_{i=1}^m \mu_{V_i, z})_{z \in \mathbb{H}^3}$ of the D -conformal densities $\mathcal{M}_{V_i} = (\mu_{V_i, z})_{z \in \mathbb{H}^3}$. The sum $\sum_{i=1}^m \mathcal{M}_{V_i}$ is also a D -conformal density. The expression $\mathcal{M}_V = 0$ means that if $\mathcal{M}_V = (\mu_{V, z})_{z \in \mathbb{H}^3}$ then $\mu_{V, z} = 0$ for every $z \in \mathbb{H}^3$. Proposition 1.3 allows to construct decompositions of the Patterson density and consequently decompositions of the Patterson-Sullivan measure corresponding to decompositions of the group Γ .

1.2 A Paradoxical decomposition of Γ and Proof of the log 3 theorem

Let ξ and η be non-commuting isometries of $\mathbb{H}^3$. Suppose that ξ and η generate a torsion-free discrete group which is topologically tame, is not co-compact and contains no parabolic. It is possible to show that $\langle \xi, \eta \rangle$ is a free group of rank 2 ([9], Proposition 9.2). Let $\Gamma = \langle \xi, \eta \rangle$. Then the proof of the log 3 theorem is divided into two parts: if $\Lambda_\Gamma = S_\infty$ or $\Lambda_\Gamma \neq S_\infty$.

Assume that $\Lambda_\Gamma = S_\infty$. Since Γ is a free group on the generating set $\Xi = \{\xi, \eta\}$, every element γ of Γ can be written uniquely as a reduced word $\psi_1 \cdots \psi_m$, where $m \geq 0$, each ψ_i is an element of $\Psi^1 = \Xi \cup \Xi^{-1}$, and $\psi_{i+1} \neq \psi_i^{-1}$ for $i = 1, \dots, m-1$. If $\gamma \neq 1$, ψ_1 is called the *initial letter* of γ . For each $\psi \in \Psi^1$, let J_ψ be the set of all non-trivial elements of Γ that have initial letter ψ . Then Γ is a disjoint union $\{1\} \cup \bigcup_{\psi \in \Psi^1} J_\psi$ which will be referred to as the *paradoxical decomposition* of Γ .

Let z_0 be a given point in $\mathbb{H}^3$. If V_ψ denotes the set $\{\gamma \cdot z_0 : \gamma \in J_\psi\}$, the orbit $W_0 = \Gamma \cdot z_0$ is a disjoint union $W_0 = \{z_0\} \cup \bigcup_{\psi \in \Psi^1} V_\psi$. Observe that W_0 is uniformly

discrete since Γ is discrete and torsion-free. Let $\mathcal{V}$ be the finite collection of all subsets of W_0 containing all the sets of the form $\bigcup_{\psi \in \Psi'} V_\psi$ or $\{z_0\} \cup \bigcup_{\psi \in \Psi'} V_\psi$ for $\Psi' \subset \Psi^1$. With these choices of W_0 and $\mathcal{V}$, Proposition 1.3 gives a number $D \in [0, 2]$ and a family $(\mathcal{M}_V)_{V \in \mathcal{V}}$ of D -conformal densities so that the conditions (i) – (iv) are satisfied. Part (iii) of Proposition 1.3 implies that $\mathcal{M}_{W_0}$ is Γ -invariant. The definition of a D -conformal density and part (i) of Proposition 1.3 show that $\mu_{W_0, z_0} \neq 0$. After normalization, it is possible to assume that $\mu_{W_0, z_0}(S_\infty) = 1$. It follows from the equality $W_0 = \{z_0\} \cup \bigcup_{\psi \in \Psi^1} V_\psi$ and part (ii) of Proposition 1.3 that $\mu_{W_0, z_0} = \mu_{\{z_0\}, z_0} + \sum_{\psi \in \Psi^1} \mu_{V_\psi, z_0}$. Since $\{z_0\}$ is finite, the measure $\mu_{\{z_0\}, z_0} = 0$ by Proposition 1.3 part (iv). Note that $\mu_{W_0, z_0} = \mu_{z_0}$. Then the decomposition of the Patterson-Sullivan measure μ_{z_0} corresponding to the paradoxical decomposition of Γ is obtained as $\mu_{z_0} = \sum_{\psi \in \Psi^1} \mu_{V_\psi, z_0}$.

The definition of the sets J_ψ gives the group theoretical identities $\psi J_{\psi^{-1}} = \Gamma - J_\psi$ so that $\psi V_{\psi^{-1}} = W_0 - V_\psi$ for every $\psi \in \Psi^1$. Since the set $W_0 - V_\psi = \{z_0\} \cup \bigcup_{\psi' \neq \psi} V_{\psi'}$ is in $\mathcal{V}$, Proposition 1.3(iii) implies that $\mathcal{M}_{V_{\psi^{-1}}} = \psi_\infty^*(\mathcal{M}_{W_0 - V_\psi})$. Proposition 1.3 part (ii) shows that $\mathcal{M}_{W_0} = \mathcal{M}_{W_0 - V_\psi} + \mathcal{M}_{V_\psi}$ for $\psi \in \Psi^1$. Therefore, the equality $\mathcal{M}_{V_{\psi^{-1}}} = \psi_\infty^*(\mathcal{M}_{W_0 - V_\psi})$ can be rewritten as $\mathcal{M}_{V_{\psi^{-1}}} = \psi_\infty^*(\mathcal{M}_{W_0} - \mathcal{M}_{V_\psi})$ which implies that $\mu_{V_{\psi^{-1}}, \psi(z_0)} = \psi_\infty^*(\mu_{z_0} - \mu_{V_\psi, z_0})$. Then it follows from the equalities given in (4) that

$$(6) \quad \int_{S_\infty} d(\psi_\infty^*(\mu_{z_0} - \mu_{V_\psi, z_0})) = \int_{S_\infty} \lambda_{\psi, z_0}^D d\mu_{V_{\psi^{-1}}, \psi(z_0)}.$$

Since $\mathcal{M}_{V_{\psi^{-1}}}$ is a D -conformal Γ -invariant density for every $\psi \in \Psi$ by Proposition 1.3(iii), the equality in (6) gives that

$$\int_{S_\infty} \lambda_{\psi, z_0}^D d\mu_{V_{\psi^{-1}}, \psi(z_0)} = 1 - \int_{S_\infty} d\mu_{V_\psi, z_0}.$$

Let us say $\mu_{V_\psi, z_0} = \nu_\psi$ for each $\psi \in \Psi^1$. With the convention that every integral in this text, unless otherwise is stated, will be calculated over the sphere at infinity, the discussion given above proves the following lemma

Lemma 1.4 ([9], Lemma 5.3) *Let Γ be a free Kleinian group on a generating set $\Xi = \{\xi, \eta\}$. Set $\Psi^1 = \Xi \cup \Xi^{-1} \subset \Gamma$. Let z_0 be any point of $\mathbb{H}^3$. Then there exists a number $D \in [0, 2]$, a Γ -invariant D -conformal density $\mathcal{M}_{\Gamma, z_0} = (\mu_z)_{z \in \mathbb{H}^3}$ for $\mathbb{H}^3$ and a family $(\nu_\psi)_{\psi \in \Psi^1}$ of Borel measures on S_∞ such that*

- (1) $\mu_{z_0}(S_\infty) = 1$; (2) $\mu_{z_0} = \sum_{\psi \in \Psi^1} \nu_\psi$; and
- (3) for each $\psi \in \Psi^1$ we have $\int \lambda_{\psi, z_0}^D d\nu_{\psi^{-1}} = 1 - \int d\nu_\psi$.

If Ξ is a two-element set $\{\xi, \eta\}$ and if $z_0 \in \ell(\xi, \eta)$ where $\ell(\xi, \eta)$ denotes the common perpendicular of the isometries ξ and η , then we also have

$$(4) \quad \int d\nu_{\xi^{-1}} = \int d\nu_{\xi} \text{ and } \int d\nu_{\eta^{-1}} = \int d\nu_{\eta}.$$

The last part of Lemma 1.4 follows from the fact that there is a unique geodesic line $\ell \subset \mathbb{H}^3$ such that the 180° rotation τ_ℓ about ℓ satisfies $\tau_\ell \xi \tau_\ell = \xi^{-1}$ and $\tau_\ell \eta \tau_\ell = \eta^{-1}$ when ξ_∞ and η_∞ have no common fixed point in S_∞ ([9], Proposition 1.2). This line is the common perpendicular $\ell(\xi, \eta)$ to the axis of ξ and η . Note that $\tau_\ell \cdot z = z$ for every $z \in \ell$. Let z_0 be a point on ℓ and $\psi \in \Psi^1$. The group theoretical identity $\tau_\ell J_\psi \tau_\ell = J_{\psi^{-1}}$ implies that $\tau_\ell V_\psi = V_{\psi^{-1}}$ where $V_\psi = \{\gamma \cdot z_0 : \gamma \in J_\psi\}$ and $\tau_\ell V_\psi = \{\tau_\ell \gamma \tau_\ell \cdot z_0 : \gamma \in J_\psi\}$. Then part (iii) of Proposition 1.3 gives that $(\tau_\ell)_\infty^* \mathcal{M}_{V_\psi} = \mathcal{M}_{V_{\psi^{-1}}}$ which means $(\tau)_\infty^* \mu_{V_\psi, z_0} = \mu_{V_{\psi^{-1}}, z_0}$ or $(\tau)_\infty^* \nu_\psi = \nu_{\psi^{-1}}$. Since $\mathcal{M}_{V_\psi}$ is a D -conformal Γ -invariant density by part (iii) of Proposition 1.3, integrating both sides of the last equality provides the part (4) of Lemma 1.4, i.e.,

$$\int d\nu_\psi = \int d\nu_{\psi^{-1}}$$

for $\psi \in \Psi^1$. For the rest of this subsection, let us assume without loss of generality that $z_0 \in \ell(\xi, \eta)$ because $\max_{\gamma \in \Xi} \{\text{dist}(z_0, \gamma \cdot z_0)\} \geq \max_{\gamma \in \Xi} \{\text{dist}(z_1, \gamma \cdot z_1)\}$ holds for z_0 and z_1 where z_1 denotes the orthogonal projection of z_0 onto $\ell(\xi, \eta)$.

Let us consider the Patterson density $\mathcal{M}_{\Gamma, z_0} = (\mu_z)_{z \in \mathbb{H}^3}$ and the Patterson-Sullivan measure μ_{z_0} introduced in Lemma 1.4. It is possible to show that $\mathcal{M}_{\Gamma, z_0}$ has a strictly positive degree, i.e., $D > 0$ ([9], Proposition 3.8). Since $\mathcal{M}_{\Gamma, z_0}$ is a D -conformal Γ -invariant density, the non-negative real-valued function $u = u_{\mathcal{M}_{\Gamma, z_0}}$ defined by $u(z) = \mu_z(S_\infty)$ is Γ -invariant, i.e., $u(\gamma \cdot z) = u(z)$ for every $\gamma \in \Gamma$. Therefore, the equality $u(z) = \mu_z(S_\infty)$ where $\mu_z(S_\infty) = \int \mathcal{P}(z_0, z, \cdot)^D d\mu_{z_0}$ can be written in the form

$$(7) \quad u(z) = \int \phi_\zeta(z) d\mu_{z_0}$$

so that $\phi_\zeta(z) = \mathcal{P}(z_0, z, \zeta)^D$. Observe that u is a C^∞ function because ϕ_ζ is a C^∞ function. Since the function $\phi_\zeta(z) = \mathcal{P}(z_0, z, \zeta)^D$ satisfies the Laplace equation (3), the differentiation of the equation in (7) under the integral sign shows that u also satisfies the Laplace equation $\Delta u = -D(3 - D - 1)u$. Note that we have $-D(3 - D - 1) < 0$ since $D > 0$. Therefore u is a superharmonic function on $\mathbb{H}^3$ (A C^∞ function f on a Riemannian manifold is called *superharmonic* if it satisfies the inequality $\Delta f < 0$). It is known that every superharmonic function on $M = \mathbb{H}^3/\Gamma$ is constant when Γ is a topologically tame, Kleinian group which contains no parabolics with $\Lambda_\Gamma = S_\infty$

([9], Proposition 6.8 and [4], Theorem 7.2). This fact implies that every Γ -invariant superharmonic function on $\mathbb{H}^3$ is constant which means that $u(z) = C$, or equivalently $\mu_z(S_\infty) = C$ for every $z \in \mathbb{H}^3$ for some constant C . On the other hand, the conformal density $C\mathcal{A} = (CA_z)_{z \in \mathbb{H}^3}$ gives the same superharmonic function $u_{C\mathcal{A}}(z) = C$ for every $z \in \mathbb{H}^3$. In other words, the expression $u_{\mathcal{M}_{\Gamma, z_0}} = u_{C\mathcal{A}}$ holds. As a result, the equality $\mathcal{M}_{\Gamma, z_0} = C\mathcal{A}$ holds, and consequently, gives that $D = 2$ ([9], Proposition 3.7). Part (1) of Lemma 1.4 provides that $C = 1$ because $\mu_{z_0}(S_\infty) = CA_{z_0}(S_\infty)$ where $A_{z_0}(S_\infty) = 1$. Therefore the Paterson density is equal to the area density $\mathcal{M}_{\Gamma, z_0} = \mathcal{A}$. In particular, the Patterson-Sullivan measure is equal to the area measure, i.e., $\mu_{z_0} = A_{z_0}$.

Conclusions (2) and (4) of Lemma 1.4 show that $1 = 2\nu_\xi(S_\infty) + 2\nu_\eta(S_\infty)$ which implies that either $\nu_\xi(S_\infty) \leq 1/4$ or $\nu_\eta(S_\infty) \leq 1/4$. If the inequality $\nu_\xi(S_\infty) \leq 1/4$ holds, parts (3) and (4) of Lemma 1.4 imply that $\int \lambda_{\xi^{-1}, z_0}^2 d\nu_\xi = 1 - \nu_\xi \geq 3/4$. Then we let $a = 1/4$, $b = 3/4$, $\nu = \nu_\xi$ and $\gamma = \xi^{-1}$. Since we have $\nu_\xi \leq A_{z_0}$ by Lemma 1.4 (1), it is possible to apply Lemma 1.1 with these choices of a and b and conclude that $\text{dist}(z_0, \xi^{-1} \cdot z_0) \geq (\log 9)/2$. Similarly, if the inequality $\nu_\eta(S_\infty) \leq 1/4$ holds, parts (3) and (4) of Lemma 1.4 imply that $\int \lambda_{\eta^{-1}, z_0}^2 d\nu_\eta = 1 - \nu_\eta \geq 3/4$. Then we let $a = 1/4$, $b = 3/4$, $\nu = \nu_\eta$ and $\gamma = \eta^{-1}$. Since we have $\nu_\eta \leq A_{z_0}$ by Lemma 1.4 (1), it is possible to apply Lemma 1.1 with these choices of a and b and see that $\text{dist}(z_0, \eta^{-1} \cdot z_0) \geq (\log 9)/2$. Remember that $\text{dist}(z_0, \xi \cdot z_0) = \text{dist}(z_0, \xi^{-1} \cdot z_0)$ and $\text{dist}(z_0, \eta \cdot z_0) = \text{dist}(z_0, \eta^{-1} \cdot z_0)$. Therefore, the following proposition completes the proof of the log 3 Theorem in the case $\Lambda_\Gamma = S_\infty$:

Proposition 1.5 ([9], Proposition 5.2) *Let ξ and η be two orientation-preserving isometries of $\mathbb{H}^3$. Suppose that the group Γ generated by ξ and η is discrete and is free on the generators ξ and η . Suppose in addition that every Γ -invariant conformal density for $\mathbb{H}^3$ is a constant multiple of the area density. If $\Gamma_1 = \{\xi, \eta\}$ and $\alpha_1 = 9$, then for any $z_0 \in \mathbb{H}^3$ we have*

$$e^{(2 \max_{\gamma \in \Gamma_1} \{\text{dist}(z_0, \gamma \cdot z_0)\})} \geq \alpha_1.$$

For the second part of the proof the log 3 Theorem, let us assume that $\Lambda_\Gamma \neq S_\infty$ where $\Gamma = \langle \xi, \eta \rangle$. Let V denotes the variety $PSL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$ which will be considered with the complex topology. Let us denote by $\mathfrak{D}$ the set of all points $(\xi', \eta') \in V$ such that the group $\langle \xi', \eta' \rangle$ generated by ξ' and η' is a free Kleinian group of rank 2. $\mathfrak{D}$ is a closed subset of V ([2]). If $\mathfrak{G}\mathfrak{F}$ denotes the set of all points $(\xi', \eta') \in \mathfrak{D}$ such that $\Lambda_{\Gamma'} \neq S_\infty$ where $\Gamma' = \langle \xi', \eta' \rangle$, then $\mathfrak{G}\mathfrak{F}$ is an open subset of V ([5], Theorem 8.1). Note that $(\xi, \eta) \in \mathfrak{G}\mathfrak{F}$. Let $\mathfrak{B} = \overline{\mathfrak{G}\mathfrak{F}} - \mathfrak{G}\mathfrak{F}$ denote the frontier of $\mathfrak{G}\mathfrak{F}$ in V . There exists a dense subset $\mathfrak{C}$ of $\mathfrak{B}$ such that every superharmonic function on $M = \mathbb{H}^3/\Gamma'$

is constant for every $(\xi', \eta') \in \mathfrak{C}$ ([9], Theorem 8.2). If (ξ', η') is in $\mathfrak{B}$, then the limit set of the group $\langle \xi', \eta' \rangle$ is the sphere at infinity. Let z_0 be a fixed point in $\mathbb{H}^3$. Let us define the function

$$\begin{aligned} f_{z_0}^1 : \mathfrak{D} &\longrightarrow \mathbb{R} \\ (\xi', \eta') &\mapsto \max(\text{dist}(z_0, \xi' \cdot z_0), \text{dist}(z_0, \eta' \cdot z_0)). \end{aligned}$$

Since $(f_{z_0}^1)^{-1}([0, B]) = \{(\xi', \eta') \in V : \max(\text{dist}(z_0, \xi' \cdot z_0), \text{dist}(z_0, \eta' \cdot z_0)) \leq B\}$ is compact for every $B \geq 0$, the function $f_{z_0}^1$ is a continuous, proper, non-negative valued function on V . Therefore, $f_{z_0}^1$ must take a minimum value on $\overline{\mathfrak{G}\mathfrak{F}}$. Let $(\xi_0, \eta_0) \in \overline{\mathfrak{G}\mathfrak{F}}$ be a point at which $f_{z_0}^1$ takes a minimum value. The fact that every superharmonic function on $M = \mathbb{H}^3/\Gamma'$ is constant for every $(\xi', \eta') \in \mathfrak{C}$ implies that $\mathcal{M}_{\Gamma', z_0} = \mathcal{A}$ for every $(\xi', \eta') \in \mathfrak{C}$ where $\Gamma' = \langle \xi', \eta' \rangle$. Then Proposition 1.5 gives that $f_{z_0}^1(\xi', \eta') \geq \log 3$ for every $(\xi', \eta') \in \mathfrak{C}$. If (ξ_0, η_0) is in $\mathfrak{B}$, then $f_{z_0}^1(\xi_0, \eta_0) \geq \log 3$ since $\mathfrak{C}$ is a dense subset of $\mathfrak{B}$. Hence, it is enough to show that $f_{z_0}^1$ has no local minimum on the open set $\mathfrak{G}\mathfrak{F}$.

Let us assume that (ξ_0, η_0) is in $\mathfrak{G}\mathfrak{F}$. Then $f_{z_0}^1$ has a local minimum at (ξ_0, η_0) . Since ξ_0, η_0 have infinite orders in $\langle \xi_0, \eta_0 \rangle$, we have $\xi_0 \cdot z_0 \neq z_0$ and $\eta_0 \cdot z_0 \neq z_0$. Then there exists a sequence (z_i) of points on the open hyperbolic line between z_0 and $\xi_0 \cdot z_0 \neq z_0$ which converges to $\xi_0 \cdot z_0$. Also there exists a sequence (w_i) of points on the open hyperbolic line between z_0 and $\eta_0 \cdot z_0 \neq z_0$ which converges to $\eta_0 \cdot z_0$. Let (ξ_i) and (η_i) be two sequences of hyperbolic isometries converging to identity such that $\xi_i \cdot z_0 = z_i$ and $\eta_i \cdot z_0 = w_i$. Then the sequence $(\xi_i \xi, \eta_i \eta)$ converges to (ξ_0, η_0) and $f_{z_0}^1(\xi_i \xi, \eta_i \eta) < f_{z_0}^1(\xi_0, \eta_0)$ for every i . Therefore, (ξ_0, η_0) is not a local minimum of $f_{z_0}^1$. Hence, (ξ_0, η_0) is in $\mathfrak{B}$. Since, the limit set Λ_Γ of $\Gamma = \langle \xi, \eta \rangle$ is not the sphere at infinity, (ξ, η) is in $\mathfrak{G}\mathfrak{F}$. Therefore, we find that $f_{z_0}^1(\xi, \eta) = \max(\text{dist}(z_0, \xi \cdot z_0), \text{dist}(z_0, \eta \cdot z_0)) \geq \log 3$. This finishes the proof of the $\log 3$ Theorem.

1.3 An alternative technique to calculate the number $\log 3$

Since the Patterson density $\mathcal{M}_{\Gamma, z_0}$ is the area density $\mathcal{A}$ when $\Lambda_\Gamma = S_\infty$, the calculation of the number $\log 3$ follows from the application of Lemma 1.1 with the choices $\gamma = \psi^{-1}$, $\nu = \nu_\psi$, $a = 1/4$ and $b = 3/4$ where $\psi \in \Gamma_1 = \{\xi, \eta\}$. These choices of a and b follow from the equalities $\mu_\xi = \mu_{\xi^{-1}}$ and $\mu_\eta = \mu_{\eta^{-1}}$ given in Lemma 1.4 part (4) which requires the use of a point z_0 on the common perpendicular $\ell(\xi, \eta)$ of ξ and η .

We calculate the number $\log 3$ without referring to $\ell(\xi, \eta)$ as follows: Let us apply Lemma 1.1 with choices $\gamma = \xi^{-1}$, $\nu = \nu_\xi$, $a = \nu_{\xi^{-1}}(S_\infty)$, $b = \int \lambda_{\xi^{-1}, z_0}^2 d\nu$ and $\gamma = \eta^{-1}$, $\nu = \nu_\eta$, $a = \nu_{\eta^{-1}}(S_\infty)$, $b = \int \lambda_{\eta^{-1}, z_0}^2 d\nu$ where $\int \lambda_{\psi^{-1}, z_0}^2 d\nu = 1 - \int d\nu_\psi$ for every $\psi \in \Gamma_1 = \{\xi, \eta\}$ by Lemma 1.4(3). Then the conclusion of Lemma 1.1 gives the inequalities

$$(8) \quad e^{2\text{dist}(z_0, \gamma \cdot z_0)} \geq \frac{(1 - \int d\nu_{\gamma^{-1}})(1 - \int d\nu_\gamma)}{\int d\nu_{\gamma^{-1}} \cdot \int d\nu_\gamma}$$

for $\gamma \in \{\xi^{-1}, \eta^{-1}\}$. We propose to consider the constants on the right-hand side of the inequalities given above as the values of the functions

$$f_1^1(x_1, x_2, x_3, x_4) = \frac{1 - x_4}{x_4} \cdot \frac{1 - x_1}{x_1} \quad \text{and} \quad f_2^1(x_1, x_2, x_3, x_4) = \frac{1 - x_3}{x_3} \cdot \frac{1 - x_2}{x_2}$$

at $(\int d\nu_\xi, \int d\nu_{\xi^{-1}}, \int d\nu_\eta, \int d\nu_{\eta^{-1}}) \in \mathbb{R}^4$ with $\int d\nu_\xi + \int d\nu_{\xi^{-1}} + \int d\nu_\eta + \int d\nu_{\eta^{-1}} = 1$. Then the calculation of the number $\log 3$ in the proof of the $\log 3$ Theorem becomes a consequence of the statement

$$\inf_{\mathbf{x} \in \Delta^3} \{ \max(f_1^1(\mathbf{x}), f_2^1(\mathbf{x})) \} = 9$$

where $\Delta^3 = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}_+^4 : x_1 + x_2 + x_3 + x_4 = 1\}$. In particular, the calculation of the number $\log 3$ reduces the calculation of the infimum of the maximum of f_1^1 and f_2^1 over the simplex Δ^3 . The functions f_1^1 and f_2^1 will be called the *displacement functions* for the paradoxical decomposition $\Gamma_{\mathcal{D}^1}$ of Γ .

This calculation technique of the number $\log 3$ combined with the arguments developed by Culler and Shalen to prove the $\log 3$ Theorem suggests a step-by-step process to find a lower bound for the maximum of the displacements under the isometries in $\Gamma_* \subseteq \Psi^* \cup \Psi_r^*$ for any given decomposition $\Gamma_{\mathcal{D}^*}$ of Γ where $\mathcal{D}^* = (\Psi^*, \Psi_r^*)$ (see § 3).

1.4 Lower bounds for displacements for a given decomposition of Γ

Let z_0 be a given point in $\mathbb{H}^3$. Let $\Gamma_{\mathcal{D}^*}$ be a decomposition (see § 3) of Γ such that $\mathcal{D}^* = (\Psi^*, \Psi_r^*)$, and Γ_* be a subset of $\Psi^* \cup \Psi_r^*$. We enumerate the elements of the subset $\Gamma_* = \{\gamma_i \in \Gamma : i \in I_*\}$ for some index set I_* .

Similar to the group-theoretical relations $\gamma J_{\gamma^{-1}} = \Gamma - J_\gamma$ for $\gamma \in \{\xi, \xi^{-1}, \eta, \eta^{-1}\}$ obtained from the paradoxical decomposition $\Gamma_{\mathcal{D}^1}$ of Γ with $\mathcal{D}^1 = (\Psi^1, \emptyset)$, the decomposition $\Gamma_{\mathcal{D}^*}$ of Γ provides group-theoretical relations between the sets of words J_ψ for $\psi \in \Psi^*$. We apply Proposition 1.3 to the orbits V_ψ of these sets J_ψ for $\psi \in \Psi^*$ and obtain an analog of Lemma 1.4 for $\Gamma_{\mathcal{D}^*}$. In other words, we

show that the Patterson-Sullivan measure μ_{z_0} can be decomposed as a sum of Borel measures ν_ψ for $\psi \in \Psi^*$ so that $\sum_{\psi \in \Psi^*} \nu_\psi = 1$ after normalization and determine all measure-theoretical relations between the Borel measures ν_ψ for $\psi \in \Psi^*$ using the group-theoretical relations obtained from $\Gamma_{\mathcal{D}^*}$.

We apply Lemma 1.1 and use the measure-theoretical relations between the Borel measures ν_ψ for $\psi \in \Psi^*$ to produce inequalities similar to (8) for $\Gamma_{\mathcal{D}^*}$. These inequalities provide lower bounds for the displacements under every isometry $\psi \in \Psi^* \cup \Psi_r^*$. The expressions on the right-hand side of these inequalities provided by Lemma 1.1 can be considered as the values of certain functions obtained by replacing each total measure with a variable as we proposed in § 1.3 at a point in the simplex defined by $\Delta^{|\Psi^*|-1} = \{\mathbf{x} \in \mathbb{R}^{|\Psi^*|} : \sum_{i=1}^{|\Psi^*|} x_i = 1, 0 < x_i < 1\}$ where $|\Psi^*|$ denotes the cardinality of Ψ^* . The functions obtained in this way will be referred to as the *displacement functions* for the decomposition $\Gamma_{\mathcal{D}^*}$. Since Γ_* is a subset of $\Psi^* \cup \Psi_r^*$, some of the displacement functions belong to the isometries in Γ_* . We calculate the infimum of maximum of these displacement functions over the simplex $\Delta^{|\Psi^*|-1}$.

As the $\log$ function is strictly increasing, the maximum of the lower bounds for the displacements under isometries in Γ_* is greater than or equal to the infimum of the maximum of the displacement functions that belong to the isometries in Γ_* over the simplex $\Delta^{|\Psi^*|-1}$. This completes the case $\Lambda_\Gamma = S_\infty$ where every Γ -invariant conformal density is a constant multiple of the area density. In other words, we conclude the case $\Lambda_\Gamma = S_\infty$ by proving an analog of Proposition 1.2 for the isometries in Γ_* in the decomposition $\Gamma_{\mathcal{D}^*}$.

The last stage of this process covers the case $\Lambda_\Gamma \neq S_\infty$. Since $\Gamma = \langle \xi, \eta \rangle$ is a free group on the generators ξ and η , every γ_i is of the form $\psi_1^i \cdots \psi_m^i$, where $m \geq 0$, each ψ_j^i is an element of $\{\xi, \xi^{-1}, \eta, \eta^{-1}\}$, and $\psi_{j+1}^i \neq (\psi_j^i)^{-1}$ for $j = 1, \dots, m-1$ and $i \in I_*$. Let $\Gamma' = \langle \xi', \eta' \rangle$ where $(\xi', \eta') \in \mathcal{D}$. Let us define the function π between the set $\{\xi, \xi^{-1}, \eta, \eta^{-1}\}$ and $\{\xi', (\xi')^{-1}, \eta', (\eta')^{-1}\}$ such that $\pi(\xi) = \xi'$, $\pi(\xi^{-1}) = (\xi')^{-1}$, $\pi(\eta) = \eta'$ and $\pi(\eta^{-1}) = (\eta')^{-1}$. Let γ'_i be the word in Γ' defined as $\pi(\psi_1^i) \cdots \pi(\psi_m^i)$ for every $i \in I_*$. Then, we show that $f_{z_0}^*$ does not have a local minimum on $\mathfrak{G}\mathfrak{F}$ where

$$\begin{aligned} f_{z_0}^* &: \mathcal{D} \longrightarrow \mathbb{R} \\ (\xi', \eta') &\longmapsto \max_{\gamma \in \Gamma'_*} (\text{dist}(z_0, \gamma \cdot z_0)). \end{aligned}$$

and $\Gamma'_* = \{\pi(\gamma) : \gamma \in \Gamma_*\}$. For the rest of this text, we will follow the steps described above for the set of isometries $\Gamma_\dagger = \{\xi, \eta, \xi\eta\}$ to prove Theorem 5.1.

The organization of this paper is as follows: In Section 2, we give the necessary calculations in detail to obtain the number $\log 3$ for the $\log 3$ Theorem following the

process we outlined above. In other words, we show that the infimum of the maximum of $f_1^1(\mathbf{x})$ and $f_2^1(\mathbf{x})$ over the simplex Δ^3 is $\alpha_1 = 9$ which follows from the facts

- A. $\inf_{\mathbf{x} \in \Delta^3} \{ \max (f_1^1(\mathbf{x}), f_2^1(\mathbf{x})) \} = \min_{\mathbf{x} \in \Delta^3} \{ \max (f_1^1(\mathbf{x}), f_2^1(\mathbf{x})) \},$
- B. $\min_{\mathbf{x} \in \Delta^3} \{ \max (f_1^1(\mathbf{x}), f_2^1(\mathbf{x})) \} = f_1^1(\mathbf{x}^*)$ for a point $\mathbf{x}^* \in \Delta_3 \subset \Delta^3,$

proved in Lemma 2.1 in § 2.1, where $\Delta_3 = \{ \mathbf{x} \in \Delta^3 : f_1^1(\mathbf{x}) = f_2^1(\mathbf{x}) \}$. The statement $\mathbf{x}^* \in \Delta_3$ is deduced from the observation that neither f_1^1 nor f_2^1 has a local minimum in Δ^3 . The number $\log 3$ is obtained by evaluating f_1^1 at the point $\mathbf{x}^*$. The coordinates of $\mathbf{x}^*$ are calculated using the fact that the determinants of the quadratic equations derived from the equality $f_1^1(\mathbf{x}) = f_2^1(\mathbf{x})$ under different parameterizations of Δ^3 is 0.

In Section 3, we give the definition of a decomposition of Γ for a given finite set of isometries $\Psi^* \subset \Gamma$. In particular, we introduce the decomposition

$$\Gamma_{\mathcal{D}^\dagger} = \{1\} \cup \Psi_r^\dagger \cup \bigcup_{\psi \in \Psi^\dagger} J_\psi$$

of Γ which concentrates on the isometries in $\Gamma_\dagger$ where $\Psi^\dagger$ denotes the set of isometries $\{\xi\eta, \xi^2, \xi\eta^{-1}, \eta, \xi^{-1}, \eta^{-1}\xi^{-1}, \eta^{-2}, \eta^{-1}\xi\}$. We prove Lemma 3.3 in § 3.2 which is the analog of Lemma 1.4 for the decomposition $\Gamma_{\mathcal{D}^\dagger}$. Using Lemma 3.3 and Lemma 1.4, we determine the displacement functions for the displacements under isometries given in $\Gamma_\dagger$ in § 3.3. There are 18 displacement functions $\{f_i^\dagger\}_{i=1}^8, \{g_j^\dagger\}_{j=1}^6$ and $\{h_k^\dagger\}_{k=1}^4$ for the decomposition $\Gamma_{\mathcal{D}^\dagger}$. It is possible to show that only the first eight $f_1^\dagger, f_2^\dagger, \dots, f_8^\dagger$ of these functions are significant to find a lower bound for the maximum of the displacements $\text{dist}(z_0, \xi \cdot z_0), \text{dist}(z_0, \eta \cdot z_0)$ and $\text{dist}(z_0, \xi\eta \cdot z_0)$. Let us say $I^\dagger = \{i \in \mathbb{Z} : 1 \leq i \leq 8\}$. The calculation of the infimum of the maximum of the functions $\{f_j^\dagger\}_{j \in I^\dagger}$ follows from the statements

- C. $\inf_{\mathbf{x} \in \Delta^7} \{ \max (f_1^\dagger(\mathbf{x}), \dots, f_8^\dagger(\mathbf{x})) \} = \min_{\mathbf{x} \in \Delta^7} \{ \max (f_1^\dagger(\mathbf{x}), \dots, f_8^\dagger(\mathbf{x})) \},$
- D. $\min_{\mathbf{x} \in \Delta^7} \{ \max (f_1^\dagger(\mathbf{x}), f_2^\dagger(\mathbf{x}), \dots, f_8^\dagger(\mathbf{x})) \} = f_1^\dagger(\mathbf{x}^*)$ for $\mathbf{x}^* \in \Delta_7 \subset \Delta^7,$

proved in Lemma 4.14 in § 4.1 where $\Delta_7 = \{ \mathbf{x} \in \Delta^7 : f_j^\dagger(\mathbf{x}) = f_k^\dagger(\mathbf{x}), j, k \in I^\dagger \}$. Although the proof of the observation that $\mathbf{x}^* \in \Delta_7 \subset \Delta^7$ also uses the fact that none of the functions $\{f_j^\dagger\}_{j \in I^\dagger}$ under any parametrization of Δ^7 has a local minimum, it is much more involved. It requires a case by case study of the values of the functions $\{f_j^\dagger\}_{j \in I^\dagger}$ at the point $\mathbf{x}^*$. Let us say $I_1^\dagger = \{1, 2, 3\}$, $I_2^\dagger = \{4, 5, 6\}$, $I_3^\dagger = \{7, 8\}$, and $\mathcal{I}_i^\dagger = \bigcup_{j \neq i} I_j^\dagger$.

In §§ 4.1.1-§§ 4.1.4, we use the first order partial derivatives of the functions $\{f_j^\dagger\}_{j \in I^\dagger}$ with different parameterizations of the simplex Δ^7 to show that there are certain

direction vectors in $\mathbb{R}^7$ so that moving along these vectors reduces the number of possible cases for the values of the functions $\{f_j^\dagger\}_{j \in I^\dagger}$ at the point $\mathbf{x}^*$ to only five main cases:

- I. $f_j^\dagger(\mathbf{x}^*) < \alpha_*$ for $j \in \mathcal{I}_1^\dagger$ and $f_j^\dagger(\mathbf{x}^*) = \alpha_*$ for $j \in I_1^\dagger$,
- II. $f_j^\dagger(\mathbf{x}^*) < \alpha_*$ for $j \in \mathcal{I}_3^\dagger$ and $f_j^\dagger(\mathbf{x}^*) = \alpha_*$ for $j \in I_3^\dagger$,
- III. $f_j^\dagger(\mathbf{x}^*) = \alpha_*$ for $j \in \mathcal{I}_3^\dagger$ and $f_j^\dagger(\mathbf{x}^*) < \alpha_*$ for $j \in I_3^\dagger$,
- IV. $f_j^\dagger(\mathbf{x}^*) = \alpha_*$ for $j \in \mathcal{I}_2^\dagger$ and $f_j^\dagger(\mathbf{x}^*) < \alpha_*$ for $j \in I_2^\dagger$,
- V. $f_j^\dagger(\mathbf{x}^*) = \alpha_*$ for every $j \in I^\dagger$,

where $\alpha_* = \min_{\mathbf{x} \in \Delta^7} \left\{ \max \left(f_1^\dagger(\mathbf{x}), f_2^\dagger(\mathbf{x}), \dots, f_8^\dagger(\mathbf{x}) \right) \right\}$. In each of the cases I, II, III and IV, we prove in §§ 4.1.6 that there exists a piece of curve in Δ^7 passing through the point $\mathbf{x}^*$ such that moving along this curve produces a point in Δ^7 at which a smaller minimum for the maximum of the functions $\{f_j^\dagger\}_{j \in I^\dagger}$ is attained. This leaves only the case $\mathbf{x}^* \in \Delta_7$ which suggest a method to find the coordinates of the point $\mathbf{x}^*$ and hence the number α_* (see §§ 4.1.6). In § 4.2, we introduce Proposition 4.1 the analog of Proposition 1.5 for the displacements $\text{dist}(z_0, \xi \cdot z_0)$, $\text{dist}(z_0, \eta \cdot z_0)$ and $\text{dist}(z_0, \xi\eta \cdot z_0)$ to complete the case $\Lambda_\Gamma = S_\infty$.

In Section 5, we show that the function $f_{z_0}^\dagger: \mathfrak{D} \rightarrow \mathbb{R}$ defined as follows $(\xi', \eta') \mapsto \max_{\gamma \in \Gamma'_\dagger} \{\text{dist}(z_0, \gamma \cdot z_0)\}$ in the proof of Theorem 5.1 has no local minimum on $\mathfrak{G}\mathfrak{F}$ to complete the case $\Lambda_\Gamma \neq S_\infty$ where $\Gamma'_\dagger = \{\pi(\gamma): \gamma \in \Gamma_\dagger\}$.

2 The log 3 theorem revisited

In this section, we present the calculations required to obtain the number $\log 3$ from the view point we proposed in § 1.4 for the case $\Lambda_\Gamma = S_\infty$.

2.1 Displacement functions for the paradoxical decomposition $\Gamma_{\mathcal{D}^1}$ of Γ

Suppose that Γ , ξ , and η satisfy the hypotheses of the Proposition 1.5. Let z_0 be a given point in $\mathbb{H}^3$. Since Γ and $\Xi = \{\xi, \eta\}$ satisfy the hypotheses of Lemma 1.4, there exists a number $D \in [0, 2]$, a conformal density $\mathcal{M}_{\Gamma, z_0}$ and a family of Borel measures $(\nu_\psi)_{\psi \in \Psi^1}$ where $\Psi^1 = \{\xi, \xi^{-1}, \eta, \eta^{-1}\}$ for which conclusions (1), (2) and, (3) of Lemma 1.4 hold. But the hypotheses of Proposition 1.5 imply that $\mathcal{M}_{\Gamma, z_0}$ is a constant multiple of the area density $\mathcal{A}$. Therefore, we get $D = 2$. Since μ_{z_0} has total

mass 1 by Lemma 1.4(1), we have $\mathcal{M}_{\Gamma, z_0} = \mathcal{A}$. We know that $\nu_\psi(S_\infty) = \int d\nu_\psi$ for every $\psi \in \{\xi, \xi^{-1}, \eta, \eta^{-1}\}$. Let us call

$$\int d\nu_{\xi^{-1}} = m_1, \int d\nu_\xi = m_4, \int d\nu_{\eta^{-1}} = m_3, \int d\nu_\eta = m_2.$$

We also know that $\nu_\xi(S_\infty) + \nu_{\xi^{-1}}(S_\infty) + \nu_{\eta^{-1}}(S_\infty) + \nu_\eta(S_\infty) = 1$ by Lemma 1.4(1) and (2). We claim that $0 < \nu_\psi(S_\infty) < 1$ for every $\psi \in \Psi^1$. Let us assume on the contrary that $\nu_\psi(S_\infty) = 0$. Then we have $\nu_{\psi^{-1}}(S_\infty) = 0$ by Lemma 1.4(4). Since we assume that $\nu_\psi(S_\infty) = 0$, we find that $\int \lambda_{\psi^{-1}, z_0}^2 d\nu_\psi = 0$. By Lemma 1.4(3), we know that $1 - \nu_{\psi^{-1}}(S_\infty) = 0$ or $\nu_{\psi^{-1}}(S_\infty) = 1$, a contradiction. Let us assume that $\nu_\psi(S_\infty) = 1$ for some $\psi \in \Psi^1$. Then we get $\nu_{\psi^{-1}}(S_\infty) = 0$ which gives a contradiction. Hence, we find that $0 < \nu_\psi(S_\infty) < 1$ for every $\psi \in \Psi^1$, or equivalently, $0 < m_i < 1$ for every $i = 1, 2, 3, 4$. In particular, we derive that $0 < \int \lambda_{\psi, z_0}^2 d\nu_{\psi^{-1}} = 1 - \int d\nu_\psi < 1$ for every $\psi \in \Psi^1$. Therefore, we can apply Lemma 1.1 for every $\psi \in \Psi^1$.

Let $\gamma = \xi$, $\nu = \nu_{\xi^{-1}}$, $a = \nu_{\xi^{-1}}(S_\infty)$, and $b = \int \lambda_{\xi, z_0}^2 d\nu_{\xi^{-1}}$. By the conclusion of Lemma 1.1 and Lemma 1.4(3), we obtain

$$(9) \quad e^{2\text{dist}(z_0, \xi \cdot z_0)} \geq \frac{(1 - \int d\nu_\xi)(1 - \int d\nu_{\xi^{-1}})}{\int d\nu_\xi \cdot \int d\nu_{\xi^{-1}}} = \frac{(1 - m_4)(1 - m_1)}{m_4 m_1}.$$

Therefore, the expression in (9) gives a lower bound for $\text{dist}(z_0, \xi \cdot z_0)$. Let us replace the constants m_1, m_2, m_3 and m_4 with the variables x_1, x_2, x_3 and x_4 respectively. Then the displacement function for the displacement $\text{dist}(z_0, \xi \cdot z_0)$ is

$$f_1^1(x_1, x_2, x_3, x_4) = \frac{1 - x_4}{x_4} \cdot \frac{1 - x_1}{x_1}.$$

Let $\gamma = \xi^{-1}$, $\nu = \nu_\xi$, $a = \nu_\xi(S_\infty)$, and $b = \int \lambda_{\xi^{-1}, z_0}^2 d\nu_\xi$. By the conclusion of Lemma 1.1 and Lemma 1.4(3), we calculate that

$$(10) \quad e^{2\text{dist}(z_0, \xi^{-1} \cdot z_0)} \geq \frac{(1 - \int d\nu_{\xi^{-1}})(1 - \int d\nu_\xi)}{\int d\nu_{\xi^{-1}} \cdot \int d\nu_\xi} = \frac{(1 - m_1)(1 - m_4)}{m_1 m_4}.$$

Therefore, the expression in (10) is a lower bound for $\text{dist}(z_0, \xi^{-1} \cdot z_0)$. In other words, the displacement function for the displacement $\text{dist}(z_0, \xi^{-1} \cdot z_0)$ is

$$f_4^1(x_1, x_2, x_3, x_4) = \frac{1 - x_1}{x_1} \cdot \frac{1 - x_4}{x_4}.$$

Let $\gamma = \eta$, $\nu = \nu_{\eta^{-1}}$, $a = \nu_{\eta^{-1}}(S_\infty)$, and $b = \int \lambda_{\eta, z_0}^2 d\nu_{\eta^{-1}}$. By the conclusion of Lemma 1.1 and Lemma 1.4(3), we calculate that

$$(11) \quad e^{2\text{dist}(z_0, \eta \cdot z_0)} \geq \frac{(1 - \int d\nu_\eta)(1 - \int d\nu_{\eta^{-1}})}{\int d\nu_\eta \cdot \int d\nu_{\eta^{-1}}} = \frac{(1 - m_2)(1 - m_3)}{m_2 m_3}.$$

Therefore, the expression in (11) gives a lower bound for $\text{dist}(z_0, \eta \cdot z_0)$. In this case, the displacement function for the displacement $\text{dist}(z_0, \eta \cdot z_0)$ is

$$f_3^1(x_1, x_2, x_3, x_4) = \frac{1 - x_2}{x_2} \cdot \frac{1 - x_3}{x_3}.$$

Similarly, let $\gamma = \eta^{-1}$, $\nu = \nu_\eta$, $a = \nu_\eta(S_\infty)$, and $b = \int \lambda_{\eta^{-1}, z_0}^2 d\nu_\eta$ and apply Lemma 1.1. Then by Lemma 1.4(3), we see that

$$(12) \quad e^{2\text{dist}(z_0, \eta^{-1} \cdot z_0)} \geq \frac{(1 - \int d\nu_{\eta^{-1}})(1 - \int d\nu_\eta)}{\int d\nu_{\eta^{-1}} \cdot \int d\nu_\eta} = \frac{(1 - m_3)(1 - m_2)}{m_3 m_2}.$$

Therefore, the expression in (12) is a lower bound for $\text{dist}(z_0, \eta \cdot z_0)$. Finally, the displacement function for the displacement $\text{dist}(z_0, \eta \cdot z_0)$ is

$$f_2^1(x_1, x_2, x_3, x_4) = \frac{1 - x_3}{x_3} \cdot \frac{1 - x_2}{x_2}.$$

The expressions on the right-hand side of the equalities (9), (10), (11), and (12) can be considered as the values of the displacement functions f_1^1 , f_2^1 , f_3^1 and f_4^1 at the point $\mathbf{m}_1 = (m_1, m_2, m_3, m_4) \in \Delta^3$ where Δ^3 is the 3-dimensional unit simplex $\{(x_1, x_2, x_3, x_4) \in \mathbb{R}_+^4 : x_1 + x_2 + x_3 + x_4 = 1\}$. In order to calculate a lower bound for the $\max(\text{dist}(z_0, \xi \cdot z_0), \text{dist}(z_0, \eta \cdot z_0))$ we need to calculate the infimum of maximum of the functions f_1^1 , f_2^1 , f_3^1 and f_4^1 over the simplex Δ^3 . Points of Δ^3 will be written in bold fonts, e.g., $\mathbf{x} = (x_1, x_2, x_3, x_4)$. Let $\mathcal{F}^1 = \{f_1^1, f_2^1, f_3^1, f_4^1\}$. Let us define the following continuous function

$$\begin{aligned} G^1 : \Delta^3 &\rightarrow \mathbb{R} \\ \mathbf{x} &\mapsto \max\{f(\mathbf{x}) : f \in \mathcal{F}^1\}. \end{aligned}$$

We aim to find $\inf_{\mathbf{x} \in \Delta^3} G^1(\mathbf{x})$. We will need different parametrization of Δ^3 . Let $\Delta^{3,s}$ be the set of points $(x_i, x_j, x_k) \in \mathbb{R}^3$ with the ordered $x_i x_j x_k$ -coordinate system so that $0 < x_i < 1$, $0 < x_j < 1 - x_i$ and $0 < x_k < 1 - x_i - x_j$ where $i < j < k$ and $i, j, k \in J - \{s\} = \{1, 2, 3, 4\} - \{s\}$. For each $s \in J$, let $\mathbf{p}_s^3 : \Delta^{3,s} \rightarrow \Delta^3$ denote the parametrization of Δ^3 where $x_s = 1 - \sum_{m \in J - \{s\}} x_m$. Then we have the following commuting diagrams

$$\begin{array}{ccc} \Delta^{3,s} & & \\ \mathbf{p}_s^3 \downarrow & \searrow f_j^{1,s} & \\ \Delta^3 & \xrightarrow{f_j^1} & \mathbb{R} \end{array}$$

for every $j, s \in J$. We shall use the notation $f_i^{1,s}$ to denote the formulas of the displacement functions f_i^1 for $i \in J$ under the parametrization $\mathbf{p}_s^3$ for $s \in J$. A generic

point in $\Delta^{3,s}$ will be written as $\mathbf{x}_s = (\mathbf{p}_s^3)^{-1}(\mathbf{x})$, e.g., $\mathbf{x}_2 = (x_1, x_3, x_4) \in \Delta^{3,2}$. Let us introduce the following index sets: $I_i^1 = \{i\}$ and $J_i^1 = J - \{i\}$ for $i \in J$. Also let $\mathcal{I}^1 = \{I_i^1 : i \in J\}$ and $\mathcal{J}^1 = \{J_i^1 : i \in J\}$.

2.2 Infimum of the maximum of the functions in $\mathcal{F}^1$ over Δ^3

Before we calculate $\inf_{\mathbf{x} \in \Delta^3} G^1(\mathbf{x})$, we shall develop some notation to shorten the equations we need to study in this section. Let $\mathbf{x} = (x_1, x_2, x_3, x_4)$ be a point and $\mathbf{x}^* = (x_1^*, x_2^*, x_3^*, x_4^*)$ and $\mathbf{x}^\circ = (x_1^\circ, x_2^\circ, x_3^\circ, x_4^\circ)$ denote two fixed points in Δ^3 . Introduce the functions $\sigma: (0, 1) \rightarrow \mathbb{R}$, $\tau: (0, 1) \rightarrow \mathbb{R}$, $\rho_k, \Sigma_k: \Delta^3 \rightarrow (0, 1)$ for each index set $I_k^1 \in \mathcal{I}^1$, and $\sigma_k: \Delta^3 \rightarrow (0, 1)$ for each $k \in J$ where

$$\sigma(x) = \frac{1-x}{x}, \tau(x) = \frac{1}{x^2}, \Sigma_k(\mathbf{x}) = \sum_{i \in I_k^1} x_i, \rho_k(\mathbf{x}) = \sigma(\Sigma_k(\mathbf{x})) \text{ and } \sigma_k(\mathbf{x}) = \sigma(x_k).$$

We have $\sigma(x) > 0$, $\tau(x) > 0$, $\sigma'(x) = -\tau(x) < 0$, and $(1/\sigma(x))' = \tau(1-x) > 0$ for every $x \in (0, 1)$.

For each parametrization $\mathbf{p}_s^3$, let us define the functions $\Sigma_i, \Sigma_{i,j}, \Sigma_{i,j,k}: \Delta^{3,s} \rightarrow (0, 1)$ for each index set $I_i^1, I_j^1, I_k^1 \in \mathcal{I}^1 - \{I_s^1\}$ so that

$$\Sigma_i(\mathbf{x}_s) = \sum_{m \in I_i^1} x_m, \Sigma_{i,j}(\mathbf{x}_s) = \sum_{m \in I_i^1 \cup I_j^1} x_m, \text{ and } \Sigma_{i,j,k}(\mathbf{x}_s) = \sum_{m \in I_i^1 \cup I_j^1 \cup I_k^1} x_m.$$

Let $\Sigma_i^* = \Sigma_i(\mathbf{x}_s^*)$, $\Sigma_i^\circ = \Sigma_i(\mathbf{x}_s^\circ)$, $\Sigma_{i,j}^* = \Sigma_{i,j}(\mathbf{x}_s^*)$, $\Sigma_{i,j}^\circ = \Sigma_{i,j}(\mathbf{x}_s^\circ)$, $\Sigma_{i,j,k}^* = \Sigma_{i,j,k}(\mathbf{x}_s^*)$, $\Sigma_{i,j,k}^\circ = \Sigma_{i,j,k}(\mathbf{x}_s^\circ)$ where $\mathbf{x}_s^* = (\mathbf{p}_s^3)^{-1}(\mathbf{x}^*)$ and $\mathbf{x}_s^\circ = (\mathbf{p}_s^3)^{-1}(\mathbf{x}^\circ)$. Let us also define $\Sigma^i: \Delta^{3,s} \rightarrow (0, 1)$ for each index set J_i^1 for every $i \in J$ such that

$$\Sigma^i(\mathbf{x}_s) = \sum_{m \in J_i^1} x_m.$$

Let us say $\Sigma_i^* = \Sigma^i(\mathbf{x}_s^*)$, $\Sigma_i^\circ = \Sigma^i(\mathbf{x}_s^\circ)$. Also define $\rho_i, \rho^{i,j,k}: \Delta^{3,s} \rightarrow (0, 1)$ for each index set $I_i^1, I_j^1, I_k^1 \in \mathcal{I}^1 - \{I_s^1\}$ where $\rho_i(\mathbf{x}_s) = \sigma(\Sigma_i(\mathbf{x}_s))$ and $\rho^{i,j,k}(\mathbf{x}_s) = 1/\sigma(\Sigma_{i,j,k}(\mathbf{x}_s))$. Let us say $\rho_i^* = \rho_i(\mathbf{x}_s^*)$, $\rho_i^\circ = \rho_i(\mathbf{x}_s^\circ)$, $\rho_{*}^{i,j,k} = \rho^{i,j,k}(\mathbf{x}_s^*)$ and $\rho_{\circ}^{i,j,k} = \rho^{i,j,k}(\mathbf{x}_s^\circ)$. Let $\sigma_i: \Delta^{3,s} \rightarrow (0, 1)$ be the function for $i \in J$ defined by $\sigma_i(\mathbf{x}_s) = \sigma(x_i)$ such that $\sigma_i^* = \sigma_i(\mathbf{x}_s^*)$ and $\sigma_i^\circ = \sigma_i(\mathbf{x}_s^\circ)$. Let us also define $\sigma^i: \Delta^{3,s} \rightarrow \mathbb{R}$ for the index sets $J_i^1 \in \mathcal{J}^1$ so that $\sigma^i(\mathbf{x}_s) = 1/\sigma(\Sigma^i(\mathbf{x}_s))$ where $\sigma_{\circ}^i = \sigma^i(\mathbf{x}_s^\circ)$ and $\sigma_{*}^i = \sigma^i(\mathbf{x}_s^*)$. Let us introduce the functions $\tau_i, \tau^i: \Delta^{3,s} \rightarrow \mathbb{R}$ for every $i \in J$ and for every $J_i^1 \in \mathcal{J}^1$ defined by $\tau_i(\mathbf{x}_s) = \tau(x_i)$ and $\tau^i(\mathbf{x}_s) = \tau(1 - \Sigma^i(\mathbf{x}_s))$. Similarly, for each index set $I_i^1, I_j^1, I_k^1 \in \mathcal{I}^1 - \{I_s^1\}$, let $\varrho_i, \varrho^{i,j,k}: \Delta^{3,s} \rightarrow \mathbb{R}$ be the functions defined by $\varrho_i(\mathbf{x}_s) = \tau(\Sigma_i(\mathbf{x}_s))$ and $\varrho^{i,j,k}(\mathbf{x}_s) = \tau(1 - \Sigma_{i,j,k}(\mathbf{x}_s))$. Note that we have the partial derivatives

$$(\sigma_i)_m = \begin{cases} -\tau_i < 0 & \text{if } m = i, \\ 0 & \text{otherwise,} \end{cases} \quad (\rho_i)_m = \begin{cases} -\varrho_i < 0 & \text{if } m \in I_i^1, \\ 0 & \text{otherwise,} \end{cases}$$

$$(\sigma^i)_m = \begin{cases} \tau^i > 0 & \text{if } m \in J_i^1, \\ 0 & \text{otherwise,} \end{cases} \quad (\rho^{i,j,k})_m = \begin{cases} \varrho^{i,j,k} > 0 & \text{if } m \in I_i^1 \cup I_j^1 \cup I_k^1, \\ 0 & \text{otherwise.} \end{cases}$$

We observe that $\inf_{\mathbf{x} \in \Delta^3} G^1(\mathbf{x}) = \inf_{\mathbf{x} \in \Delta^3} \{\max(f_1^1(\mathbf{x}), f_2^1(\mathbf{x}))\}$ because $f_1^1(\mathbf{x}) = f_4^1(\mathbf{x})$ and $f_2^1(\mathbf{x}) = f_3^1(\mathbf{x})$ for every $\mathbf{x} \in \Delta^3$. Therefore, it is enough to prove the following:

Lemma 2.1 *If $f_1^1(\mathbf{x}) = \rho_4(\mathbf{x})\sigma_1(\mathbf{x})$ and $f_2^1(\mathbf{x}) = \rho_3(\mathbf{x})\sigma_2(\mathbf{x})$ are the displacement functions on Δ^3 defined in §2.1, then we have $\inf_{\mathbf{x} \in \Delta^3} \{\max(f_1^1(\mathbf{x}), f_2^1(\mathbf{x}))\} = 9$.*

Proof Define the continuous function $F^1: \Delta^3 \rightarrow \mathbb{R}$ so that $\mathbf{x} \mapsto \max(f_1^1(\mathbf{x}), f_2^1(\mathbf{x}))$. Note that the inequalities $f_1^1(\mathbf{x}) > 1$ and $f_2^1(\mathbf{x}) > 1$ hold for every $\mathbf{x} \in \Delta^3$. Because if we have $f_1^1(\mathbf{x}) = 1$ for some point $\mathbf{x} \in \Delta^3$, then we get $x_1 + x_4 = 1$. This implies that $x_2 = 0$ and $x_3 = 0$. But the point $(x_1, 0, 0, x_4)$ is not in Δ^3 . Similarly, if we have $f_2^1(\mathbf{x}) = 1$ for some point $\mathbf{x} \in \Delta^3$, then we get $x_2 + x_3 = 1$. This implies that $x_1 = 0$ and $x_4 = 0$. But the point $(0, x_2, x_3, 0)$ is not in Δ^3 . Therefore, we have $\inf_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) \geq 1$. We need to show that $\inf_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = 9$. First, we prove that

$$\inf_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}).$$

Let $\Delta_n = \{(x_1, x_2, x_3, x_4) \in \Delta^3: 1/n \leq x_i \leq 1 - 1/(2n) \text{ for } i = 1, 2, 3, 4\}$ for $n \geq 2$, then F^1 has an absolute minimum α_n at a point $\mathbf{x}_n$ in Δ_n since Δ_n is compact. The sequence (α_n) is decreasing since we have $\Delta_n \subset \Delta_{n+1}$. Let us consider the sequence $(\mathbf{x}_n)$. We claim that the sequence $(\mathbf{x}_n)$ cannot have a limit point on the boundary of Δ^3 . Let us assume to the contrary that $\mathbf{x}_n \rightarrow \mathbf{x}_0 \in \partial\Delta^3$ as $n \rightarrow \infty$. If b_1, b_2, b_3, b_4 denote the coordinates of the point $\mathbf{x}_0$, then $b_i = 0$ for some $i = 1, 2, 3, 4$. Suppose, for example, that $b_1 = 0$. This implies that $b_2 = 1$ since otherwise F^1 would tend to infinity through the sequence $\mathbf{x}_n$. But it is supposed to be tending to an infimum. Therefore, we must have $b_3 = 0$ and $b_4 = 0$. If we repeat this argument for $b_3 = 0$, we see that $b_4 = 1$ which gives a contradiction.

We claim that there exists a positive integer n_0 so that $\mathbf{x}_n = \mathbf{x}_{n+1}$ for every $n \geq n_0$. Let us assume otherwise that we have a subsequence $(\mathbf{x}_{n_j})$ of the sequence $(\mathbf{x}_n)$ so that $\mathbf{x}_{n_j} \in \Delta_{n_{j+1}} - \Delta_{n_j}$ for every integer $j > 0$. Since we have $\bigcup_{n=2}^{\infty} \Delta_n = \Delta^3$, we get $F^1(\mathbf{x}_{n_j}) \rightarrow \infty$ as $j \rightarrow \infty$. But this contradicts with the fact that (α_n) is a decreasing sequence. Therefore, we have $\mathbf{x}_n = \mathbf{x}_{n+1}$ for every $n \geq n_0$. In other words, the absolute minimum $F^1(\mathbf{x}_{n_0})$ of F^1 exists on Δ^3 . Then, we obviously have $\inf_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x})$. Thus, we need to calculate the $\min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x})$. To this purpose, we find a point in Δ^3 where the minimum is attained. Let $\mathbf{x}^* = (x_1^*, x_2^*, x_3^*, x_4^*) \in \Delta^3$ be a point so that $F^1(\mathbf{x}^*) = \min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x})$ and $\Delta_3 = \{\mathbf{x} \in \Delta^3: f_1^1(\mathbf{x}) = f_2^1(\mathbf{x})\}$. We claim that $\mathbf{x}^* \in \Delta_3$.

Let us assume otherwise that $f_1^1(\mathbf{x}^*) \neq f_2^1(\mathbf{x}^*)$. For any parametrization $\mathbf{p}_s^3$, we have the following commuting diagram

$$\begin{array}{ccc} \Delta^{3,s} & & \\ \mathbf{p}_s^3 \downarrow & \searrow F^{1,s} & \\ \Delta^3 & \xrightarrow{F^1} & \mathbb{R} \end{array}$$

where $F^{1,s}(\mathbf{x}_s) = \max_{\mathbf{x}_s \in \Delta^{3,s}} \{f_1^{1,s}(\mathbf{x}_s), f_2^{1,s}(\mathbf{x}_s)\}$. Using the diagram above, we derive that $F^1(\mathbf{x}^*) = F^{1,s}(\mathbf{x}_s^*)$ such that $\mathbf{x}_s^* = (\mathbf{p}_s^3)^{-1}(\mathbf{x}^*)$. We use the parametrization $\mathbf{p}_4^3$ of Δ^3 . We have either $f_1^{1,4}(\mathbf{x}_4^*) > f_2^{1,4}(\mathbf{x}_4^*)$ or $f_1^{1,4}(\mathbf{x}_4^*) < f_2^{1,4}(\mathbf{x}_4^*)$. If we assume that $f_1^{1,4}(\mathbf{x}_4^*) > f_2^{1,4}(\mathbf{x}_4^*)$, then there exists a neighborhood U of $\mathbf{x}_4^*$ in $\Delta^{3,4}$ so that $f_1^{1,4}(\mathbf{x}_4) > f_2^{1,4}(\mathbf{x}_4)$ for every $\mathbf{x}_4 \in U$ since $f_1^{1,4}$ is continuous on $\Delta^{3,4}$. Hence, we get $F^{1,4}(\mathbf{x}_4) = f_1^{1,4}(\mathbf{x}_4)$ for every $\mathbf{x}_4 \in U$. Since we have $f_1^{1,4}(\mathbf{x}_4^*) = F^{1,4}(\mathbf{x}_4^*)$, the functions $f_1^{1,4}$ has a local minimum at $\mathbf{x}_4^* \in U$. We know that $f_1^{1,4} = \rho^{1,2,3}\sigma_1$ such that $(f_1^{1,4})_1 = \varrho^{1,2,3}\sigma_1 - \rho^{1,2,3}\tau_1$, $(f_1^{1,4})_2 = \varrho^{1,2,3}\sigma_1$ and $(f_1^{1,4})_3 = \varrho^{1,2,3}\sigma_1$. We have $(f_1^{1,4})_2 > 0$ for every $\mathbf{x}_4 \in \Delta^{3,4}$. Therefore, $f_1^{1,4}$ has no critical point in $\Delta^{3,4}$. Since $f_1^{1,4}$ is differentiable everywhere in $\Delta^{3,4}$ and has no local minimum, we obtain a contradiction.

Let us assume that $f_1^{1,4}(\mathbf{x}_4^*) < f_2^{1,4}(\mathbf{x}_4^*)$. Since $f_2^{1,4}$ is a continuous function on $\Delta^{3,4}$, we get $f_1^{1,4}(\mathbf{x}_4) < f_2^{1,4}(\mathbf{x}_4)$ for every $\mathbf{x}_4$ in some neighborhood V of $\mathbf{x}_4^*$ where we have $F^{1,4}(\mathbf{x}_4) = f_2^{1,4}(\mathbf{x}_4)$ for every $\mathbf{x}_4 \in V$. Then, $f_2^{1,4}$ has a local minimum at $\mathbf{x}_4^* \in V$. We know that $f_2^{1,4} = \rho_3\sigma_2$ so that $(f_2^{1,4})_1 = 0$, $(f_2^{1,4})_2 = -\rho_3\tau^1$, $(f_2^{1,4})_3 = -\varrho_3\sigma_2$. We see that $(f_2^{1,4})_2 < 0$ for every $\mathbf{x}_4 \in \Delta^{3,4}$. Therefore, $f_2^{1,4}$ has no critical point in $\Delta^{3,4}$. Since $f_2^{1,4}$ is also differentiable everywhere in $\Delta^{3,4}$ and has no local minimum, we obtain another contradiction. Hence, we conclude that $\mathbf{x}^* \in \Delta_3$.

Using the parametrization $\mathbf{p}_4^3$ of Δ^3 , we consider the equation $f_1^{1,4}(\mathbf{x}_4^*) = f_2^{1,4}(\mathbf{x}_4^*)$ or $\rho_*^{1,2,3}\sigma_1^* = \rho_3^*\sigma_2^*$. We find that

$$\frac{x_1^* + x_2^* + x_3^*}{1 - x_1^* - x_2^* - x_3^*} \cdot \frac{1 - x_1^*}{x_1^*} = \frac{1 - x_3^*}{x_3^*} \cdot \frac{1 - x_2^*}{x_2^*},$$

or $(1 - \Sigma_{2,3}^*)(x_1^*)^2 - (1 - \Sigma_{2,3}^*)^2 x_1^* + x_2^* x_3^* \Sigma_{2,3}^* = 0$. In other words, we can consider x_1^* as a solution of the equation $q_1^1(x) = 0$ where $q_1^1(x) = (1 - \Sigma_{2,3}^*)x^2 - (1 - \Sigma_{2,3}^*)^2 x + x_2^* x_3^* \Sigma_{2,3}^*$. Let $D_1^1(\mathbf{x}_4) = D_1^1(x_1, x_2, x_3)$ where $D_1^1(x_1, x_2, x_3) = (1 - \Sigma_{2,3}^*)^4 - 4x_2 x_3 \Sigma_{2,3}^* (1 - \Sigma_{2,3}^*)$. Since $D_1(\mathbf{x}_4^*)$ corresponds to the discriminant of $q_1^1(x)$, we have $D_1^1(\mathbf{x}_4^*) \geq 0$. In particular, we get the expression

$$x_1^* = \frac{(1 - x_2^* - x_3^*)^2 \pm \sqrt{D_1^1(\mathbf{x}_4^*)}}{2(1 - x_2^* - x_3^*)}.$$

We claim that $D_1^1(\mathbf{x}_4^*) = 0$. Let us assume otherwise that $D_1^1(\mathbf{x}_4^*) > 0$. Since D_1^1 is continuous, there exists an $\epsilon > 0$ so that $D_1^1(x_1^\circ, x_2^\circ, x_3^\circ) > 0$ for $x_2^\circ = x_2^*$ and $x_3^\circ = x_3^* + \epsilon$. Therefore, if we say $\tilde{q}_1^1(x) = (1 - \Sigma_{2,3}^\circ)x^2 - (1 - \Sigma_{2,3}^\circ)^2x + x_2^\circ x_3^\circ(\Sigma_{2,3}^\circ)$, we see that the quadratic equation $\tilde{q}_1^1(x) = 0$ has two real roots. These roots are given as

$$x_1^\circ = \frac{(1 - x_2^\circ - x_3^\circ)^2 \pm \sqrt{D_1^1(\mathbf{x}_4^\circ)}}{2(1 - x_2^\circ - x_3^\circ)}.$$

In fact, we can choose $\epsilon > 0$ so that the coefficients of $q_1^1(x)$ and $\tilde{q}_1^1(x)$ are sufficiently close to each other to conclude that $0 < x_i^\circ < 1$ for every $i = 1, 2, 3, 4$ where we set $x_4^\circ = 1 - x_1^\circ - x_2^\circ - x_3^\circ$. Therefore, if we consider the point $\mathbf{x}^\circ = (x_1^\circ, x_2^\circ, x_3^\circ, x_4^\circ) \in \Delta^3$, we see that $f_1^1(\mathbf{x}^\circ) = f_2^1(\mathbf{x}^\circ)$. But we obtain $F^1(\mathbf{x}^\circ) = f_2^1(\mathbf{x}^\circ)$ where

$$F^1(\mathbf{x}^\circ) = \sigma(x_3^* + \epsilon)\sigma(x_2^*) < \rho_3^*\sigma_2^* = F^1(\mathbf{x}^*).$$

This contradicts with the minimality of $F^1(\mathbf{x}^*)$. Thus, we get $D_1^1(x_1^*, x_2^*, x_3^*) = 0$. Then, we know that $2x_1^* = 1 - x_2^* - x_3^*$. Since we have $x_4^* = 1 - x_1^* - x_2^* - x_3^*$, we obtain $x_1^* = x_4^*$.

We repeat a similar argument using the parametrization $\mathbf{p}_2^3$. In other words, we consider the equation $f_1^{1,2}(\mathbf{x}_2^*) = f_2^{1,2}(\mathbf{x}_2^*)$ or $\rho_4^*\sigma_1^* = \rho_3^*\sigma_2^{1,3,4}$. We see that

$$\frac{1 - x_1^*}{x_1^*} \cdot \frac{1 - x_4^*}{x_4^*} = \frac{1 - x_3^*}{x_3^*} \cdot \frac{x_1^* + x_3^* + x_4^*}{1 - x_1^* - x_3^* - x_4^*}.$$

or $(1 - \Sigma_{1,4}^*)(x_3^*)^2 - (1 - \Sigma_{1,4}^*)^2x_3^* + x_1^*x_4^*\Sigma_{1,4}^* = 0$. In other words, we can consider x_3^* as a solution of the equation $q_2^1(x) = 0$ where $q_2^1(x) = (1 - \Sigma_{1,4}^*)x^2 - (1 - \Sigma_{1,4}^*)^2x + x_1^*x_4^*\Sigma_{1,4}^*$. Let $D_2^1(\mathbf{x}_2) = D_1^1(x_1, x_3, x_4)$ where $D_2^1(x_1, x_3, x_4) = (1 - \Sigma_{1,4})^4 - 4x_1x_4\Sigma_{1,4}(1 - \Sigma_{1,4})$. Since $D_2(\mathbf{x}_2^*)$ corresponds to the discriminant of $q_2^1(x)$, we have $D_2^1(\mathbf{x}_2^*) \geq 0$. In particular, we get the expression

$$x_3^* = \frac{(1 - x_1^* - x_4^*)^2 \pm \sqrt{D_2^1(\mathbf{x}_2^*)}}{2(1 - x_1^* - x_4^*)}.$$

We claim that $D_2^1(\mathbf{x}_2^*) = 0$. Let us assume otherwise that $D_2^1(\mathbf{x}_2^*) > 0$. Since D_2^1 is continuous, there exists an $\epsilon > 0$ so that $D_2^1(x_1^\circ, x_3^\circ, x_4^\circ) > 0$ for $x_1^\circ = x_1^*$ and $x_4^\circ = x_4^* + \epsilon$. Therefore, if we say $\tilde{q}_2^1(x) = (1 - \Sigma_{1,4}^\circ)x^2 - (1 - \Sigma_{1,4}^\circ)^2x + x_1^\circ x_4^\circ(\Sigma_{1,4}^\circ)$, we see that the quadratic equation $\tilde{q}_2^1(x) = 0$ has two real roots. These roots are given as

$$x_3^\circ = \frac{(1 - x_1^\circ - x_4^\circ)^2 \pm \sqrt{D_2^1(\mathbf{x}_2^\circ)}}{2(1 - x_1^\circ - x_4^\circ)}.$$

In fact, we can choose $\epsilon > 0$ so that the coefficients of $q_2^1(x)$ and $\tilde{q}_2^1(x)$ are sufficiently close to each other to conclude that $0 < x_i^\circ < 1$ for every $i = 1, 2, 3, 4$ where we set

$x_2^\circ = 1 - x_1^\circ - x_3^\circ - x_4^\circ$. Therefore, if we consider the point $\mathbf{x}^\circ = (x_1^\circ, x_2^\circ, x_3^\circ, x_4^\circ) \in \Delta^3$, we see that $f_1^1(\mathbf{x}^\circ) = f_2^1(\mathbf{x}^\circ)$. But we obtain $F^1(\mathbf{x}^\circ) = f_1^1(\mathbf{x}^\circ)$ where

$$F^1(\mathbf{x}^\circ) = \sigma(x_4^* + \epsilon)\sigma(x_1^*) < \rho_4^*\sigma_1^* = F^1(\mathbf{x}^*).$$

This contradicts with the minimality of $F^1(\mathbf{x}^*)$. Thus, we get $D_2^1(x_1^*, x_3^*, x_4^*) = 0$. Then, we know that $2x_3^* = 1 - x_1^* - x_4^*$. Since we have $x_2^* = 1 - x_1^* - x_3^* - x_4^*$, we obtain $x_2^* = x_3^*$.

We know that $\mathbf{x}^* = (x_1^*, x_2^*, x_2^*, x_1^*) \in \Delta_3$. Using the equality $f_1^1(\mathbf{x}^*) = f_2^1(\mathbf{x}^*)$ once more, we calculate that $(\sigma_1^*)^2 = (\sigma_2^*)^2$. Since we know that $x_1^* > 0$ and $x_2^* > 0$, we obtain that $x_1^* = x_2^*$. Hence, we get $\mathbf{x}^* = (x_1^*, x_1^*, x_1^*, x_1^*)$. As a result, x_1^* satisfies the equation $x_1^* + x_1^* + x_1^* + x_1^* = 1$. In other words, we get that $x_1^* = 1/4$. Then we obtain that $\mathbf{x}^* = (1/4, 1/4, 1/4, 1/4)$. Finally, we calculate that $\min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = F^1(\mathbf{x}^*)$ where $F^1(\mathbf{x}^*) = f_1^1(\mathbf{x}^*)$ and $f_1^1(\mathbf{x}^*) = 9$. $\square$

2.3 Proof of Proposition 1.5

Lemma 2.1 suggests a slightly different proof for Proposition 1.5 than the one presented in [9]. This proof is given as follows:

Proof We know that $\text{dist}(z_0, \gamma \cdot z_0) = \text{dist}(z_0, \gamma^{-1} \cdot z_0)$ for every isometry γ . Therefore, it is enough to show that $\max(\text{dist}(z_0, \xi \cdot z_0), \text{dist}(z_0, \eta \cdot z_0)) \geq \log 3$. Let $\mathbf{m}_1 = (m_1, m_2, m_3, m_4) \in \Delta^3$ where m_1, m_2, m_3 and m_4 are the total measures defined in §2.1. By the discussion given in §2.1, we have

$$\max(\text{dist}(z_0, \xi \cdot z_0), \text{dist}(z_0, \eta \cdot z_0)) \geq \frac{1}{2} \log \left(\max \left(\prod_{i=4,1} \frac{1-m_i}{m_i}, \prod_{i=3,2} \frac{1-m_i}{m_i} \right) \right).$$

Using the fact that $(\log t)/2$ is a strictly increasing function, we obtain that

$$\frac{1}{2} \log \left(\max \left(\prod_{i=4,1} \frac{1-m_i}{m_i}, \prod_{i=3,2} \frac{1-m_i}{m_i} \right) \right) \geq \frac{1}{2} \log \left(\inf_{\mathbf{x} \in \Delta^3} \{ \max(f_1^1(\mathbf{x}), f_2^1(\mathbf{x})) \} \right).$$

In Lemma 2.1, we showed that $\inf_{\mathbf{x} \in \Delta^3} \{ \max(f_1^1(\mathbf{x}), f_2^1(\mathbf{x})) \} = 9$. Hence, we get

$$e^{(2 \max \{ \text{dist}(z_0, \xi \cdot z_0), \text{dist}(z_0, \eta \cdot z_0) \})} \geq \inf_{\mathbf{x} \in \Delta^3} \{ \max(f_1^1(\mathbf{x}), f_2^1(\mathbf{x})) \} = 9.$$

$\square$

We would like to emphasize a few key points used in the proof of Lemma 2.1. Remember that f_1^1 and f_2^1 have no local minimum under any parametrization of the simplex Δ^3 . This fact implied that $\mathbf{x}^* \in \Delta_3$. The proof of Lemma 2.1 showed that the main ingredients used to calculate the number $\log 3$ are

- A. $\inf_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x})$,
- B. There exists a point $\mathbf{x}^* \in \Delta_3 \subset \Delta^3$ such that $\min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = f_1^1(\mathbf{x}^*)$.

Lemma 2.1 also establishes that the point $\mathbf{x}^* \in \Delta^3$ is unique. If we assume the uniqueness of the point $\mathbf{x}^*$ a priori together with (A), it is possible to suggest an alternative way of finding the coordinates of the point $\mathbf{x}^*$ which reduces the number of calculations necessary to find the number $\log 3$.

Let $T_1, T_2: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be the maps defined by $(x_1, x_2, x_3, x_4) \mapsto (x_4, x_3, x_2, x_1)$ and $(x_1, x_2, x_3, x_4) \mapsto (x_2, x_1, x_4, x_3)$, respectively. The simplex Δ^3 is preserved by the transformations T_1 and T_2 . We have $f_i^1(T_1(\mathbf{x})) = f_i^1(\mathbf{x})$ for every $\mathbf{x} \in \Delta^3$ for $i = 1, 2$. Let $H_1: \Delta^3 \rightarrow \mathbb{R}$ be the function so that $H_1(\mathbf{x}) = \max\{(f_i^1 \circ T_1)(\mathbf{x}): i = 1, 2\}$. We see that $F^1(\mathbf{x}) = H_1(\mathbf{x})$ for every $\mathbf{x} \in \Delta^3$, $\min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^3} H_1(\mathbf{x})$ and $\{f_i^1(\mathbf{x}^*): i = 1, 2\} = \{f_i^1(T_1(\mathbf{x}^*)): i = 1, 2\}$. Since F^1 takes its minimum value at the point $\mathbf{x}^*$, H_1 takes its minimum value at the point $T_1^{-1}(\mathbf{x}^*)$. We assumed that $\mathbf{x}^*$ is unique. Therefore, we obtain that $T_1^{-1}(\mathbf{x}^*) = \mathbf{x}^*$ which implies that $x_1^* = x_4^*$, and $x_2^* = x_3^*$.

Similarly, we have $f_1^1(T_2(\mathbf{x})) = f_2^1(\mathbf{x})$ and $f_2^1(T_2(\mathbf{x})) = f_1^1(\mathbf{x})$ for every $\mathbf{x} \in \Delta^3$. Let $H_2: \Delta^3 \rightarrow \mathbb{R}$ be the function so that $H_2(\mathbf{x}) = \max\{(f_i^1 \circ T_2)(\mathbf{x}): i = 1, 2\}$. We see that $F^1(\mathbf{x}) = H_2(\mathbf{x})$ for every $\mathbf{x} \in \Delta^3$ and $\min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^3} H_2(\mathbf{x})$. The function H_2 takes its minimum value at the point $T_2^{-1}(\mathbf{x}^*)$ because we have $\{f_i^1(\mathbf{x}^*): i = 1, 2\} = \{f_i^1(T_2(\mathbf{x}^*)): i = 1, 2\}$ and F^1 takes its minimum value at the point $\mathbf{x}^*$. We assumed that $\mathbf{x}^*$ is unique. Hence, we obtain that $T_2^{-1}(\mathbf{x}^*) = \mathbf{x}^*$. This means that $x_1^* = x_2^*$, and $x_3^* = x_4^*$. In other words, we find that $x_i^* = x_j^*$ for every $i, j = 1, 2, 3, 4$. Since we know $\mathbf{x}^* \in \Delta^3$, we see that $4x_i^* = 1$ for every i . Finally, we find that $\mathbf{x}^* = (1/4, 1/4, 1/4, 1/4)$. Then a simple evaluation shows that $\min_{\mathbf{x} \in \Delta^3} F^1(\mathbf{x}) = 9$. In this calculation method, we don't refer to the statement $\mathbf{x}^* \in \Delta_3$ given in (B).

3 Decompositions of 2-generator free groups

Let Γ be a group which is free on a finite generating set Ξ . Every element γ of Γ can be written uniquely as a reduced word $\psi_1 \cdots \psi_m$, where $m \geq 0$, each ψ_i is an element

of $\Xi \cup \Xi^{-1}$, and $\psi_{i+1} \neq \psi_i^{-1}$ for $i = 1, \dots, m-1$. If $k \leq m$ is a positive integer and $\gamma \neq 1$, we shall call $\psi_1 \dots \psi_k$ the *initial word of length k* of γ .

Let Ψ^* be a finite set of words in Γ . For each word $\psi \in \Psi^*$, let J_ψ denote the set of non-trivial elements of Γ that have initial word ψ . Depending on the number of elements in Ξ and lengths of words in Ψ^* there may be a set of words which are not contained in any of J_ψ . Let us call this set the *residue set* of Ψ^* and denote it by Ψ_r^* .

Definition 3.1 For a given pair $\mathcal{D}^* = (\Psi^*, \Psi_r^*)$ of finite, disjoint sets of words Ψ^* and Ψ_r^* in Γ , if $\Gamma = \{1\} \cup \Psi_r^* \cup \bigcup_{\psi \in \Psi^*} J_\psi$, then $\Gamma_{\mathcal{D}^*}$ will be called a *decomposition* of Γ .

Definition 3.2 A decomposition $\Gamma_{\mathcal{D}^*}$ with $\mathcal{D}^* = (\Psi^*, \Psi_r^*)$ is *symmetric* if Ψ^* and Ψ_r^* are preserved by every bijection of $\Xi \cup \Xi^{-1}$.

3.1 Decomposition of $\Gamma = \langle \xi, \eta \rangle$ for the isometries in $\Gamma_\dagger$

We know that $\Gamma = \langle \xi, \eta \rangle$ is a free group on the generating set $\Xi = \{\xi, \eta\}$ ([9], Proposition 9.2). The decomposition $\Gamma_{\mathcal{D}^0}$ where $\mathcal{D}^0 = (\Xi \cup \Xi^{-1}, \emptyset)$ corresponds to the paradoxical decomposition of Γ used in the proof of the log 3 Theorem. In this section, we introduce a decomposition of Γ that contains the isometries in $\Gamma_\dagger = \{\xi, \eta, \xi\eta\}$.

Let $\Psi^\dagger$ be the set $\{\xi\eta, \xi^2, \xi\eta^{-1}, \eta, \xi^{-1}, \eta^{-1}\xi^{-1}, \eta^{-2}, \eta^{-1}\xi\}$. Then the residue set of $\Psi^\dagger$ is $\Psi_r^\dagger = \{\xi, \eta^{-1}\}$. It is straightforward to see that $\Gamma = \{1\} \cup \Psi_r^\dagger \cup \bigcup_{\psi \in \Psi^\dagger} J_\psi$. Clearly, $\Gamma_{\mathcal{D}^\dagger}$ is a decomposition of Γ with $\mathcal{D}^\dagger = (\Psi^\dagger, \Psi_r^\dagger)$. We will concentrate on the decomposition $\Gamma_{\mathcal{D}^\dagger}$ to prove Theorem 5.1. Notice that $\Gamma_{\mathcal{D}^1}$ is a symmetric decomposition. On the other hand, $\Gamma_{\mathcal{D}^\dagger}$ is not symmetric.

3.2 Γ -invariant conformal densities for the decomposition $\Gamma_{\mathcal{D}^\dagger}$

In this section, we prove Lemma 3.3 an analog of Lemma 1.4 for the decomposition $\Gamma_{\mathcal{D}^\dagger}$. Let $J_\Psi := \bigcup_{\psi \in \Psi} J_\psi$ where $\cup$ denotes the disjoint union.

Let $\pi: J \rightarrow \{\xi, \xi^{-1}, \eta, \eta^{-1}\}$ be a bijection where $J = \{1, 2, 3, 4\}$. Then π determines a map $\pi_1: A_1 \rightarrow \Psi^\dagger$ where $A_1 = \{(i, j) \in J \times J: \pi(i)\pi(j) \in \Psi^\dagger\}$. Note that we have $\pi_1(i, j) = \pi(i)\pi(j)$ for every $(i, j) \in A_1$. Let $A_2 = \{(i, j) \in A_1: (\pi_1(i, j))^{-1} \in \Psi^\dagger\}$. For every $(i, j) \in A_2$, we define the terms $\gamma_{i,j} = \pi_1(i, j)$, $\gamma^{i,j} = (\gamma_{i,j})^{-1}$, $\gamma_i = \pi(i)$, and $\Psi_i = \{\pi_1(i, k): (i, k) \in A_1\}$. Let $A_0^\dagger = \{\gamma_{i,j}: (i, j) \in A_2\}$.

Let $A_3 = \{j \in J : \pi(j) \in \Psi^\dagger\}$. Then for every $j \in A_3$, we define $\gamma_j = (\pi(j))^{-1}$, $\gamma^j = \pi(j)$, $\Upsilon_j = \{\pi_1(k, l) : \pi(k) = \gamma_j, (k, l) \in A_1\}$. Let $A_1^\dagger = \{\gamma_j : j \in A_3\}$. For every $(i, j) \in A_4 = A_1 - A_2$, let $\beta_{ij} = (\pi(i))^{-1}$ and $\beta^{ij} = \pi_1(i, j)$, $\beta_j = \pi(j)$, $\Phi_j = \Psi^\dagger - \Psi_j$, and $A_2^\dagger = \{\beta_{ij} : (i, j) \in A_4\}$. For every $(i, j) \in A_2$, let $\zeta_{ij} = \pi_1(i, j)$, $\zeta^{ij} = (\pi(j))^{-1}$, $\zeta_i = \pi(i)$ and $A_3^\dagger = \{\zeta_{ij} : (i, j) \in A_2\}$. Let $A_5 = \{((i, k), (l, j)) \in A_1 \times A_1 : \pi(k)\pi(l) = 1, \pi(i)\pi(j) \neq 1\}$. Let us denote any element $((i, k), (l, j))$ in A_5 by (i, k, j) . Let $\zeta_{i,k,j} = \pi(i)\pi(k)$, $\zeta^{i,k,j} = \pi(k)^{-1}\pi(j)$, and $\Omega_{i,k,j} = \Psi^\dagger - \{\pi_1(i, j)\}$ for every $(i, k, j) \in A_5$. Let $A_4^\dagger = \{\zeta_{i,k,j} : (i, k, j) \in A_5\}$. Let us also define $A_6 = \{(i, j) \in A_1 : \pi(j) \in \Psi_r^\dagger\}$. For every $(i, j) \in A_6$, let $\delta_{ij} = \pi_1(i, j)$ and $\delta^{ij} = (\pi(j))^{-1}$. Let us define $A_5^\dagger = \{\delta_{ij} : (i, j) \in A_6\}$. Then we prove

Lemma 3.3 *Let Γ be a Kleinian group which is free on a generating set $\{\xi, \eta\}$ and $\Gamma_{\mathcal{D}^\dagger}$ be the decomposition of Γ described in §3.1. Let z_0 be any point of $\mathbb{H}^3$. Let γ_{ij} , γ^{ij} , γ_i and Ψ_i for $(i, j) \in A_2$, γ_j , γ^j and Υ_j for $j \in A_3$, β_{ij} , β^{ij} , β_j and Φ_i for $(i, j) \in A_4$, ζ_{ij} , ζ^{ij} , ζ_j for $(i, j) \in A_2$, $\zeta_{i,k,j}$, $\zeta^{i,k,j}$ and $\Omega_{i,k,j}$ for $(i, k, j) \in A_5$, δ_{ij} , δ^{ij} for $(i, j) \in A_6$ be the expressions defined above and*

$$\begin{aligned} s(\gamma_{ij}) &= \gamma^{ij}, s(\gamma_j) = \gamma^j, s(\beta_{ij}) = \beta^{ij}, s(\zeta_{ij}) = \zeta^{ij}, s(\zeta_{i,k,j}) = \zeta^{i,k,j}, \\ s(\delta_{ij}) &= \delta^{ij}, S(\gamma_{ij}) = \{\gamma_i\} \cup \Psi_i, S(\gamma_j) = \{\gamma_j\} \cup \Upsilon_j, S(\beta_{ij}) = \{\beta_j\} \cup \Phi_j, \\ S(\zeta_{ij}) &= \{\zeta_i\} \cup \Phi_i, S(\zeta_{i,k,j}) = \Psi_r^\dagger \cup \Omega_{i,k,j}, S(\delta_{ij}) = \{\delta_{ij}\}. \end{aligned}$$

Then there exists a number $D \in [0, 2]$, a Γ -invariant D -conformal density for $\mathbb{H}^3$ $\mathcal{M}_{\Gamma, z_0} = (\mu_z)$ and a family $(\nu_\psi)_{\psi \in \Psi}$ of Borel measures on S_∞ such that

- (1) $\mu_{z_0}(S_\infty) = 1$; (2) $\mu_{z_0} = \sum_{\psi \in \Psi^\dagger} \nu_\psi$; and (3) for every $\gamma \in A^\dagger = \bigcup_{m=0}^5 A_m^\dagger$

$$\int \lambda_{\gamma, z_0}^D d\mu_{V_{s(\gamma)}} = 1 - \sum_{\psi \in S(\gamma)} \int d\nu_\psi.$$

Proof We know that $\Gamma_{\mathcal{D}^\dagger}$ is a decomposition of Γ . Since Γ is discrete and torsion-free, it acts freely on $\mathbb{H}^3$. Hence the orbit $W = \Gamma \cdot z_0$ is a disjoint union

$$W = \{z_0\} \cup V_0 \cup \bigcup_{\psi \in \Psi^\dagger} V_\psi$$

where $V_0 = \{\gamma \cdot z_0 : \gamma \in \Psi_r^\dagger\}$ and $V_\psi = \{\gamma \cdot z_0 : \gamma \in J_\psi\}$. Note that V_0 is a finite set which is equal to $V_\xi \cup V_{\eta^{-1}}$ where $V_\xi = \{\xi \cdot z_0\}$ and $V_{\eta^{-1}} = \{\eta^{-1} \cdot z_0\}$. Let $\mathcal{V}$ denote the finite collection of all subsets of W which are the sub-unions of the disjoint union given above; that is, $\mathcal{V}$ consists of all sets of the form $\bigcup_{\psi \in \Psi} V_\psi$ or $V_0 \cup \bigcup_{\psi \in \Psi} V_\psi$ or $\{z_0\} \cup \bigcup_{\psi \in \Psi} V_\psi$ or $\{z_0\} \cup V_0 \cup \bigcup_{\psi \in \Psi} V_\psi$ for $\Psi \subset \Psi^\dagger$. We shall apply Proposition 1.3 with these choices of W and $\mathcal{V}$. We take D to be a number, and $(\mathcal{M}_V)_{V \in \mathcal{V}}$ a family

of conformal densities, for which conditions (i)-(iv) of Proposition 1.3 are satisfied. We write $\mathcal{M}_V = (\mu_{V,z})_{z \in \mathbb{H}^3}$. We set $\mathcal{M}_{\Gamma,z_0} = \mathcal{M}_W$, and $\nu_\psi = \mu_{V_\psi,z_0}$ for each $\psi \in \Psi^\dagger$. According to Proposition 1.2, $\mathcal{M}$ is Γ -invariant. It follows from condition (i) of Proposition 1.3 and the definition of a conformal density that $\mu_{z_0} = \mu_{W,z_0} \neq 0$; hence after multiplying by a constant we may assume that μ_{z_0} has total mass 1. This is conclusion (1) of Lemma 3.3. We shall show that the other conclusions of Lemma 3.3 also hold with these definitions of $\mathcal{M}$ and ν_ψ .

By (1) of Lemma 3.3 and condition (ii) of Proposition 1.3, we have $\mu_{z_0} = \mu_{W,z_0}$ where

$$\mu_{W,z_0} = \mu_{\{z_0\},z_0} + \mu_{V_0,z_0} + \sum_{\psi \in \Psi^\dagger} \mu_{V_\psi,z_0}.$$

But we know that $\mu_{\{z_0\},z_0} = 0$ and $\mu_{V_0,z_0} = 0$ by condition (iv) of Proposition 1.3. Hence we get $\mu_{z_0} = \sum_{\psi \in \Psi^\dagger} \nu_\psi$, which is conclusion (2) of Lemma 3.3. In order to complete part (3) of Lemma 3.3, we determine all of the group theoretical relations between the set of words J_ψ for $\psi \in \Psi^\dagger$.

Let us consider $\delta_{i,j}J_{\delta^{i,j}}$ for $(i,j) \in A_6$. We know that $(\delta_{i,j})^{-1} \in J_{\delta^{i,j}}$ and $\delta^{i,j} \in J_{\delta^{i,j}}$. Therefore, we have $1 \in \delta_{i,j}J_{\delta^{i,j}}$ and $\pi(i) \in J_{\delta^{i,j}}$. Let w be a word in $J_{\delta^{i,j}}$ so that $w \neq \delta^{i,j}$. Then we have $w = \delta^{i,j}w_1$ or $w = (\pi(j))^{-1}w_1$ for some word $w_1 \in \Gamma$. We get $\delta_{i,j}w = \pi(i)w_1$. The first letter of w_1 cannot be $\pi(j)$. Therefore, $\delta_{i,j}J_{\delta^{i,j}}$ cannot contain any word with initial word $\pi(i)\pi(j) = \delta_{i,j}$. Note that we have either $\pi(i) = \pi(j)$ or $\pi(i) \neq \pi(j)$. Assume that $\pi(i) = \pi(j)$. The first letter of w_1 can be one of the isometries $\pi(i)^{-1}, \pi(k), \pi(k)^{-1}$ where $k \in \{1, 2, 3, 4\}$ so that $k \neq i$. If the first letter of w_1 is $\pi(i)^{-1}$, then we have $w_1 = \pi(i)^{-1}w_2$ for some word $w_2 \in \Gamma$. The first letter of w_2 cannot be $\pi(i)$. But it can be one of the isometries $\pi(i)^{-1}, \pi(k), \pi(k)^{-1}$. We know that $\pi(i) \in \Psi_r^\dagger$. Note that either $\pi(k)$ or $\pi(k)^{-1}$ is in $\Psi_r^\dagger$. Without loss of generality, let us assume that $\pi(k)^{-1} \in \Psi_r^\dagger$. Since we have $\delta_{i,j}w = w_2$, we conclude that $\delta_{i,j}J_{\delta^{i,j}}$ contains the words $1, \pi(i), \pi(k)$ and every word starting with $\pi(i)^{-1}, \pi(k)^{-1}, \pi(k)\pi(i), \pi(k)\pi(k)$ and $\pi(k)\pi(i)^{-1}$. If the first letter of w_2 is not $\pi(i)^{-1}$, then $\delta_{i,j}J_{\delta^{i,j}}$ contains all the words starting with $\pi(i)\pi(k)$ and $\pi(i)\pi(k)^{-1}$.

Assume that $\pi(i) \neq \pi(j)$. By the definition of A_6 , we know that $\pi(i)\pi(j) \neq 1$. The first letter of w_1 can be one of the isometries $\pi(i), \pi(i)^{-1}, \pi(j)^{-1}$. If the first letter of w_1 is $\pi(i)^{-1}$, then we have $w_1 = \pi(i)^{-1}w_2$ for some word $w_2 \in \Gamma$. The first letter of w_2 cannot be $\pi(i)$. But it can be one of the isometries $\pi(i)^{-1}, \pi(j), \pi(j)^{-1}$. We know that $\pi(j) \in \Psi_r^\dagger$. Note that either $\pi(i)$ or $\pi(i)^{-1}$ is in $\Psi_r^\dagger$. Without loss of generality, let us assume that $\pi(i)^{-1} \in \Psi_r^\dagger$. Since we have $\delta_{i,j}w = w_2$, we conclude that $\delta_{i,j}J_{\delta^{i,j}}$ contains the words $1, \pi(i), \pi(j)$ and every word starting with $\pi(i)^{-1}, \pi(j)^{-1}, \pi(j)\pi(i), \pi(j)\pi(j)$ and $\pi(j)\pi(i)^{-1}$. If the first letter of w_2 is not $\pi(i)^{-1}$, then all words starting with $\pi(i)\pi(i)$ and $\pi(i)\pi(j)^{-1}$. In either case, $\delta_{i,j}J_{\delta^{i,j}}$ contains every word in Γ except

the ones that start with $\delta_{i,j}$. Thus, it follows from the definitions of the sets J_ψ for each $\psi \in \Psi^\dagger$ and the discussion presented above that for every isometry $\gamma \in A^\dagger$ we obtain

$$(13) \quad \gamma J_{s(\gamma)} = \Gamma - J_{s(\gamma)}.$$

Let V_Ψ denotes the union $\bigcup_{\gamma \in \Psi} V_\gamma$ where Ψ is a subset of $\Psi^\dagger \cup \Psi_r^\dagger$. Using the group-theoretical relations in (13), we derive the following measure-theoretical relations

$$(14) \quad \gamma V_{s(\gamma)} = W - V_{s(\gamma)}$$

between the orbits V_ψ for every isometry $\gamma \in A^\dagger$ so that $s(\gamma) = \psi$ for every $\psi \in \Psi^\dagger$.

Since we have $W - V_{s(\gamma)} = V_{s(\gamma)} \in \mathcal{V}$, the condition (iii) of Proposition 1.3 gives $\mathcal{M}_{V_{s(\gamma)}} = \gamma_\infty^* (\mathcal{M}_{W-V_{s(\gamma)}})$. On the other hand, by Proposition 1.3(ii) we find that $\mathcal{M}_{\Gamma, z_0} = \mathcal{M}_{W-V_{s(\gamma)}} + \mathcal{M}_{V_{s(\gamma)}}$. Therefore, we obtain $\mathcal{M}_{V_{s(\gamma)}} = \gamma_\infty^* (\mathcal{M}_{\Gamma, z_0} - \mathcal{M}_{V_{s(\gamma)}})$. Proposition 1.3(ii) implies that $\mu_{V_{s(\gamma)}, \gamma \cdot z_0} = \gamma_\infty^* (\mu_{z_0} - \sum_{\psi \in S(\gamma)} \nu_\psi)$ which gives that $d\mu_{V_{s(\gamma)}, \gamma \cdot z_0} = d(\gamma_\infty^* (\mu_{z_0} - \sum_{\psi \in S(\gamma)} \nu_\psi))$ or $d\mu_{V_{s(\gamma)}, \gamma \cdot z_0} = d(\mu_{z_0} - \sum_{\psi \in S(\gamma)} \nu_\psi)$. For every isometry $\gamma \in A^\dagger$, equating the total masses of the two sides of the last equality given above, we obtain that

$$\int \lambda_{\gamma, z_0}^D d\mu_{V_{s(\gamma)}} = 1 - \sum_{\psi \in S(\gamma)} \int d\nu_\psi$$

where we use the equality $d\mu_{V_{s(\gamma)}, \gamma \cdot z_0} = \mathcal{P}(z_0, s(\gamma) \cdot z_0, \cdot)^D d\mu_{V_{s(\gamma)}}$ which follows from the facts that $\mathcal{M}_{V_{s(\gamma)}}$ is a D -conformal density and $d\mu_{V_{s(\gamma)}, \gamma \cdot z_0} = \lambda_{\gamma, z_0}^D d\mu_{V_{s(\gamma)}}$ by (1). $\square$

3.3 Displacement functions for the decomposition $\Gamma_{\mathcal{D}^\dagger}$

Suppose that Γ , ξ , and η satisfy the hypotheses of Proposition 1.3. Let z_0 be any point in $\mathbb{H}^3$. Since Γ and $\Xi = \{\xi, \eta\}$ satisfy the hypotheses of Lemma 3.3, there exists a number $D \in [0, 2]$, a conformal density $\mathcal{M}_{\Gamma, z_0}$ and a family of Borel measures $(\nu_\psi)_{\psi \in \Psi^\dagger}$ where $\Psi^\dagger = \{\xi\eta, \xi^2, \xi\eta^{-1}, \eta, \xi^{-1}, \eta^{-1}\xi^{-1}, \eta^{-2}, \eta^{-1}\xi\}$ for which conclusions (1), (2), and (3) of Lemma 3.3 hold. Hypotheses of Proposition 1.3 imply that $\mathcal{M}_{\Gamma, z_0}$ is a constant multiple of the area density $\mathcal{A}$. Therefore, we get $D = 2$. Since μ_{z_0} has total mass 1 by Lemma 3.3(1), we have $\mathcal{M}_{\Gamma, z_0} = \mathcal{A}$. We know that $\nu_\psi(S_\infty) = \int d\nu_\psi$ for every $\psi \in \Psi^\dagger$. By Lemma 3.3(1), we obtain that $\sum_{\psi \in \Psi^\dagger} \nu_\psi(S_\infty) = 1$. We see that $0 \leq \nu_\psi(S_\infty) \leq 1$ for every $\psi \in \Psi^\dagger$.

We show that $0 < \nu_\psi(S_\infty) < 1$ for every $\psi \in \Psi^\dagger$. To this purpose, it is enough to show that $\nu_{\psi_0}(S_\infty) \neq 0$ for every $\psi_0 \in \Psi^\dagger$ because if we have $\nu_{\psi_0}(S_\infty) = 1$ for some $\psi_0 \in \Psi^\dagger$, then we obtain that $\nu_\psi(S_\infty) = 0$ for every $\psi \in \Psi^\dagger - \{\psi_0\}$. Assume that

$\nu_{\psi_0} = 0$ for a given $\psi_0 \in \{\xi^{-1}, \eta\}$. Let us consider the following lists where each entry in a list is assigned for $\psi_0, \gamma_0, S(\gamma_0), \psi_1, \psi_2, \gamma_2$ and $S(\gamma_2)$, respectively:

$$\begin{aligned} &(\xi^{-1}, \eta^{-1}\xi, \{\eta^{-1}\xi\}, \eta^{-1}\xi, \eta, \xi\eta^{-1}, \{\xi\eta^{-1}\}), \\ &(\eta, \xi\eta^{-1}, \{\xi\eta^{-1}\}, \xi\eta^{-1}, \xi^{-1}, \eta^{-1}\xi, \{\eta^{-1}\xi\}). \end{aligned}$$

Since we assume that $\nu_{\psi_0}(S_\infty) = 0$ and $\psi_0 = s(\gamma_0)$, we get $\sum_{\psi \in S(\gamma_0)} \nu_\psi = 1$ by Lemma 3.3(3). We see that $\nu_{\psi_1}(S_\infty) \neq 0$ for some $\psi_1 \in S(\gamma_0)$. Let ψ_2 be an element in $\Psi^\dagger - S(\gamma_0)$. Then there is an isometry $\gamma_2 \in \Psi^\dagger$ so that $\psi_2 = s(\gamma_2)$ and $S(\gamma_0) \cap S(\gamma_2) = \emptyset$. Since we have $\sum_{\psi \in \Psi^\dagger} \nu_\psi = 1$ and $\sum_{\psi \in S(\gamma_0)} \nu_\psi = 1$, we derive that $\nu_{\psi_2}(S_\infty) = 0$. By Lemma 3.3, we obtain that $\sum_{\psi \in S(\gamma_2)} \nu_\psi = 1$. Using the facts that $\sum_{\psi \in \Psi^\dagger} \nu_\psi = 1$ and $S(\gamma_0) \cap S(\gamma_2) = \emptyset$, we conclude that $\nu_{\psi_1}(S_\infty) = 0$, a contradiction. Therefore, we conclude that $\nu_{\xi^{-1}}(S_\infty) \neq 0$ and $\nu_\eta(S_\infty) \neq 0$. Assume that $\nu_{\psi_0} = 0$ for a given $\psi_0 \in \{\xi\eta, \xi\eta^{-1}, \xi^2, \eta^{-1}\xi, \eta^{-1}\xi^{-1}, \eta^{-2}\}$. Let us consider the following lists where each entry in a list is assigned for $\psi_0, \gamma_0, S(\gamma_0), \psi_1$, respectively:

$$\begin{aligned} &(\xi\eta, \xi^{-1}, \Psi^\dagger - \{\eta\}, \xi^{-1}), (\xi\eta^{-1}, \xi^{-1}, \Psi^\dagger - \{\eta^{-1}, \eta^{-1}\xi, \eta^{-1}\xi^{-1}, \eta^{-2}\}, \eta), \\ &(\xi^2, \xi^{-1}, \Psi^\dagger - \{\xi\}, \xi^{-1}), (\eta^{-2}, \eta, \Psi^\dagger - \{\eta^{-1}, \eta^{-1}\xi, \eta^{-1}\xi^{-1}, \eta^{-2}\}, \xi^{-1}), \\ &(\eta^{-1}\xi^{-1}, \eta, \Psi^\dagger - \{\xi^{-1}\}, \eta), (\eta^{-1}\xi, \eta, \Psi^\dagger - \{\xi, \xi\eta^{-1}, \xi\eta, \xi^2\}, \xi^{-1}). \end{aligned}$$

Since we assume that $\nu_{\psi_0}(S_\infty) = 0$ and $\psi_0 = s(\gamma_0)$, we get $\sum_{\psi \in S(\gamma_0)} \nu_\psi = 1$ by Lemma 3.3(3). We have $\psi_1 \notin S(\gamma_0)$. Therefore, we obtain that $\nu_{\psi_1}(S_\infty) = 0$ for some $\psi_1 \in \{\xi^{-1}, \eta\}$. This is a contradiction because $\nu_{\xi^{-1}}(S_\infty) \neq 0$ and $\nu_\eta(S_\infty) \neq 0$. We conclude that $0 < \nu_\psi(S_\infty) < 1$ for every $\psi \in \{\xi\eta, \xi\eta^{-1}, \xi^2, \eta^{-1}\xi, \eta^{-1}\xi^{-1}, \eta^{-2}\}$. Since we have $\psi = s(\gamma)$ for some $\gamma \in A^\dagger$ and $S(\gamma) \subset \Psi^\dagger$, we also conclude by Lemma 3.3(3) that

$$0 < \int \lambda_{\gamma, z_0}^2 d\mu_{V_{S(\gamma)}} = 1 - \sum_{\psi \in S(\gamma)} \int d\nu_\psi < 1$$

for every $\gamma \in A^\dagger$. In other words, $\nu_{S(\gamma)}$ and $\int \lambda_{\gamma, z_0}^2 d\mu_{V_{S(\gamma)}}$ satisfy the hypothesis of Lemma 1.1. We apply Lemma 1.1 to every $\gamma \in A^\dagger$ with $\nu = \nu_{S(\gamma)}$, $a = \nu_{S(\gamma)}(S_\infty)$ and $b = \int \lambda_{\gamma, z_0}^2 d\mu_{V_{S(\gamma)}}$. We obtain lower bounds

$$e^{2\text{dist}(z_0, \gamma \cdot z_0)} \geq \frac{\int \lambda_{\gamma, z_0}^2 d\mu_{V_{S(\gamma)}} \cdot (1 - \int d\nu_{S(\gamma)})}{\int d\nu_{S(\gamma)} \cdot (1 - \int \lambda_{\gamma, z_0}^2 d\mu_{V_{S(\gamma)}})}$$

for every isometry in $\Psi^\dagger \cup \Psi_r^\dagger$. Using Lemma 3.3 part (3), we calculate lower bounds

$$(15) \quad e^{2\text{dist}(z_0, \gamma \cdot z_0)} \geq \frac{\left(1 - \sum_{\psi \in S(\gamma)} m_{\sigma^\dagger(\psi)}\right) \cdot (1 - m_{\sigma^\dagger(S(\gamma))})}{\left(\sum_{\psi \in S(\gamma)} m_{\sigma^\dagger(\psi)}\right) \cdot m_{\sigma^\dagger(S(\gamma))}}$$

where $\int d\nu_\psi = m_{\sigma^\dagger(\psi)}$ for a bijection $\sigma^\dagger: \Psi^\dagger \rightarrow I^\dagger = \{i \in \mathbb{Z}: 1 \leq i \leq 8\}$. Note that we have 18 lower bounds given in the expression (15) for the displacements under the

isometries in $\Psi_r^\dagger \cup \Psi^\dagger$ since there are 18 group theoretical relations listed in Lemma 3.3. We replace each constant $m_{\sigma^\dagger(\psi)}$ appearing in (15) with the variable $x_{\sigma^\dagger(\psi)}$ for every $\psi \in S(\gamma) \cup \{s(\gamma)\}$ for every $\gamma \in A_1^\dagger \cup A_2^\dagger \cup A_3^\dagger$. Then we introduce the following $8 = |A_1^\dagger| + |A_2^\dagger| + |A_3^\dagger|$ displacement functions

$$f_{\sigma^\dagger(s(\gamma))}^\dagger = \frac{\left(1 - \sum_{\psi \in S(\gamma)} x_{\sigma^\dagger(\psi)}\right) \cdot (1 - x_{\sigma^\dagger(s(\gamma))})}{\left(\sum_{\psi \in S(\gamma)} x_{\sigma^\dagger(\psi)}\right) \cdot x_{\sigma^\dagger(s(\gamma))}}$$

for $\text{dist}(z_0, \xi \cdot z_0)$, $\text{dist}(z_0, \eta \cdot z_0)$, $\text{dist}(z_0, \xi^{-1} \cdot z_0)$, $\text{dist}(z_0, \eta^{-1} \cdot z_0)$, $\text{dist}(z_0, \xi\eta \cdot z_0)$ and $\text{dist}(z_0, \eta^{-1}\xi^{-1} \cdot z_0)$. Similarly, we replace each $m_{\sigma^\dagger(\psi)}$ appearing in (15) with the variable $x_{\sigma^\dagger(\psi)}$ for every $\psi \in S(\gamma) \cup \{s(\gamma)\}$ for every $\gamma \in A_4^\dagger \cup A_5^\dagger$. As a result, we obtain $6 = |A_4^\dagger| + |A_5^\dagger|$ more displacement functions $g_{\sigma^\dagger(s(\gamma))}$ for $\text{dist}(z_0, \xi^{-1} \cdot z_0)$, $\text{dist}(z_0, \eta \cdot z_0)$, $\text{dist}(z_0, \eta^{-1} \cdot z_0)$, and $\text{dist}(z_0, \xi\eta \cdot z_0)$.

Finally, we replace $m_{\sigma^\dagger(\psi)}$ appearing in (15) with the variable $x_{\sigma^\dagger(\psi)}$ for every $\psi \in S(\gamma) \cup \{s(\gamma)\}$ for every $\gamma \in A_6^\dagger$. We find $4 = |A_6^\dagger|$ more displacement functions $h_{\sigma^\dagger(s(\gamma))}$ for the displacements $\text{dist}(z_0, \xi^2 \cdot z_0)$, $\text{dist}(z_0, \eta^{-2} \cdot z_0)$, $\text{dist}(z_0, \xi\eta^{-1} \cdot z_0)$, and $\text{dist}(z_0, \eta\xi \cdot z_0)$. There are totally 18 displacement functions. Since we know that $\text{dist}(z_0, \gamma \cdot z_0) = \text{dist}(z_0, \gamma^{-1} \cdot z_0)$ for every isometry γ and are interested in the displacements under the isometries in $\Gamma_\dagger = \{\xi, \eta, \xi\eta\}$, we will concentrate on the first 14 displacement functions $f_1^\dagger, f_2^\dagger, \dots, f_8^\dagger, g_1^\dagger, g_2^\dagger, \dots, g_6^\dagger$ for the proof of Theorem 5.1.

The constants obtained on the right hand-side of the inequalities in the expression (15) can be considered as the values of the functions $f_1^\dagger, \dots, f_8^\dagger, g_1^\dagger, \dots, g_6^\dagger$ at the point $\mathbf{m}_\dagger = (m_1, m_2, \dots, m_8) \in \Delta^7$ where Δ^7 denotes the 7-dimensional simplex $\{(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8) \in \mathbb{R}_+^8 : \sum_{i=1}^8 x_i = 1\}$. To calculate a lower bound for the maximum of $\text{dist}(z_0, \xi \cdot z_0)$, $\text{dist}(z_0, \eta \cdot z_0)$, and $\text{dist}(z_0, \xi\eta \cdot z_0)$, it is required to find the greatest lower bound for the maximum of the functions $f_1^\dagger, \dots, f_8^\dagger, g_1^\dagger, \dots, g_6^\dagger$ over the simplex Δ^7 . Points of Δ^3 will be written in bold fonts, e.g., $\mathbf{x} = (x_1, \dots, x_8)$. Let $\mathcal{F}^\dagger = \{f_1^\dagger, f_2^\dagger, \dots, f_8^\dagger, g_1^\dagger, g_2^\dagger, \dots, g_6^\dagger\}$. Let $G^\dagger$ be the continuous function defined as

$$\begin{aligned} G^\dagger : \quad \Delta^7 &\rightarrow \mathbb{R} \\ \mathbf{x} &\mapsto \max\{f(\mathbf{x}) : f \in \mathcal{F}^\dagger\}. \end{aligned}$$

We aim to calculate $\inf_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x})$. We will need different parametrization of Δ^7 . Let $\Delta^{7,s}$ be the set of points $(x_{i_1}, \dots, x_{i_7}) \in \mathbb{R}^7$ with the ordered $x_{i_1}x_{i_2} \cdots x_{i_7}$ -coordinate system so that $0 < x_{i_1} < 1$, $0 < x_{i_2} < 1 - x_{i_1}$, $\dots$, $0 < x_{i_7} < 1 - x_{i_1} - \dots - x_{i_6}$ where $i_1 < i_2 < \dots < i_7$ and $i_1, i_2, \dots, i_7 \in I^\dagger - \{s\}$. For each $s \in I^\dagger$, let $p_s^7 : \Delta^{7,s} \rightarrow \Delta^7$

denote the parametrization of Δ^7 where $x_s = 1 - \sum_{m \in I^\dagger - \{s\}} x_m$. Then we have the following commuting diagrams

$$\begin{array}{ccc} \Delta^{7,s} & & \\ \downarrow p_s^7 & \searrow f_j^{\dagger,s} & \\ \Delta^7 & \xrightarrow{f_j^\dagger} & \mathbb{R} \end{array}$$

for every $j, s \in I^\dagger$. We shall use the notation $f_i^{\dagger,s}$ to denote the formulas of the displacement functions $f_i^\dagger$ for $i \in I^\dagger$ under the parametrization p_s^7 for $s \in I^\dagger$. A generic point in $\Delta^{7,s}$ will be written as $\mathbf{x}_s = (p_s^7)^{-1}(\mathbf{x})$, e.g., $\mathbf{x}_1 = (x_2, \dots, x_8) \in \Delta^{7,1}$. Let us introduce the index sets $I_1^\dagger = \{1, 2, 3\}$, $I_2^\dagger = \{4, 5, 6\}$, $I_3^\dagger = \{7, 8\}$, $I_4^\dagger = I_2^\dagger \cup \{7\}$, $I_5^\dagger = I_1^\dagger \cup \{8\}$ and $J_i^\dagger = I^\dagger - \{i\}$ for $i \in I^\dagger$. Let $\mathfrak{J}^\dagger = \{I_j^\dagger : j = 1, 2, 3, 4, 5\}$ and $\mathfrak{J}^\dagger = \{J_i^\dagger : i \in I^\dagger\}$.

4 Displacements under the isometries in $\Gamma_\dagger$ when $\Lambda_\Gamma = S_\infty$

In this section, we prove an analog of Proposition 1.5 for the set of isometries in $\Gamma_\dagger = \{\xi, \eta, \xi\eta\} \subset \Psi_r^\dagger \cup \Psi^\dagger$ which is stated as follows

Proposition 4.1 *Let ξ and η be two orientation-preserving isometries of $\mathbb{H}^3$. Suppose that the group Γ generated by ξ and η is discrete and is free on the generators ξ and η . Suppose in addition that every Γ -invariant conformal density for $\mathbb{H}^3$ is a constant multiple of the area density. Then for any $z \in \mathbb{H}^3$ we have*

$$e\left(2 \max_{\gamma \in \Gamma_\dagger} \{\text{dist}(z_0, \gamma \cdot z_0)\}\right) \geq \alpha_\dagger$$

Proposition 4.1 is required to complete the proof of Theorem 5.1 when $\Lambda_\Gamma = S_\infty$. The lower bounds given in the proposition above are obtained by calculating $\inf_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x})$. To this purpose, we gather some facts that are necessary to complete this task in the following subsections. Before we calculate $\inf_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x})$, we shall develop some notation to shorten the equations we need to study in this section.

Let $\mathbf{x} = (x_i)_{i \in I^\dagger}$ be a point in Δ^7 and $\mathbf{x}^* = (x_i^*)_{i \in I^\dagger}$, $\mathbf{x}^\circ = (x_i^\circ)_{i \in I^\dagger}$ be two fixed points in Δ^7 . We shall need the following functions in addition to σ and τ we introduced in §2.2.: $\rho_i, \Sigma_i, \Sigma_{i,j} : \Delta^7 \rightarrow (0, 1)$ for each index set $I_i^\dagger, I_j^\dagger \in \mathfrak{J}^\dagger$, $\rho^i, \Sigma^i : \Delta^7 \rightarrow (0, 1)$ for each index set $J_i^\dagger \in \mathfrak{J}^\dagger$ and $\sigma_i : \Delta^7 \rightarrow (0, 1)$ for each $i \in I^\dagger$ where

$$\Sigma_i(\mathbf{x}) = \sum_{k \in I_i^\dagger} x_k, \Sigma_{i,j}(\mathbf{x}) = \sum_{k \in I_i^\dagger \cup I_j^\dagger} x_k, \Sigma^i(\mathbf{x}) = \sum_{k \in J_i^\dagger} x_k,$$

$\rho_i(\mathbf{x}) = \sigma(\Sigma_i(\mathbf{x}))$, $\rho_{i,j}(\mathbf{x}) = \sigma(\Sigma_{i,j}(\mathbf{x}))$, $\rho^i(\mathbf{x}) = \sigma(\Sigma^i(\mathbf{x}))$, and $\sigma_i(\mathbf{x}) = \sigma(x_i)$. Let us define $\rho_i^* = \rho_i(\mathbf{x}^*)$ and $\rho_{i,j}^* = \rho_{i,j}(\mathbf{x}^*)$ for every $I_i^\dagger, I_j^\dagger \in \mathcal{J}^\dagger$ and $\sigma_i^* = \sigma_i(\mathbf{x}^*)$ for every $i \in I^\dagger$. For each parametrization $\mathbf{p}_s^7$, let us define $\Sigma_{\mathbf{i}}, \Sigma_{\mathbf{i},j}: \Delta^{7,s} \rightarrow (0,1)$ for index sets $I_i^\dagger, I_j^\dagger \in \mathcal{J}^\dagger - \{I_s^\dagger\}$ and $\Sigma^{\mathbf{i}}: \Delta^{7,s} \rightarrow (0,1)$ for any index set $J_i^\dagger \in \mathcal{J}^\dagger$ so that

$$\Sigma_{\mathbf{i}}(\mathbf{x}_s) = \sum_{k \in I_i^\dagger} x_k, \quad \Sigma_{\mathbf{i},j}(\mathbf{x}_s) = \sum_{k \in I_i^\dagger \cup I_j^\dagger} x_k, \quad \text{and} \quad \Sigma^{\mathbf{i}}(\mathbf{x}_s) = \sum_{k \in J_i^\dagger} x_k.$$

Let $\mathbf{x}_s^* = (\mathbf{p}_s^7)^{-1}(\mathbf{x}_s)$ and $\mathbf{x}_s^\circ = (\mathbf{p}_s^7)^{-1}(\mathbf{x}_s)$. Then we set $\Sigma_{\mathbf{i}}^* = \Sigma_{\mathbf{i}}(\mathbf{x}_s^*)$, $\Sigma_{\mathbf{i}}^\circ = \Sigma_{\mathbf{i}}(\mathbf{x}_s^\circ)$, $\Sigma_{\mathbf{i},j}^* = \Sigma_{\mathbf{i},j}(\mathbf{x}_s^*)$, $\Sigma_{\mathbf{i},j}^\circ = \Sigma_{\mathbf{i},j}(\mathbf{x}_s^\circ)$, $\Sigma^{\mathbf{i}}(\mathbf{x}_s^*) = \Sigma_{\mathbf{i}}^*$ and $\Sigma^{\mathbf{i}}(\mathbf{x}_s^\circ) = \Sigma_{\mathbf{i}}^\circ$. We will use $\Sigma_{\mathbf{0}}^*$ to denote $\Sigma_{i \in I^\dagger} x_i^* = 1$. Let us introduce the functions $\tau_i, \tau^i, \sigma_i: \Delta^{7,s} \rightarrow (0,1)$ for every $i \in J_s^\dagger$ defined by $\tau_i(\mathbf{x}_s) = \tau(x_i)$, $\tau^i(\mathbf{x}_s) = \tau(1 - x_i)$ and $\sigma_i(\mathbf{x}_s) = \sigma(x_i)$ so that $\tau_i^* = \tau_i(\mathbf{x}_s^*)$, $\sigma_i^* = \sigma_i(\mathbf{x}_s^*)$ and $\sigma_i^\circ = \sigma_i(\mathbf{x}_s^\circ)$. Also introduce $\tau^{\mathbf{i}}, \sigma^{\mathbf{i}}: \Delta^{7,s} \rightarrow (0,1)$ for every index set $J_i^\dagger \in \mathcal{J}^\dagger$ defined by $\tau^{\mathbf{i}}(\mathbf{x}_s) = \tau(1 - \Sigma^{\mathbf{i}}(\mathbf{x}_s))$ and $\sigma^{\mathbf{i}}(\mathbf{x}_s) = 1/\sigma(\Sigma^{\mathbf{i}}(\mathbf{x}_s))$ where $\sigma_{\mathbf{i}}^* = \sigma^{\mathbf{i}}(\mathbf{x}_s^*)$ and $\sigma_{\mathbf{i}}^\circ = \sigma^{\mathbf{i}}(\mathbf{x}_s^\circ)$.

Let us define the functions $\rho_{\mathbf{i}}, \rho^{\mathbf{i}}, \rho_{\mathbf{i},j}, \rho^{\mathbf{i},j}: \Delta^{7,s} \rightarrow (0,1)$ for every $I_i^\dagger, I_j^\dagger \in \mathcal{J}^\dagger$ so that $\rho_{\mathbf{i}}(\mathbf{x}_s) = \sigma(\Sigma_{\mathbf{i}}(\mathbf{x}_s))$, $\rho_{\mathbf{i}}^* = \rho_{\mathbf{i}}(\mathbf{x}_s^*)$, $\rho_{\mathbf{i}}^\circ = \rho_{\mathbf{i}}(\mathbf{x}_s^\circ)$, $\rho^{\mathbf{i}}(\mathbf{x}_s) = 1/\rho_{\mathbf{i}}(\mathbf{x}_s)$, $\rho_{\mathbf{i}}^{\mathbf{i}} = \rho^{\mathbf{i}}(\mathbf{x}_s)$, $\rho_{\mathbf{i},j}(\mathbf{x}_s) = \sigma(\Sigma_{\mathbf{i},j}(\mathbf{x}_s))$, $\rho_{\mathbf{i},j}^* = \rho_{\mathbf{i},j}(\mathbf{x}_s^*)$, $\rho_{\mathbf{i},j}^\circ = \rho_{\mathbf{i},j}(\mathbf{x}_s^\circ)$, $\rho^{\mathbf{i},j}(\mathbf{x}_s) = 1/\rho_{\mathbf{i},j}(\mathbf{x}_s)$, $\rho_{\mathbf{i},j}^{\mathbf{i},j} = \rho^{\mathbf{i},j}(\mathbf{x}_s)$, and $\rho_{\mathbf{i},j}^{\mathbf{i},j} = \rho^{\mathbf{i},j}(\mathbf{x}_s^\circ)$. Let us also define $\varrho_{\mathbf{i}}, \varrho^{\mathbf{i}}, \varrho_{\mathbf{i},j}, \varrho^{\mathbf{i},j}: \Delta^{7,s} \rightarrow (0,1)$ for the index sets $I_i^\dagger, I_j^\dagger \in \mathcal{J}^\dagger$ such that $\varrho_{\mathbf{i}}(\mathbf{x}_s) = \tau(\Sigma_{\mathbf{i}}(\mathbf{x}_s))$, $\varrho^{\mathbf{i}}(\mathbf{x}_s) = \tau(1 - \Sigma_{\mathbf{i}}(\mathbf{x}_s))$, $\varrho_{\mathbf{i},j}(\mathbf{x}_s) = \tau(\Sigma_{\mathbf{i},j}(\mathbf{x}_s))$, and $\varrho^{\mathbf{i},j}(\mathbf{x}_s) = \tau(1 - \Sigma_{\mathbf{i},j}(\mathbf{x}_s))$. Then we get the partial derivatives

$$(\sigma_i)_m = \begin{cases} -\tau_i < 0 & \text{if } m = i, \\ 0 & \text{otherwise,} \end{cases} \quad (\sigma^{\mathbf{i}})_m = \begin{cases} \tau^{\mathbf{i}} > 0 & \text{if } m \in J_i^\dagger, \\ 0 & \text{otherwise,} \end{cases}$$

$$(\rho_{\mathbf{i}})_m = \begin{cases} -\varrho_{\mathbf{i}} < 0 & \text{if } m \in I_i^\dagger, \\ 0 & \text{otherwise,} \end{cases} \quad (\rho^{\mathbf{i}})_m = \begin{cases} \varrho^{\mathbf{i}} > 0 & \text{if } m \in I_i^\dagger, \\ 0 & \text{otherwise,} \end{cases}$$

$$(\rho_{\mathbf{i},j})_m = \begin{cases} -\varrho_{\mathbf{i},j} < 0 & \text{if } m \in I_i^\dagger \cup I_j^\dagger, \\ 0 & \text{otherwise.} \end{cases} \quad (\rho^{\mathbf{i},j})_m = \begin{cases} \varrho^{\mathbf{i},j} > 0 & \text{if } m \in I_i^\dagger \cup I_j^\dagger, \\ 0 & \text{otherwise.} \end{cases}$$

4.1 Infimum of the maximum of the functions in $\mathcal{F}^\dagger$ over Δ^7

There is a bijection $\sigma^\dagger: \Psi^\dagger \rightarrow I^\dagger$ so that we obtain the following displacement functions $f_i^\dagger: \Delta^7 \rightarrow \mathbb{R}_+$ for every $i \in I^\dagger$ where

$$\begin{aligned} f_1^\dagger(\mathbf{x}) &= \frac{1-x_4-x_5-x_6}{x_4+x_5+x_6} \cdot \frac{1-x_1}{x_1}, f_2^\dagger(\mathbf{x}) = \frac{1-x_4-x_5-x_6-x_7-x_8}{x_4+x_5+x_6+x_7+x_8} \cdot \frac{1-x_2}{x_2}, \\ f_3^\dagger(\mathbf{x}) &= \frac{1-x_1-x_2-x_3-x_7-x_8}{x_1+x_2+x_3+x_7+x_8} \cdot \frac{1-x_3}{x_3}, f_4^\dagger(\mathbf{x}) = \frac{1-x_1-x_2-x_3}{1-x_1-x_2-x_3} \cdot \frac{1-x_4}{x_4}, \\ f_5^\dagger(\mathbf{x}) &= \frac{1-x_1-x_2-x_3-x_7-x_8}{x_1+x_2+x_3+x_7+x_8} \cdot \frac{1-x_5}{x_5}, f_7^\dagger(\mathbf{x}) = \frac{1-x_4-x_5-x_6}{x_4+x_5+x_6} \cdot \frac{1-x_7}{x_7}, \\ f_6^\dagger(\mathbf{x}) &= \frac{1-x_4-x_5-x_6-x_7-x_8}{x_4+x_5+x_6+x_7+x_8} \cdot \frac{1-x_6}{x_6}, f_8^\dagger(\mathbf{x}) = \frac{1-x_1-x_2-x_3}{x_1+x_2+x_3} \cdot \frac{1-x_8}{x_8}. \end{aligned}$$

We define the following continuous function

$$\begin{aligned} F^\dagger : \Delta^7 &\rightarrow \mathbb{R} \\ \mathbf{x} &\mapsto \max \left(f_1^\dagger(\mathbf{x}), \dots, f_8^\dagger(\mathbf{x}) \right). \end{aligned}$$

We show that $\inf_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x}) = \inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$ (see Lemma 4.14). Therefore, it is enough to calculate $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$. Since we have $f_i^\dagger(\mathbf{x}) > 1$ for every $\mathbf{x} \in \Delta^7$ for $i \in I^\dagger$, we have $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) \geq 1$.

Let us define the subsets $\Delta_n = \{\mathbf{x} \in \Delta^7: 1/n \leq x_i \leq 1 - 1/(2n) \text{ for } i \in I^\dagger\}$ of Δ^7 for every $n \geq 2$. Then $F^\dagger$ has an absolute minimum α_n at some point $\mathbf{x}_n$ in each Δ_n since Δ_n is compact. Let us consider the sequences (α_n) and $(\mathbf{x}_n)$. It is clear that (α_n) is a decreasing sequence because $\Delta_n \subset \Delta_{n+1}$. Since (α_n) is bounded below by 1, it converges to a finite number greater than or equal to 1. This means that $F^\dagger(\mathbf{x}_n)$ approaches to $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$.

We claim that the sequence $(\mathbf{x}_n)$ cannot have a limit point on the boundary of Δ^7 . Let us assume contrary that $\mathbf{x}_n \rightarrow \mathbf{b} \in \partial\Delta^7$ as $n \rightarrow \infty$. If $(b_1, b_2, \dots, b_8)$ denotes the coordinates of the point $\mathbf{b}$, then $b_i = 0$ for some $i \in I^\dagger$. Let us assume that $b_i = 0$ for some $i \in \{1, 7\}$. Then using the function $f_i^\dagger$, we conclude that $b_4 + b_5 + b_6 = 1$. Because otherwise $f_i^\dagger$ would tend to infinity through the sequence $(\mathbf{x}_n)$. But it is supposed to be tending to an infimum. Therefore, we must have $b_j = 0$ for every $j \in \{2, 3, 8\}$. Then, we get that $b_4 + b_5 + b_6 + b_7 + b_8 = 1$ and $b_1 + b_2 + b_3 + b_7 + b_8 = 1$ and $b_1 + b_2 + b_3 = 1$. Because otherwise $f_j^\dagger$ would tend to infinity through the sequence $(\mathbf{x}_n)$ when it is supposed to tend to an infimum. In any case, we obtain that $b_4 + b_5 + b_6 + b_i > 1$ for some $i \in \{1, 2, 3, 7, 8\}$. This is a contradiction.

Let us assume that $b_i = 0$ for some $i \in \{2, 6\}$, then we get $b_4 + b_5 + b_6 + b_7 + b_8 = 1$ using the function $f_i^\dagger$. Because otherwise $f_i^\dagger$ would tend to infinity through the sequence

$(\mathbf{x}_n)$. But it is supposed to be tending to an infimum. Therefore, we must have $b_j = 0$ for every $j \in \{1, 3, 4, 5, 7, 8\}$. We already showed above that $b_1 \neq 0$, a contradiction. Let us assume that $b_i = 0$ for some $i \in \{3, 5\}$. Since the function $f_i^\dagger$ cannot tend to infinity through the sequence $(\mathbf{x}_n)$, we conclude that $b_1 + b_2 + b_3 + b_7 + b_8 = 1$. This means that $b_j = 0$ for every $j \in \{4, 5, 6\}$. Then, we get that $b_4 + b_5 + b_6 + b_7 + b_8 = 1$ and $b_1 + b_2 + b_3 + b_7 + b_8 = 1$ and $b_1 + b_2 + b_3 = 1$. Because otherwise $f_j^\dagger$ would tend to infinity through the sequence $(\mathbf{x}_n)$ when it is supposed to tend to an infimum. In any case, we obtain that $b_1 + b_2 + b_3 + b_7 + b_8 + b_i > 1$ for some $i \in \{4, 5, 6\}$, a contradiction.

Let us assume that $b_i = 0$ for some $i \in \{4, 8\}$. Using the function $f_i^\dagger$, we conclude that $b_1 + b_2 + b_3 = 1$. Otherwise, $f_i^\dagger$ would tend to infinity through the sequence $(\mathbf{x}_n)$. Therefore, we get that $b_j = 0$ for every $j \in \{5, 6, 7\}$. Since the function $f_j^\dagger$ cannot approach to infinity through the sequence $(\mathbf{x}_n)$, we obtain that $b_4 + b_5 + b_6 + b_7 + b_8 = 1$ and $b_1 + b_2 + b_3 + b_7 + b_8 = 1$ and $b_4 + b_5 + b_6 = 1$. In any case, we obtain that $b_1 + b_2 + b_3 + b_i > 1$ for some $i \in \{5, 6, 7\}$. This gives a contradiction. Hence, the sequence $(\mathbf{x}_n)$ cannot have a limit point on the boundary of Δ^7 .

We claim that there exists a positive integer n_0 so that $\mathbf{x}_n = \mathbf{x}_{n+1}$ for $n \geq n_0$. Otherwise, we would have a subsequence $(\mathbf{x}_{n_j})$ of the sequence $(\mathbf{x}_n)$ so that $\mathbf{x}_{n_j} \in \Delta_{n_j+1} - \Delta_{n_j}$ for every integer $j > 0$. Since we have $\bigcup_{n=2}^{\infty} \Delta_n = \Delta^7$, we get $F^\dagger(\mathbf{x}_{n_j}) = \alpha_{n_j} \rightarrow \infty$ as $j \rightarrow \infty$. This is a contradiction. Therefore, we have $\mathbf{x}_n = \mathbf{x}_{n+1}$ for every $n \geq n_0$. In other words, the absolute minimum $F^\dagger(\mathbf{x}_{n_0})$ of $F^\dagger$ exists on Δ^7 . Then, we obviously have $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$. In the following sections, we calculate $\alpha_* = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$.

It is possible to find an interval that contains α_* . If we have $x_i = (\sqrt{2} - 1)/2$ for $i = 1, 4, 7, 8$ and $x_i = (3 - 2\sqrt{2})/4$ for $i = 2, 3, 5, 6$, then $\bar{\mathbf{x}} = (x_i)_{i \in I^+}$ is a point in Δ^7 such that $f_i(\bar{\mathbf{x}}) = 5 + 3\sqrt{2}$ for every $i \in I^+$. Therefore, we get $F^\dagger(\bar{\mathbf{x}}) = 5 + 3\sqrt{2}$ which implies that $\alpha_* \leq 5 + 3\sqrt{2}$. To find a lower bound for α_* we consider the functions $f_7^\dagger$ and $f_8^\dagger$. It is clear that $\min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) \geq \inf_{\mathbf{x} \in \Delta^7} \{\max(f_7^\dagger(\mathbf{x}), f_8^\dagger(\mathbf{x}))\}$. We apply the following substitution $X_4 = x_4 + x_5 + x_6$, $X_1 = x_7$, $X_3 = x_1 + x_2 + x_3$, $X_2 = x_8$. Then we see that

$$f_7^\dagger(\mathbf{x}) = \frac{(1 - X_4)(1 - X_1)}{X_4 X_1}, \quad f_8^\dagger(\mathbf{x}) = \frac{(1 - X_3)(1 - X_2)}{X_3 X_2}$$

where $\sum_{i=1}^4 X_i = 1$. By Lemma 2.1, we have $\inf_{\mathbf{x} \in \Delta^3} \{\max(f_7^\dagger(\mathbf{x}), f_8^\dagger(\mathbf{x}))\} = 9$. Hence, we obtain that $9 \leq \alpha_*$. This means that the minimum value of $F^\dagger$ lies in the interval $I_\alpha = [9, 5 + 3\sqrt{2}]$.

In order to calculate α_* , we need to consider the extreme values of the functions $\{f_i^\dagger\}_{i \in I^+}$ on the simplex Δ^7 . For any parametrization $\mathbf{p}_s^7$, we have the following

commuting diagram

$$\begin{array}{ccc} & \Delta^{7,s} & \\ p_s^7 \downarrow & \searrow F^{\dagger,s} & \\ \Delta^7 & \xrightarrow{F^\dagger} & \mathbb{R} \end{array}$$

where $F^{\dagger,s}(\mathbf{x}_s) = \max_{\mathbf{x}_s \in \Delta^{7,s}} \{f_i^{\dagger,s}(\mathbf{x}_s) : i \in I^\dagger\}$. By the diagram above, we derive that $F^\dagger(\mathbf{x}^*) = F^{\dagger,s}(\mathbf{x}_s^*)$ such that $\mathbf{x}_s^* = (p_s^7)^{-1}(\mathbf{x}^*)$. We use the parametrization p_8^7 . Then we obtain that

$$\begin{aligned} f_1^{\dagger,8} &= \rho_2 \sigma_1, & f_2^{\dagger,8} &= \rho^1 \sigma_2, & f_3^{\dagger,8} &= \rho^2 \sigma_3, & f_4^{\dagger,8} &= \rho_1 \sigma_4, \\ f_5^{\dagger,8} &= \rho^2 \sigma_5, & f_6^{\dagger,8} &= \rho^1 \sigma_6, & f_7^{\dagger,8} &= \rho_2 \sigma_7, & f_8^{\dagger,8} &= \rho_1 \sigma^8. \end{aligned}$$

Some of the first order partial derivatives of the functions $\{f_i^{\dagger,8}\}_{i \in I^{\dagger,8}}$ are calculated as $(f_4^{\dagger,8})_1 = -\varrho_1 \sigma_4$, $(f_1^{\dagger,8})_1 = -\rho_2 \tau_1$, $(f_7^{\dagger,8})_4 = -\varrho_2 \sigma_7$, $(f_3^{\dagger,8})_3 = -\rho^2 \tau_3$ which are all negative and $(f_8^{\dagger,8})_4 = \rho_1 \tau^8$, $(f_6^{\dagger,8})_1 = \varrho^1 \sigma_6$, $(f_2^{\dagger,8})_1 = \varrho^1 \sigma_2$, $(f_5^{\dagger,8})_4 = \varrho^2 \sigma_5$ which are all positive for every $\mathbf{x}_8 \in \Delta^{7,8}$. Therefore, the following equations $(f_j^{\dagger,8})_i = 0$ for $i \in I^\dagger$ has no solution in $\Delta^{7,8}$ for any $j \in I^\dagger$. Consequently, none of the functions $f_1^{\dagger,8}, f_2^{\dagger,8}, \dots, f_8^{\dagger,8}$ has a local extremum in $\Delta^{7,8}$.

Let us consider the set $\{f_1^\dagger(\mathbf{x}^*), f_2^\dagger(\mathbf{x}^*), \dots, f_8^\dagger(\mathbf{x}^*)\}$. If we have $f_j^\dagger(\mathbf{x}^*) \neq f_k^\dagger(\mathbf{x}^*)$ for every $j \neq k$, then the set of real numbers $\{f_1^\dagger(\mathbf{x}^*), f_2^\dagger(\mathbf{x}^*), \dots, f_8^\dagger(\mathbf{x}^*)\}$ has a unique largest element. By renumbering the functions, we can assume that $f_1^\dagger(\mathbf{x}^*)$ is the largest value. Thus, we derive that $f_1^\dagger(\mathbf{x}^*) = \alpha_*$ or $f_1^{\dagger,8}(\mathbf{x}_8^*) = \alpha_*$. By continuity, there exists a neighborhood U of $\mathbf{x}_8^*$ contained in $\Delta^{7,8}$ so that $F^{\dagger,8}(\mathbf{x}) = f_1^{\dagger,8}(\mathbf{x}_8)$ for every $\mathbf{x}_8 \in U$. Since $F^{\dagger,8}$ has a minimum at $\mathbf{x}_8^*$, then $f_1^{\dagger,8}$ has a local minimum at $\mathbf{x}_8^*$. But this contradicts with the fact that $f_1^{\dagger,8}$ has no local extremum in $\Delta^{7,8}$. Therefore, there exists two functions $f_{j_1}^\dagger, f_{j_2}^\dagger$ such that $f_{j_1}^\dagger(\mathbf{x}^*) = f_{j_2}^\dagger(\mathbf{x}^*)$ where $j_1, j_2 \in I^\dagger$ and $j_1 \neq j_2$. Hence, at least two of the functions attain the value α_* at the point $\mathbf{x}^*$. In the following sections, we give a list of all possible cases for the values of the functions $f_1^\dagger, f_2^\dagger, \dots, f_8^\dagger$ at the point $\mathbf{x}^*$.

4.1.1 The relationship between the functions $f_1^\dagger, f_2^\dagger$ and $f_3^\dagger$

Let p_s^7 be a given parametrization of Δ^7 so that $s \neq 7$ and $s \neq 8$. For every $i, j \in J_s^\dagger$, $i \neq j$, we define the vectors $\vec{u}_i^{j,s}$ in $\mathbb{R}^7$ with the ordered $x_{i_1} x_{i_2} \dots x_{i_7}$ -coordinate system so that $i_1 < i_1 < \dots < i_7$ and $i_1, i_1, \dots, i_7 \in I^\dagger - \{s\}$ where

$$\vec{u}_i^{j,s} = \begin{cases} -1 & \text{if } n = i, \\ 1 & \text{if } n = j, \\ 0 & \text{otherwise.} \end{cases}$$

Let us assume that $s \in I_1^\dagger$. We use the parametrization $\mathbf{p}_1^7$ of Δ^7 . Then we derive that

$$\begin{aligned} f_1^{\dagger,1} &= \rho_2 \sigma^1, & f_2^{\dagger,1} &= \rho_{2,3} \sigma_2, & f_3^{\dagger,1} &= \rho^2 \sigma_3, & f_4^{\dagger,1} &= \rho^{2,3} \sigma_4, \\ f_5^{\dagger,1} &= \rho^2 \sigma_5, & f_6^{\dagger,1} &= \rho_{2,3} \sigma_6, & f_7^{\dagger,1} &= \rho_2 \sigma_7, & f_8^{\dagger,1} &= \rho^{2,3} \sigma_8. \end{aligned}$$

We have the following first order partial derivatives: $(f_l^{\dagger,1})_2 = (f_l^{\dagger,1})_3$, $(f_l^{\dagger,1})_3 = 0$ for every $l \in \mathcal{I}_1^\dagger$, $(f_1^{\dagger,1})_2 = (f_1^{\dagger,1})_3$, $(f_1^{\dagger,1})_3 = \rho_2 \tau^1$, $(f_2^{\dagger,1})_2 = -\rho_{2,3} \tau_2$, $(f_2^{\dagger,1})_3 = 0$, $(f_3^{\dagger,1})_2 = 0$, and $(f_3^{\dagger,1})_3 = -\rho^2 \tau_3$ for every $\mathbf{x}_1 \in \Delta^{7,1}$. If we use the parametrization $\mathbf{p}_2^7$ of Δ^7 , then we see that

$$\begin{aligned} f_1^{\dagger,2} &= \rho_2 \sigma_1, & f_2^{\dagger,2} &= \rho_{2,3} \sigma^2, & f_3^{\dagger,2} &= \rho^2 \sigma_3, & f_4^{\dagger,2} &= \rho^{2,3} \sigma_4, \\ f_5^{\dagger,2} &= \rho^2 \sigma_5, & f_6^{\dagger,2} &= \rho_{2,3} \sigma_6, & f_7^{\dagger,2} &= \rho_2 \sigma_7, & f_8^{\dagger,2} &= \rho^{2,3} \sigma_8. \end{aligned}$$

We have the following first order partial derivatives: $(f_l^{\dagger,2})_1 = (f_l^{\dagger,2})_3$, $(f_l^{\dagger,2})_3 = 0$ for every $l \in \mathcal{I}_1^\dagger$, $(f_1^{\dagger,2})_1 = -\rho_2 \tau_1$, $(f_1^{\dagger,2})_3 = 0$, $(f_2^{\dagger,2})_1 = (f_2^{\dagger,2})_3$, $(f_2^{\dagger,2})_3 = \rho_{2,3} \tau^2$, $(f_3^{\dagger,2})_1 = 0$, and $(f_3^{\dagger,2})_3 = -\rho^2 \tau_3$ for every $\mathbf{x}_2 \in \Delta^{7,2}$. If we use the parametrization $\mathbf{p}_3^7$, then we find that

$$\begin{aligned} f_1^{\dagger,3} &= \rho_2 \sigma_1, & f_2^{\dagger,3} &= \rho_{2,3} \sigma_2, & f_3^{\dagger,3} &= \rho^2 \sigma^3, & f_4^{\dagger,3} &= \rho^{2,3} \sigma_4, \\ f_5^{\dagger,3} &= \rho^2 \sigma_5, & f_6^{\dagger,3} &= \rho_{2,3} \sigma_6, & f_7^{\dagger,3} &= \rho_2 \sigma_7, & f_8^{\dagger,3} &= \rho^{2,3} \sigma_8. \end{aligned}$$

We have the following first order partial derivatives: $(f_l^{\dagger,3})_1 = (f_l^{\dagger,3})_2$, $(f_l^{\dagger,3})_2 = 0$ for every $l \in \mathcal{I}_1^\dagger$, $(f_1^{\dagger,3})_1 = -\rho_2 \tau_1$, $(f_1^{\dagger,3})_2 = 0$, $(f_2^{\dagger,3})_1 = 0$, $(f_2^{\dagger,3})_2 = -\rho_{2,3} \tau_2$, $(f_3^{\dagger,3})_1 = (f_3^{\dagger,3})_2$ and $(f_3^{\dagger,3})_2 = \rho^2 \tau^3$ for every $\mathbf{x}_3 \in \Delta^{7,3}$.

Assume that $f_i^\dagger(\mathbf{x}^*) < \alpha_*$, $f_j^\dagger(\mathbf{x}^*) = \alpha_*$, $f_k^\dagger(\mathbf{x}^*) \leq \alpha_*$ for $i, j, k \in I_1^\dagger$ where $i \neq j$, $j \neq k$, $i \neq k$ and $f_l^\dagger(\mathbf{x}^*) \leq \alpha_*$ for $l \in \mathcal{I}_1^\dagger$. We find a point $\mathbf{z} \in \Delta^7$ so that $f_l^\dagger(\mathbf{z}) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^\dagger(\mathbf{z}) = f_j^\dagger(\mathbf{x}^*)$ for every $l \in \mathcal{I}_1^\dagger$ as follows: We use the parametrization $\mathbf{p}_7^7$. We calculate the directional derivatives of all of the functions $f_1^{\dagger,k}, f_2^{\dagger,k}, \dots, f_8^{\dagger,k}$ in the direction of the vector $\vec{u}_i^{j,k}$. We get

$$D_{\vec{u}_i^{j,k}} f_l^{\dagger,k}(\mathbf{x}_k) > 0, \quad D_{\vec{u}_i^{j,k}} f_l^{\dagger,k}(\mathbf{x}_k) = 0 \text{ for } l \in \{k\} \cup \mathcal{I}_1^\dagger, \quad D_{\vec{u}_i^{j,k}} f_j^{\dagger,k}(\mathbf{x}_k) < 0$$

for every $\mathbf{x}_k \in \Delta^{7,k}$. Then we conclude that $f_i^{\dagger,k}$ is increasing, $f_j^{\dagger,k}$ is decreasing and $f_l^{\dagger,k}$ is constant for every $l \in \{k\} \cup \mathcal{I}_1^\dagger$ on the line segment $\ell_{j,k} \cap \Delta^{7,k}$ where we let $\ell_{j,k} = \{\vec{\mathbf{x}}_k^* + t \vec{u}_i^{j,k} : t \in \mathbb{R}\}$ and $\mathbf{x}_k^* = (\mathbf{p}_k^7)^{-1}(\mathbf{x}^*)$. Since $f_i^{\dagger,k}$ is continuous on $\ell_{j,k} \cap \Delta^7$ and $f_i^{\dagger,k}(\mathbf{x}_k^*) < \alpha_*$, there exists an $\epsilon_i > 0$ so that $f_i^{\dagger,k}(\mathbf{x}_k) < \alpha_*$ for every $\mathbf{x}_k \in \ell_{j,k}^{\epsilon_i} = \{\vec{\mathbf{x}}_k^* + t \vec{u}_i^{j,k} : t \in (0, \epsilon_i)\}$. Let $\mathbf{y}_k$ be a point on $\ell_{j,k}^{\epsilon_i}$. Then we get $f_i^{\dagger,k}(\mathbf{y}_k) < \alpha_*$. Since $f_j^{\dagger,k}$ is decreasing on the line segment $\ell_{j,k}^{\epsilon_i} \subset \ell_i$ and $f_j^{\dagger,k}(\mathbf{x}_k^*) = \alpha_*$, we see that $f_j^{\dagger,k}(\mathbf{y}_k) < \alpha_*$. In other words, if we move along the line segment starting at $\mathbf{x}_k^*$ in the direction of the vector $\vec{u}_i^{j,k}$ for a sufficiently small amount, we get a point

$\mathbf{y}_k \in \Delta^{7,k}$ so that the values of $f_l^{\dagger,k}$ and $f_j^{\dagger,k}$ at $\mathbf{y}_k$ are less than α_* and the values of the functions $f_l^{\dagger,k}$ for $l \in \{k\} \cup \mathcal{I}_1^\dagger$ at the point $\mathbf{y}_k$ are the same as at the point $\mathbf{x}_k^*$, i.e. $f_l^{\dagger,k}(\mathbf{y}_k) = f_l^{\dagger,k}(\mathbf{x}_k^*)$ for every $l \in \{k\} \cup \mathcal{I}_1^\dagger$. Let us call the perturbation performed in the way we described above when $f_l^{\dagger,k}(\mathbf{x}_k) < \alpha_*$, $f_j^{\dagger,k}(\mathbf{x}_k) = \alpha_*$, $f_k^{\dagger,k}(\mathbf{x}_k) \leq \alpha_*$ and $f_l^{\dagger,k}(\mathbf{x}_k) \leq \alpha_*$ for $l \in \mathcal{I}_1^\dagger$ for some $\mathbf{x}_k \in \Delta^{7,k}$ $\text{pert}_i^{j,k}$ and the point obtained by applying the perturbation $\text{pert}_i^{j,k}$ at the point $\mathbf{x}_k$ $\text{pert}_i^{j,k}(\mathbf{x}_k)$. In the case above, we have $\mathbf{y}_k = \text{pert}_i^{j,k}(\mathbf{x}_k^*)$.

If we have $f_k^{\dagger}(\mathbf{x}^*) < \alpha_*$, then we let $\mathbf{z} = (\mathbf{p}_k^7)^{-1}(\mathbf{y}_k) \in \Delta^7$. If we have $f_k^{\dagger}(\mathbf{x}^*) = \alpha_*$, then we get $f_k^{\dagger}(\mathbf{x}^*) = f_k^{\dagger,k}(\mathbf{y}_k)$. We need to apply one more perturbation. We apply the perturbation $\text{pert}_i^{k,j}$ or $\text{pert}_j^{k,i}$ at the points $\mathbf{y}_j = \mathbf{p}_j^7((\mathbf{p}_k^7)^{-1}(\mathbf{y}_k))$ or $\mathbf{y}_i = \mathbf{p}_i^7((\mathbf{p}_k^7)^{-1}(\mathbf{y}_k))$, respectively. We obtain a point $\mathbf{z}_j = \text{pert}_i^{k,j}(\mathbf{y}_j) \in \Delta^{7,j}$ or $\mathbf{z}_i = \text{pert}_j^{k,i}(\mathbf{y}_i) \in \Delta^{7,i}$ so that $f_l^{\dagger,j}(\mathbf{z}_j) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^{\dagger,j}(\mathbf{z}_j) \leq \alpha_*$ for every $l \in \mathcal{I}_1^\dagger$ or $f_l^{\dagger,i}(\mathbf{z}_i) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^{\dagger,i}(\mathbf{z}_i) \leq \alpha_*$ for every $l \in \mathcal{I}_1^\dagger$. Then we let $\mathbf{z} = (\mathbf{p}_j^7)^{-1}(\mathbf{z}_j) \in \Delta^7$ or $\mathbf{z} = (\mathbf{p}_i^7)^{-1}(\mathbf{z}_i) \in \Delta^7$ and conclude that $f_l^{\dagger}(\mathbf{z}) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^{\dagger}(\mathbf{z}) = f_j^{\dagger}(\mathbf{x}^*)$ for every $l \in \mathcal{I}_1^\dagger$.

Let us assume that $f_i^{\dagger}(\mathbf{x}^*) < \alpha_*$, $f_j^{\dagger}(\mathbf{x}^*) < \alpha_*$, $f_k^{\dagger}(\mathbf{x}^*) = \alpha_*$ for $i, j, k \in I_1^\dagger$ where $i \neq j$, $j \neq k$, $i \neq k$ and $f_l^{\dagger}(\mathbf{x}^*) \leq \alpha_*$ for $l \in \mathcal{I}_1^\dagger$. Then we find a point $\mathbf{z} \in \Delta^7$ so that $f_l^{\dagger}(\mathbf{z}) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^{\dagger}(\mathbf{z}) = f_l^{\dagger}(\mathbf{x}^*)$ for every $l \in \mathcal{I}_1^\dagger$. We use the parametrization $\mathbf{p}_i^7$ or $\mathbf{p}_j^7$. Then we apply the perturbation $\text{pert}_i^{k,j}$ or $\text{pert}_j^{k,i}$ at the points $\mathbf{y}_j = (\mathbf{p}_j^7)^{-1}(\mathbf{x}^*)$ or $\mathbf{y}_i = (\mathbf{p}_i^7)^{-1}(\mathbf{x}^*)$, respectively. We obtain a point $\mathbf{z}_j = \text{pert}_i^{k,j}(\mathbf{y}_j) \in \Delta^{7,j}$ or $\mathbf{z}_i = \text{pert}_j^{k,i}(\mathbf{y}_i) \in \Delta^{7,i}$ so that $f_l^{\dagger,j}(\mathbf{z}_j) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^{\dagger,j}(\mathbf{z}_j) \leq \alpha_*$ for every $l \in \mathcal{I}_2^\dagger$ or $f_l^{\dagger,i}(\mathbf{z}_i) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^{\dagger,i}(\mathbf{z}_i) \leq \alpha_*$ for every $l \in \mathcal{I}_2^\dagger$. Then we let $\mathbf{z} = (\mathbf{p}_j^7)^{-1}(\mathbf{z}_j) \in \Delta^7$ or $\mathbf{z} = (\mathbf{p}_i^7)^{-1}(\mathbf{z}_i) \in \Delta^7$ and conclude that $f_l^{\dagger}(\mathbf{z}) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_l^{\dagger}(\mathbf{z}) = f_l^{\dagger}(\mathbf{x}^*)$ for every $l \in \mathcal{I}_1^\dagger$.

Finally, let us assume that $f_i^{\dagger}(\mathbf{x}^*) < \alpha_*$ for every $i \in I_1^\dagger$ and $f_j^{\dagger}(\mathbf{x}^*) \leq \alpha_*$ for $j \in \mathcal{I}_1^\dagger$. If this is the case, we know that there exists two distinct indices $j_1, j_2 \in \mathcal{I}_1^\dagger$ so that $f_{j_1}^{\dagger}$ and $f_{j_2}^{\dagger}$ take the value α_* at the point $\mathbf{x}^*$. Hence, we conclude that if one or two of the functions $f_1^{\dagger}, f_2^{\dagger}, f_3^{\dagger}$ take a value less than α_* at the point $\mathbf{x}^*$, it is possible to find a point $\mathbf{z}$ obtained by applying one or two perturbations chosen from the list

$$\mathcal{L}_1 = \{\text{pert}_i^{j,s} : i, j, s \in I_1^\dagger, i \neq j, j \neq s, i \neq s\}$$

so that all of the functions $f_1^{\dagger}, f_2^{\dagger}, f_3^{\dagger}$ take a value less than α_* at that point and the values of the functions $f_4^{\dagger}, f_5^{\dagger}, f_6^{\dagger}, f_7^{\dagger}, f_8^{\dagger}$ at that point are the same as at the point $\mathbf{x}^*$.

The relationship between the functions $f_1^\dagger$, $f_2^\dagger$ and $f_3^\dagger$ described in this section implies the following: if we assume $f_l^\dagger(\mathbf{x}^*) < \alpha_*$ for every $l \in \mathcal{I}_1^\dagger$, we see that $f_l^\dagger(\mathbf{x}^*) = \alpha_*$ for every $l \in I_1^\dagger$. Because, if we have $f_l^\dagger(\mathbf{x}^*) < \alpha_*$ for some $l \in I_1^\dagger$ and $f_l^\dagger(\mathbf{x}^*) < \alpha_*$ for every $l \in \mathcal{I}_1^\dagger$, then there exists a point $\mathbf{z} \in \Delta^7$ so that $f_j(\mathbf{z}) < \alpha_*$ for every $l \in I_1^\dagger$ and $f_j^\dagger(\mathbf{z}) = f_j^\dagger(\mathbf{x}^*) < \alpha_*$ for every $j \in \mathcal{I}_1^\dagger$. This contradicts with the minimality of α_* . Therefore, the following case

$$\text{I. } f_l^\dagger(\mathbf{x}^*) < \alpha_* \text{ for } l \in \mathcal{I}_1^\dagger \text{ and } f_l^\dagger(\mathbf{x}^*) = \alpha_* \text{ for } l \in I_1^\dagger$$

requires further investigation. We will denote the set $\{f_1^\dagger, f_2^\dagger, f_3^\dagger\}$ with $\{f_1\}$.

4.1.2 The relationship between the functions $f_4^\dagger$, $f_5^\dagger$ and $f_6^\dagger$

The functions $f_4^\dagger$, $f_5^\dagger$ and $f_6^\dagger$ are also related in a similar manner to $f_1^\dagger$, $f_2^\dagger$ and $f_3^\dagger$. If we use the parametrization $\mathfrak{p}_4^7$ of Δ^7 , then we derive that

$$\begin{aligned} f_1^{\dagger,4} &= \rho^{1,3}\sigma_1, & f_2^{\dagger,4} &= \rho^1\sigma_2, & f_3^{\dagger,4} &= \rho_{1,3}\sigma_3, & f_4^{\dagger,4} &= \rho_1\sigma^4, \\ f_5^{\dagger,4} &= \rho_{1,3}\sigma_5, & f_6^{\dagger,4} &= \rho^1\sigma_6, & f_7^{\dagger,4} &= \rho^{1,3}\sigma_7, & f_8^{\dagger,4} &= \rho_1\sigma_8. \end{aligned}$$

We have the following first order partial derivatives: $(f_l^{\dagger,4})_5 = (f_l^{\dagger,4})_6$, $(f_l^{\dagger,4})_6 = 0$ for every $l \in \mathcal{I}_2^\dagger$, $(f_4^{\dagger,4})_5 = (f_4^{\dagger,4})_6$, $(f_4^{\dagger,4})_6 = \rho_1\tau^4$, $(f_5^{\dagger,4})_5 = -\rho_{1,3}\tau_5$, $(f_5^{\dagger,4})_6 = 0$, $(f_6^{\dagger,4})_5 = 0$, and $(f_6^{\dagger,4})_6 = -\rho^1\tau_6$ for every $\mathbf{x}_4 \in \Delta^{7,4}$. If we use the parametrization $\mathfrak{p}_5^7$ of Δ^7 , then we see that

$$\begin{aligned} f_1^{\dagger,5} &= \rho^{1,3}\sigma_1, & f_2^{\dagger,5} &= \rho^1\sigma_2, & f_3^{\dagger,5} &= \rho_{1,3}\sigma_3, & f_4^{\dagger,5} &= \rho_1\sigma_4, \\ f_5^{\dagger,5} &= \rho_{1,3}\sigma^5, & f_6^{\dagger,5} &= \rho^1\sigma_6, & f_7^{\dagger,5} &= \rho^{1,3}\sigma_7, & f_8^{\dagger,5} &= \rho_1\sigma_8. \end{aligned}$$

We have the following first order partial derivatives: $(f_l^{\dagger,5})_4 = (f_l^{\dagger,5})_6$, $(f_l^{\dagger,5})_6 = 0$ for every $l \in \mathcal{I}_2^\dagger$, $(f_4^{\dagger,5})_4 = -\rho_1\tau_4$, $(f_4^{\dagger,5})_6 = 0$, $(f_5^{\dagger,5})_4 = (f_5^{\dagger,5})_6$, $(f_5^{\dagger,5})_6 = \rho_{1,3}\tau^5$, $(f_6^{\dagger,5})_4 = 0$, and $(f_6^{\dagger,5})_6 = -\rho^1\tau_6$ for every $\mathbf{x}_5 \in \Delta^{7,5}$. If we use the parametrization $\mathfrak{p}_6^7$, then we find that

$$\begin{aligned} f_1^{\dagger,6} &= \rho^{1,3}\sigma_1, & f_2^{\dagger,6} &= \rho^1\sigma_2, & f_3^{\dagger,6} &= \rho_{1,3}\sigma_3, & f_4^{\dagger,6} &= \rho_1\sigma_4, \\ f_5^{\dagger,6} &= \rho_{1,3}\sigma_5, & f_6^{\dagger,6} &= \rho^1\sigma^6, & f_7^{\dagger,6} &= \rho^{1,3}\sigma_7, & f_8^{\dagger,6} &= \rho_1\sigma_8. \end{aligned}$$

We have the following first order partial derivatives: $(f_l^{\dagger,6})_4 = (f_l^{\dagger,6})_5$, $(f_l^{\dagger,6})_5 = 0$ for every $l \in \mathcal{I}_2^\dagger$, $(f_4^{\dagger,6})_4 = -\rho_1\tau_4$, $(f_4^{\dagger,6})_5 = 0$, $(f_5^{\dagger,6})_4 = 0$, $(f_5^{\dagger,6})_5 = -\rho_{1,3}\tau_5$, $(f_6^{\dagger,6})_4 = (f_6^{\dagger,6})_5$ and $(f_6^{\dagger,6})_5 = \rho^1\tau^6$ for every $\mathbf{x}_6 \in \Delta^{7,6}$.

Assume that $f_i^\dagger(\mathbf{x}^*) < \alpha_*$, $f_j^\dagger(\mathbf{x}^*) = \alpha_*$, $f_k^\dagger(\mathbf{x}^*) \leq \alpha_*$ for $i, j, k \in I_2^\dagger$ where $i \neq j$, $j \neq k$, $i \neq k$ and $f_l^\dagger(\mathbf{x}^*) \leq \alpha_*$ for $l \in \mathcal{I}_2^\dagger$. We find a point $\mathbf{w} \in \Delta^7$ so that $f_l^\dagger(\mathbf{w}) < \alpha_*$ for every

$l \in I_2^\dagger$ and $f_l^\dagger(\mathbf{w}) = f_l^\dagger(\mathbf{x}^*)$ for every $l \in I_2^\dagger$ as follows: We use the parametrization $\mathbf{p}_k^7$. We calculate the directional derivatives of all of the functions $f_1^{\dagger,k}, f_2^{\dagger,k}, \dots, f_8^{\dagger,k}$ in the direction of the vector $\vec{u}_i^{j,k}$. We get

$$D_{\vec{u}_i^{j,k}} f_i^{\dagger,k}(\mathbf{x}_k) > 0, D_{\vec{u}_i^{j,k}} f_l^{\dagger,k}(\mathbf{x}_k) = 0 \text{ for } l \in \{k\} \cup I_2^\dagger, D_{\vec{u}_i^{j,k}} f_j^{\dagger,k}(\mathbf{x}_k) < 0$$

for every $\mathbf{x}_k \in \Delta^{7,k}$. Then we conclude that $f_i^{\dagger,k}$ is increasing, $f_j^{\dagger,k}$ is decreasing and $f_l^{\dagger,k}$ is constant for every $l \in \{k\} \cup I_2^\dagger$ on the line segment $\ell_{j,k} \cap \Delta^{7,k}$ where we let $\ell_{j,k} = \{\mathbf{x}_k^* + t\vec{u}_i^{j,k} : t \in \mathbb{R}\}$ and $\mathbf{x}_k^* = (\mathbf{p}_k^7)^{-1}(\mathbf{x}^*)$. Since $f_i^{\dagger,k}$ is continuous on $\ell_{j,k} \cap \Delta^7$ and $f_i^{\dagger,k}(\mathbf{x}_k^*) < \alpha_*$, there exists an $\epsilon_i > 0$ so that $f_i^{\dagger,k}(\mathbf{x}_k) < \alpha_*$ for every $\mathbf{x}_k \in \ell_{j,k}^{\epsilon_i} = \{\mathbf{x}_k^* + t\vec{u}_i^{j,k} : t \in (0, \epsilon_i)\}$. Let $\mathbf{y}_k$ be a point on $\ell_{j,k}^{\epsilon_i}$. Then we get $f_i^{\dagger,k}(\mathbf{y}_k) < \alpha_*$. Since $f_j^{\dagger,k}$ is decreasing on the line segment $\ell_{j,k}^{\epsilon_i} \subset \ell_j$ and $f_j^{\dagger,k}(\mathbf{x}_k^*) = \alpha_*$, we see that $f_j^{\dagger,k}(\mathbf{y}_k) < \alpha_*$. In other words, if we move along the line segment starting at $\mathbf{x}_k^*$ in the direction of the vector $\vec{u}_i^{j,k}$ for a sufficiently small amount, we get a point $\mathbf{y}_k \in \Delta^{7,k}$ so that the values of $f_i^{\dagger,k}$ and $f_j^{\dagger,k}$ at $\mathbf{y}_k$ are less than α_* and the values of the functions $f_l^{\dagger,k}$ for $l \in \{k\} \cup I_2^\dagger$ at the point $\mathbf{y}_k$ are the same as at the point $\mathbf{x}_k^*$, i.e. $f_l^{\dagger,k}(\mathbf{y}_k) = f_l^{\dagger,k}(\mathbf{x}_k^*)$ for every $l \in \{k\} \cup I_2^\dagger$. Let us call the perturbation performed in the way we described above when $f_i^{\dagger,k}(\mathbf{x}_k) < \alpha_*$, $f_j^{\dagger,k}(\mathbf{x}_k) = \alpha_*$, $f_k^{\dagger,k}(\mathbf{x}_k) \leq \alpha_*$ and $f_l^{\dagger,k}(\mathbf{x}_k) \leq \alpha_*$ for $l \in I_2^\dagger$ for some $\mathbf{x}_k \in \Delta^{7,k}$ $\text{pert}_i^{j,k}$ and the point obtained by applying the perturbation $\text{pert}_i^{j,k}$ at the point $\mathbf{x}_k$ $\text{pert}_i^{j,k}(\mathbf{x}_k)$. In the case above, we have $\mathbf{y}_k = \text{pert}_i^{j,k}(\mathbf{x}_k^*)$.

If we have $f_k^{\dagger}(\mathbf{x}^*) < \alpha_*$, then we let $\mathbf{w} = (\mathbf{p}_k^7)^{-1}(\mathbf{y}_k) \in \Delta^7$. If we have $f_k^{\dagger}(\mathbf{x}^*) = \alpha_*$, then we get $f_k^{\dagger}(\mathbf{x}^*) = f_k^{\dagger,k}(\mathbf{y}_k)$. We need to apply one more perturbation. We apply the perturbation $\text{pert}_i^{k,j}$ or $\text{pert}_j^{k,i}$ at the points $\mathbf{y}_j = \mathbf{p}_j^7((\mathbf{p}_k^7)^{-1}(\mathbf{y}_k))$ or $\mathbf{y}_i = \mathbf{p}_i^7((\mathbf{p}_k^7)^{-1}(\mathbf{y}_k))$, respectively. We obtain a point $\mathbf{w}_j = \text{pert}_i^{k,j}(\mathbf{y}_j) \in \Delta^{7,j}$ or $\mathbf{w}_i = \text{pert}_j^{k,i}(\mathbf{y}_i) \in \Delta^{7,i}$ so that $f_l^{\dagger,j}(\mathbf{w}_j) < \alpha_*$ for every $l \in I_2^\dagger$ and $f_l^{\dagger,j}(\mathbf{w}_j) \leq \alpha_*$ for every $l \in I_2^\dagger$ or $f_l^{\dagger,i}(\mathbf{w}_i) < \alpha_*$ for every $l \in I_2^\dagger$ and $f_l^{\dagger,i}(\mathbf{w}_i) \leq \alpha_*$ for every $l \in I_2^\dagger$. Then we let $\mathbf{w} = (\mathbf{p}_j^7)^{-1}(\mathbf{w}_j) \in \Delta^7$ or $\mathbf{w} = (\mathbf{p}_i^7)^{-1}(\mathbf{w}_i) \in \Delta^7$ and conclude that $f_l^{\dagger}(\mathbf{w}) < \alpha_*$ for every $l \in I_2^\dagger$ and $f_l^{\dagger}(\mathbf{w}) = f_l^{\dagger}(\mathbf{x}^*)$ for every $l \in I_2^\dagger$.

Let us assume that $f_i^{\dagger}(\mathbf{x}^*) < \alpha_*$, $f_j^{\dagger}(\mathbf{x}^*) < \alpha_*$, $f_k^{\dagger}(\mathbf{x}^*) = \alpha_*$ for $i, j, k \in I_2^\dagger$ where $i \neq j$, $j \neq k$, $i \neq k$ and $f_l^{\dagger}(\mathbf{x}^*) \leq \alpha_*$ for $l \in I_1^\dagger$. Then we find a point $\mathbf{w} \in \Delta^7$ so that $f_l^{\dagger}(\mathbf{w}) < \alpha_*$ for every $l \in I_2^\dagger$ and $f_l^{\dagger}(\mathbf{w}) = f_l^{\dagger}(\mathbf{x}^*)$ for every $l \in I_2^\dagger$. We use the parametrization $\mathbf{p}_i^7$ or $\mathbf{p}_j^7$. Then we apply the perturbation $\text{pert}_i^{k,j}$ or $\text{pert}_j^{k,i}$ at the points $\mathbf{y}_j = (\mathbf{p}_j^7)^{-1}(\mathbf{x}^*)$ or $\mathbf{y}_i = (\mathbf{p}_i^7)^{-1}(\mathbf{x}^*)$, respectively. We obtain a point $\mathbf{w}_j = \text{pert}_i^{k,j}(\mathbf{y}_j) \in \Delta^{7,j}$ or $\mathbf{w}_i = \text{pert}_j^{k,i}(\mathbf{y}_i) \in \Delta^{7,i}$ so that $f_l^{\dagger,j}(\mathbf{w}_j) < \alpha_*$ for every $l \in I_2^\dagger$

and $f_l^{\dagger,j}(\mathbf{w}_j) \leq \alpha_*$ for every $l \in \mathcal{I}_2^{\dagger}$ or $f_l^{\dagger,i}(\mathbf{w}_i) < \alpha_*$ for every $l \in I_2^{\dagger}$ and $f_l^{\dagger,i}(\mathbf{w}_i) \leq \alpha_*$ for every $l \in \mathcal{I}_2^{\dagger}$. Then we let $\mathbf{w} = (\mathbf{p}_j^7)^{-1}(\mathbf{w}_j) \in \Delta^7$ or $\mathbf{w} = (\mathbf{p}_i^7)^{-1}(\mathbf{w}_i) \in \Delta^7$ and conclude that $f_l^{\dagger}(\mathbf{w}) < \alpha_*$ for every $l \in I_1^{\dagger}$ and $f_l^{\dagger}(\mathbf{w}) = f_j^{\dagger}(\mathbf{x}^*)$ for every $l \in \mathcal{I}_2^{\dagger}$.

Finally, let us assume that $f_i^{\dagger}(\mathbf{x}^*) < \alpha_*$ for every $i \in I_2^{\dagger}$ and $f_j^{\dagger}(\mathbf{x}^*) \leq \alpha_*$ for $j \in \mathcal{I}_2^{\dagger}$. If this is the case, we know that there exists two distinct indices $j_1, j_2 \in \mathcal{I}_2^{\dagger}$ so that $f_{j_1}^{\dagger}$ and $f_{j_2}^{\dagger}$ take the value α_* at the point $\mathbf{x}^*$. Hence, we conclude that if one or two of the functions $f_1^{\dagger}, f_2^{\dagger}, f_3^{\dagger}$ take a value less than α_* at the point $\mathbf{x}^*$, it is possible to find a point $\mathbf{w}$ obtained by applying one or two perturbations chosen from the list

$$\mathcal{L}_2 = \{\text{pert}_i^{j,s} : i, j, s \in I_2^{\dagger}, i \neq j, j \neq s, i \neq s\}$$

so that all of the functions $f_1^{\dagger}, f_2^{\dagger}, f_3^{\dagger}$ take a value less than α_* at $\mathbf{w}$ and the values of the functions $f_4^{\dagger}, f_5^{\dagger}, f_6^{\dagger}, f_7^{\dagger}, f_8^{\dagger}$ at $\mathbf{w}$ are the same as at the point $\mathbf{x}^*$.

The relationship between the functions $f_1^{\dagger}, f_2^{\dagger}$ and $f_3^{\dagger}$ described in this section implies the following: if we assume $f_l^{\dagger}(\mathbf{x}^*) < \alpha_*$ for every $l \in \mathcal{I}_2^{\dagger}$, we see that $f_l^{\dagger}(\mathbf{x}^*) = \alpha_*$ for every $l \in I_2^{\dagger}$. Because, if we have $f_l^{\dagger}(\mathbf{x}^*) < \alpha_*$ for some $l \in I_2^{\dagger}$ and $f_l^{\dagger}(\mathbf{x}^*) < \alpha_*$ for every $l \in \mathcal{I}_2^{\dagger}$, then there exists a point $\mathbf{w} \in \Delta^7$ so that $f_j(\mathbf{w}) < \alpha_*$ for every $l \in I_2^{\dagger}$ and $f_j^{\dagger}(\mathbf{w}) = f_j^{\dagger}(\mathbf{x}^*) < \alpha_*$ for every $j \in \mathcal{I}_2^{\dagger}$. This contradicts with the minimality of α_* . Therefore, the following case

$$I_1. \quad f_l^{\dagger}(\mathbf{x}^*) < \alpha_* \text{ for } l \in \mathcal{I}_2^{\dagger} \text{ and } f_l^{\dagger}(\mathbf{x}^*) = \alpha_* \text{ for } l \in I_2^{\dagger}$$

requires further investigation. We will denote the set $\{f_4^{\dagger}, f_5^{\dagger}, f_6^{\dagger}\}$ with $\{f_2\}$.

4.1.3 The relationship between the functions $f_7^{\dagger}$ and $f_8^{\dagger}$

Let $\mathbf{p}_s^7$ be a given parametrization of Δ^7 so that $s = 7$ or $s = 8$. Let us consider the vectors $\vec{u}_i^{j,s}$ in $\mathbb{R}^7$ with the ordered $x_{i_1}x_{i_2} \cdots x_{i_7}$ -coordinate system defined by

$$\vec{u}_7^{8,8} = \begin{cases} -1 & \text{if } n = 7, \\ 0 & \text{otherwise,} \end{cases} \quad \text{and} \quad \vec{u}_8^{7,7} = \begin{cases} -1 & \text{if } n = 8, \\ 0 & \text{otherwise,} \end{cases}$$

where $i_1 < i_1 < \cdots < i_7$ and $i_1, i_1, \dots, i_7 \in I^{\dagger} - \{s\}$. If we use the parametrization $\mathbf{p}_7^7$ of Δ^7 , then we obtain the formulas below for the displacements functions:

$$\begin{aligned} f_1^{\dagger,7} &= \rho_2 \sigma_1, & f_2^{\dagger,7} &= \rho^1 \sigma_2, & f_3^{\dagger,7} &= \rho^2 \sigma_3, & f_4^{\dagger,7} &= \rho_1 \sigma_4, \\ f_5^{\dagger,7} &= \rho^2 \sigma_5, & f_6^{\dagger,7} &= \rho^1 \sigma_6, & f_7^{\dagger,7} &= \rho_2 \sigma^7, & f_8^{\dagger,7} &= \rho_1 \sigma_8. \end{aligned}$$

We have the first order partial derivatives: $(f_l^{\dagger,7})_8 = 0$ for every $l \in \mathcal{I}_3^\dagger$, $(f_7^{\dagger,7})_8 = \rho_2 \tau^7$, and $(f_8^{\dagger,7})_8 = -\rho_1 \tau_8$ for every $\mathbf{x}_7 \in \Delta^{7,7}$. If we use the parametrization $\mathbf{p}_7^7$ of Δ^7 , then we see that

$$\begin{aligned} f_1^{\dagger,8} &= \rho_2 \sigma_1, & f_2^{\dagger,8} &= \rho^1 \sigma_2, & f_3^{\dagger,8} &= \rho^2 \sigma_3, & f_4^{\dagger,8} &= \rho_1 \sigma_4, \\ f_5^{\dagger,8} &= \rho^2 \sigma_5, & f_6^{\dagger,8} &= \rho^1 \sigma_6, & f_7^{\dagger,8} &= \rho_2 \sigma_7, & f_8^{\dagger,8} &= \rho_1 \sigma^8. \end{aligned}$$

We have the following first order partial derivatives: $(f_l^{\dagger,8})_7 = 0$ for every $l \in \mathcal{I}_3^\dagger$, $(f_7^{\dagger,8})_7 = -\rho_2 \tau_7$, and $(f_8^{\dagger,8})_8 = \rho_1 \tau^8$ for every $\mathbf{x}_8 \in \Delta^{7,8}$.

Assume that $f_i^\dagger(\mathbf{x}^*) < \alpha_*$ and $f_j^\dagger(\mathbf{x}^*) = \alpha_*$ for $i, j \in I_3^\dagger$ where $i \neq j$ and $f_l^\dagger(\mathbf{x}^*) \leq \alpha_*$ for $l \in \mathcal{I}_3^\dagger$. We find a point $\mathbf{t} \in \Delta^7$ so that $f_l^\dagger(\mathbf{t}) < \alpha_*$ for every $l \in I_3^\dagger$ and $f_l^\dagger(\mathbf{t}) = f_l^\dagger(\mathbf{x}^*)$ for every $l \in \mathcal{I}_3^\dagger$ as follows: We use the parametrization $\mathbf{p}_j^7$. We calculate the directional derivatives of all of the functions $f_1^{\dagger,j}, f_2^{\dagger,j}, \dots, f_8^{\dagger,j}$ in the direction of the vector $\bar{u}_i^{\dagger,j}$. We get

$$D_{\bar{u}_i^{\dagger,j}} f_i^{\dagger,j}(\mathbf{x}_j) > 0, \quad D_{\bar{u}_i^{\dagger,j}} f_l^{\dagger,j}(\mathbf{x}_j) = 0 \text{ for } l \in \{j\} \cup \mathcal{I}_3^\dagger, \quad D_{\bar{u}_i^{\dagger,j}} f_j^{\dagger,j}(\mathbf{x}_j) < 0$$

for every $\mathbf{x}_j \in \Delta^{7,j}$. Then we conclude that $f_i^{\dagger,j}$ is increasing, $f_j^{\dagger,j}$ is decreasing and $f_l^{\dagger,j}$ is constant for every $l \in \{j\} \cup \mathcal{I}_3^\dagger$ on the line segment $\ell_{j,j} \cap \Delta^{7,j}$ where we let $\ell_{j,j} = \{\bar{\mathbf{x}}_j^* + t \bar{u}_i^{\dagger,j} : t \in \mathbb{R}\}$ and $\bar{\mathbf{x}}_j^* = (\mathbf{p}_j^7)^{-1}(\mathbf{x}^*)$. Since $f_i^{\dagger,j}$ is continuous on $\ell_{j,j} \cap \Delta^7$ and $f_i^{\dagger,j}(\bar{\mathbf{x}}_j^*) < \alpha_*$, there exists an $\epsilon_i > 0$ so that $f_i^{\dagger,j}(\mathbf{x}_j) < \alpha_*$ for every $\mathbf{x}_j \in \ell_{j,j}^{\epsilon_i} = \{\bar{\mathbf{x}}_j^* + t \bar{u}_i^{\dagger,j} : t \in (0, \epsilon_i)\}$. Let $\mathbf{t}_j$ be a point on $\ell_{j,j}^{\epsilon_i}$. Then we get $f_i^{\dagger,j}(\mathbf{t}_j) < \alpha_*$. Since $f_j^{\dagger,j}$ is decreasing on the line segment $\ell_{j,j}^{\epsilon_i} \subset \ell_i$ and $f_j^{\dagger,j}(\bar{\mathbf{x}}_j^*) = \alpha_*$, we see that $f_j^{\dagger,j}(\mathbf{t}_j) < \alpha_*$. In other words, if we move along the line segment starting at $\bar{\mathbf{x}}_j^*$ in the direction of the vector $\bar{u}_i^{\dagger,j}$ for a sufficiently small amount, we get a point $\mathbf{t}_j \in \Delta^{7,j}$ so that the values of $f_i^{\dagger,j}$ and $f_j^{\dagger,j}$ at $\mathbf{t}_j$ are less than α_* and the values of the functions $f_l^{\dagger,j}$ for $l \in \{j\} \cup \mathcal{I}_3^\dagger$ at the point $\mathbf{t}_j$ are the same as at the point $\bar{\mathbf{x}}_j^*$, i.e. $f_l^{\dagger,j}(\mathbf{t}_j) = f_l(\bar{\mathbf{x}}_j^*)$ for every $l \in \{j\} \cup \mathcal{I}_3^\dagger$. Let us call the perturbation performed in the way we described above when $f_i^{\dagger,j}(\mathbf{x}_j) < \alpha_*$, $f_j^{\dagger,j}(\mathbf{x}_j) = \alpha_*$, $f_j^{\dagger,j}(\mathbf{x}_j) \leq \alpha_*$ and $f_l^{\dagger,j}(\mathbf{x}_j) \leq \alpha_*$ for $l \in \mathcal{I}_3^\dagger$ for some $\mathbf{x}_j \in \Delta^{7,j}$ $\text{pert}_i^{\dagger,j}$ and the point obtained by applying the perturbation $\text{pert}_i^{\dagger,j}$ at the point $\mathbf{x}_j$ $\text{pert}_i^{\dagger,j}(\mathbf{x}_j)$. In the case above, we have $\mathbf{t}_j = \text{pert}_i^{\dagger,j}(\bar{\mathbf{x}}_j^*)$. Then we let $\mathbf{t} = (\mathbf{p}_j^7)^{-1}(\mathbf{t}_j) \in \Delta^7$ and conclude that $f_l^\dagger(\mathbf{t}) < \alpha_*$ for every $l \in I_3^\dagger$ and $f_l^\dagger(\mathbf{t}) = f_l^\dagger(\mathbf{x}^*)$ for every $l \in \mathcal{I}_3^\dagger$.

Finally, let us assume that $f_l^\dagger(\mathbf{x}^*) < \alpha_*$ for every $l \in I_3^\dagger$ and $f_l^\dagger(\mathbf{x}^*) \leq \alpha_*$ for $l \in \mathcal{I}_3^\dagger$. If this is the case, we know that there exists two distinct indices $j_1, j_2 \in \mathcal{I}_3^\dagger$ so that $f_{j_1}^\dagger$ and $f_{j_2}^\dagger$ take the value α_* at the point $\mathbf{x}^*$. Hence, we conclude that if one of the functions $f_j^\dagger$

or $f_8^\dagger$ take a value less than α_* at the point $\mathbf{x}^*$, it is possible to find a point $\mathbf{t}$ obtained by applying a perturbations chosen from the list

$$\mathcal{L}_3 = \{pert_7^{8,8}, pert_8^{7,7}\}$$

so that both of the functions $f_7^\dagger$ and $f_8^\dagger$ take a value less than α_* at $\mathbf{t}$ and the values of the functions $f_1^\dagger, f_2^\dagger, f_3^\dagger, f_4^\dagger, f_5^\dagger, f_6^\dagger$ at $\mathbf{t}$ are the same as at $\mathbf{x}^*$.

The relationship between the functions $f_7^\dagger$ and $f_8^\dagger$ described in this section implies the following: If we assume that $f_l^\dagger(\mathbf{x}^*) < \alpha_*$ for $l \in \mathcal{I}_3^\dagger$, we see that $f_i^\dagger(\mathbf{x}^*) = \alpha_*$ and $f_j^\dagger(\mathbf{x}^*) = \alpha_*$ for every $i, j \in I_3^\dagger$ where $i \neq j$. Because if we have $f_i^\dagger(\mathbf{x}^*) = \alpha_*$ and $f_l^\dagger(\mathbf{x}^*) < \alpha_*$ for every $l \in \mathcal{I}_3^\dagger \cup (I_3^\dagger - \{i\})$, then there exists a point $\mathbf{t} \in \Delta^7$ so that $f_i^\dagger(\mathbf{t}) < \alpha_*$, $f_j^\dagger(\mathbf{t}) < \alpha_*$ and $f_l(\mathbf{t}) = f_l(\mathbf{x}^*) < \alpha_*$ for every $l \in \mathcal{I}_3^\dagger$. This contradicts with the minimality of α_* . Therefore, the following case

$$\text{II. } f_l^\dagger(\mathbf{x}^*) < \alpha_* \text{ for } l \in \mathcal{I}_3^\dagger \text{ and } f_l(\mathbf{x}^*) = \alpha_* \text{ for } l \in I_3^\dagger$$

needs further investigation. We will denote the set $\{f_7^\dagger, f_8^\dagger\}$ with $\{f_3\}$. Note that Case II can be obtained directly from the fact that there exists two functions $f_{j_1}^\dagger, f_{j_2}^\dagger$ such that $f_{j_1}^\dagger(\mathbf{x}^*) = f_{j_2}^\dagger(\mathbf{x}^*)$ where $j_1, j_2 \in I^\dagger$ and $j_1 \neq j_2$. If we assume that $f_l^\dagger(\mathbf{x}^*) < \alpha_*$ for every $l \in \mathcal{I}_3^\dagger$, then it is clear that $j_1 = 7$ and $j_2 = 8$.

In the rest of this paper, we will drop the superscript $\dagger$ from the displacement functions $\{f_i^\dagger\}_{i \in I^\dagger}$, $\{g_j^\dagger\}_{j \in I^\dagger - \{7,8\}}$ and some of the index sets. We will use the expressions $f_i^\dagger(\mathbf{x}) = f_i(\mathbf{x})$, $g_j^\dagger(\mathbf{x}) = g_j(\mathbf{x})$ for every $\mathbf{x} \in \Delta^7$ and for every $i \in I^\dagger$ and $j \in I^\dagger - \{7,8\}$. Similarly, we will have $I_i^\dagger = I_i$ for $i = 1, 2, 3, 4, 5$ and $J_i^\dagger = J_i$ for $i \in I^\dagger$ and $I^\dagger = I$.

4.1.4 The relationship between the sets $\{f_1\}$, $\{f_2\}$ and $\{f_3\}$

Case I_1 in §§4.1.2 is equivalent to Case I in §§4.1.1 in the sense that the existence of a point in Δ^7 which satisfies Case I implies the existence of a point in Δ^7 which satisfies Case I_1 . Similarly, the existence of a point in Δ^7 which satisfies Case I_1 implies the existence of a point in Δ^7 which satisfies Case I.

Let us consider the transformation $L: \Delta^7 \rightarrow \Delta^7$ defined by $x_1 \leftrightarrow x_4$, $x_2 \leftrightarrow x_5$, $x_3 \leftrightarrow x_6$, and $x_7 \leftrightarrow x_8$. We see that all the functions $\{f_i\}_{i \in I}$ are preserved and $f_1(\mathbf{x}^*) = f_4(L(\mathbf{x}^*))$, $f_2(\mathbf{x}^*) = f_5(L(\mathbf{x}^*))$, $f_3(\mathbf{x}^*) = f_6(L(\mathbf{x}^*))$, $f_7(\mathbf{x}^*) = f_8(L(\mathbf{x}^*))$. Hence, if we have $f_j(\mathbf{x}^*) < \alpha_*$ for $j \in \mathcal{I}_2$ and $f_j(\mathbf{x}^*) = \alpha_*$ for $j \in I_1$, then we have $f_j(L(\mathbf{x}^*)) = \alpha_*$ for $j \in I_2$ and $f_j(L(\mathbf{x}^*)) < \alpha_*$ for $j \in \mathcal{I}_2$. Similarly, if we have $f_j(\mathbf{x}^*) = \alpha_*$ for $j \in I_2$ and $f_j(\mathbf{x}^*) < \alpha_*$ for $j \in \mathcal{I}_2$, then we have $f_j(L(\mathbf{x}^*)) = \alpha_*$ for

$j \in I_1$ and $f_j(L(\mathbf{x}^*)) < \alpha_*$ for $j \in \mathcal{I}_2$. Therefore, we consider Cases I and I_1 as one case.

The relationships between the functions $\{f_i: i \in I\}$ we presented above show that if we have $f_l(\mathbf{x}^*) < \alpha_*$ for some $l \in I_1$, we can find a point $\mathbf{z} \in \Delta^7$ using perturbations chosen from the list $\mathcal{L}_1$ so that $f_l(\mathbf{z}) < \alpha_*$ for every $l \in I_1$ and $F^\dagger(\mathbf{z}) = \alpha_*$. Similarly, if we have $f_l(\mathbf{x}^*) < \alpha_*$ for some $l \in I_2$, then we can find a point $\mathbf{w} \in \Delta^7$ using perturbations from the list $\mathcal{L}_2$ so that $f_l(\mathbf{w}) < \alpha_*$ for every $l \in I_2$ and $F^\dagger(\mathbf{w}) = \alpha_*$. If there exists an $l \in I_3$ so that $f_l(\mathbf{x}^*) < \alpha_*$, then we can find a point $\mathbf{t} \in \Delta^7$ using one or two perturbations from the list $\mathcal{L}_3$ so that $f_l(\mathbf{t}) < \alpha_*$ for every $l \in I_3$ and $F^\dagger(\mathbf{t}) = \alpha_*$.

Let assume that $f_l(\mathbf{x}^*) < \alpha_*$ for every $l \in I_3$. If there exists an $l_1 \in I_1$ such that $f_{l_1}(\mathbf{x}^*) < \alpha_*$, then we have $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_2$. Otherwise, we apply perturbations chosen from the list $\mathcal{L}_2$. We obtain a point $\mathbf{w} \in \Delta^7$ so that $f_l(\mathbf{w}) < \alpha_*$ for every $l \in I_2$. Then we apply one or two perturbations chosen from the list $\mathcal{L}_1$. We find a point $\mathbf{z} \in \Delta^7$ so that $f_l(\mathbf{z}) < \alpha_*$ for every $l \in I$. This contradicts with the minimality of α_* . Therefore, we see that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_2$. But then by applying permutations from the list $\mathcal{L}_1$, we obtain Case I_1 , or equivalently Case I.

If there exists an $l_2 \in I_2$ such that $f_{l_2}(\mathbf{x}^*) < \alpha_*$, then we have $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_1$. Otherwise, we apply perturbations chosen from the list $\mathcal{L}_1$. We obtain a point $\mathbf{z} \in \Delta^7$ so that $f_l(\mathbf{z}) < \alpha_*$ for every $l \in I_1$. Then we apply one or two perturbations chosen from the list $\mathcal{L}_2$. We find a point $\mathbf{w} \in \Delta^7$ so that $f_l(\mathbf{w}) < \alpha_*$ for every $l \in I$. This contradicts with the minimality of α_* . Thus, we see that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_1$. But then this is Case I. Therefore, a new case arises if we assume that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in \mathcal{I}_3$. Let us numerate this case as

III. $f_j(\mathbf{x}^*) = \alpha_*$ for $j \in \mathcal{I}_3$ and $f_j(\mathbf{x}^*) < \alpha_*$ for $j \in I_3$.

Let us assume that $f_l(\mathbf{x}^*) < \alpha_*$ for every $l \in I_2$. If there exists $l_1 \in I_1$ so that $f_{l_1}(\mathbf{x}^*) < \alpha_*$, then we claim that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_3$. Because if there exists $l_3 \in I_3$ such that $f_{l_3}(\mathbf{x}^*) < \alpha_*$, we apply a perturbation from the list $\mathcal{L}_3$. We obtain a point $\mathbf{t} \in \Delta^7$ so that $f_l(\mathbf{t}) < \alpha_*$ for every $l \in I_3$. Then we apply perturbations from the list $\mathcal{L}_1$. We get a point $\mathbf{z} \in \Delta^7$ so that $f_l(\mathbf{z}) < \alpha_*$ for every $l \in I$. This contradicts with the minimality of α_* . Therefore, we see that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_3$. But this gives Case II.

If there exists a $l_3 \in I_3$ such that $f_{l_3}(\mathbf{x}^*) < \alpha_*$, then we have $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_1$. Because, otherwise, we apply perturbations from the list $\mathcal{L}_1$ and obtain a point $\mathbf{z} \in \Delta^7$ so that $f_k(\mathbf{z}) < \alpha_*$ for every $l \in I_1$. Then we apply perturbations from the list $\mathcal{L}_3$. We find a point $\mathbf{t} \in \Delta^7$ so that $f_l(\mathbf{t}) < \alpha_*$ for every $l \in I$. This contradicts with

the minimality of α_* . Hence, we conclude that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_1$. But then this is Case I. Therefore, another new case arises if we assume that $f_l(\mathbf{x}^*) = \alpha_*$ for every $k \in \mathcal{I}_2$. Specifically, we obtain the following case

IV. $f_j(\mathbf{x}^*) = \alpha_*$ for $j \in \mathcal{I}_2$ and $f_j(\mathbf{x}^*) < \alpha_*$ for $j \in I_2$,

which needs further investigation.

Let us assume that $f_l(\mathbf{x}^*) < \alpha_*$ for every $l \in I_1$. If there exists $l_3 \in I_3$ so that $f_{l_3}(\mathbf{x}^*) < \alpha_*$, then we claim that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_2$. Because if there exists $l_2 \in I_2$ such that $f_{l_2}(\mathbf{x}^*) < \alpha_*$, we apply a perturbation from the list $\mathcal{L}_2$. We obtain a point $\mathbf{w} \in \Delta^7$ so that $f_l(\mathbf{w}) < \alpha_*$ for every $l \in I_2$. Then we apply perturbations from the list $\mathcal{L}_3$. We find a point $\mathbf{t} \in \Delta^7$ so that $f_l(\mathbf{t}) < \alpha_*$ for every $l \in I$. This contradicts with the minimality of α_* . Thus, we see that $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_2$. But then this gives Case I_1 .

If there exists an $l_2 \in I_2$ such that $f_{l_2}(\mathbf{x}^*) < \alpha_*$, then we have $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_3$. Otherwise, we apply perturbations from the list $\mathcal{L}_3$. We obtain a point $\mathbf{t} \in \Delta^7$ so that $f_l(\mathbf{t}) < \alpha_*$ for every $l \in I_3$. Then we apply perturbations from the list $\mathcal{L}_2$. We get a point $\mathbf{w} \in \Delta^7$ so that $f_l(\mathbf{w}) < \alpha_*$ for every $l \in I$. This contradicts with the minimality of α_* . Therefore, we conclude $f_l(\mathbf{x}^*) = \alpha_*$ for every $l \in I_3$. But then this is Case II. Therefore, another new case arises if we assume that $f_k(\mathbf{x}^*) = \alpha_*$ for every $k \in \mathcal{I}_2$. In particular, we obtain the case

IV₁. $f_j(\mathbf{x}^*) = \alpha_*$ for $j \in \mathcal{I}_1$ and $f_j(\mathbf{x}^*) < \alpha_*$ for $j \in I_1$.

In fact, Case IV₁ is not a separate case. Because if we have $f_l(\mathbf{x}^*) = \alpha_*$ for $l \in \mathcal{I}_2$ and $f_l(\mathbf{x}^*) < \alpha_*$ for $l \in I_1$, then we see that $f_l(L(\mathbf{x}^*)) < \alpha_*$ for $l \in I_2$ and $f_l(L(\mathbf{x}^*)) = \alpha_*$ for $l \in \mathcal{I}_2$. Similarly, if we have $f_l(\mathbf{x}^*) < \alpha_*$ for $l \in I_2$ and $f_l(\mathbf{x}^*) = \alpha_*$ for $l \in \mathcal{I}_2$ then we find $f_l(L(\mathbf{x}^*)) = \alpha_*$ for $l \in \mathcal{I}_2$ and $f_l(L(\mathbf{x}^*)) < \alpha_*$ for $l \in I_1$. Therefore, we consider Cases IV and IV₁ as one case.

4.1.5 Cases I, II, III and IV are not possible at the point $\mathbf{x}^*$

Let us define the following subset $\Delta_7 = \{\mathbf{y} \in \Delta^7 : f_j(\mathbf{y}) = f_k(\mathbf{y}), j, k \in I\}$ of Δ^7 . We know that Δ_7 is not empty because we have $\bar{\mathbf{x}} \in \Delta_7$ (See §4.1). We aim to prove the following lemma

Lemma 4.2 *If the equality $F^\dagger(\mathbf{y}) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$ holds for some point $\mathbf{y} \in \Delta^7$, then we have $\mathbf{y} \in \Delta_7$. In particular, we get $\mathbf{x}^* \in \Delta_7$.*

To this purpose, we need to show that Cases I, II, III and IV are not possible at the point $\mathbf{x}^*$. We start with Case I.

Lemma 4.3 *Case I is not possible at the point $\mathbf{x}^*$.*

Proof We use the parametrization $\mathbf{p}_8^7$. Remember that $J_8 = I - \{8\}$. Then Case I is expressed as

$$\begin{aligned} \rho_2^* \sigma_1^* &= \alpha_*, & \rho_*^1 \sigma_2^* &= \alpha_*, & \rho_*^2 \sigma_3^* &= \alpha_*, & \rho_1^* \sigma_4^* &< \alpha_*, \\ \rho_*^2 \sigma_5^* &< \alpha_*, & \rho_*^1 \sigma_6^* &< \alpha_*, & \rho_2^* \sigma_7^* &< \alpha_* & \rho_1^* \sigma_*^8 &< \alpha_*. \end{aligned}$$

We obtain $(1 - \alpha_*)(x_2^*)^2 - (1 - \alpha_*)(1 - x_1^* - x_3^*)x_2^* - (x_1^* + x_3^*) = 0$ by simplifying the equation $\rho_*^1 \sigma_2^* = \alpha_*$. Hence x_2^* is a solution of the quadratic equation $q_1(x) = 0$ where we set $q_1(x) = (1 - \alpha_*)x^2 - (1 - \alpha_*)(1 - x_1^* - x_3^*)x - (x_1^* + x_3^*)$. Let $p_1: \Delta^{7,8} \rightarrow \mathbb{R}$ denote the map defined by $\mathbf{x}_8 \mapsto x_1 + x_3$. Let us define the function $D: \mathbb{R} \rightarrow \mathbb{R}$ so that $D(x) = (1 - \alpha_*)^2(1 - x)^2 + 4(1 - \alpha_*)x$. Note that $D(p_1(\mathbf{x}_8^*))$ corresponds to the discriminant of $q_1(x)$ where $\mathbf{x}_8^* = (\mathbf{p}_8^7)^{-1}(\mathbf{x}^*)$. Then, we see that

$$x_2^* = \frac{(1 - \alpha_*)(1 - x_1^* - x_3^*) \pm \sqrt{D(p_1(\mathbf{x}_8^*))}}{2(1 - \alpha_*)}.$$

We have $D(p_1(\mathbf{x}_8^*)) \geq 0$. Assume that the discriminant $D(p_1(\mathbf{x}_8^*)) > 0$. Since D is continuous, there exists an $\epsilon > 0$ so that $D(x_1^\circ + x_3^\circ) > 0$ for $x_1^\circ = x_1^* + \epsilon$ and $x_3^\circ = x_3^* + \epsilon$. We define the expression $\tilde{q}_1(x) = (1 - \alpha_*)x^2 - (1 - \alpha_*)(1 - x_1^\circ - x_3^\circ)x - (x_1^\circ + x_3^\circ)$. Then we see that the quadratic equation $\tilde{q}_1(x) = 0$ has two real roots

$$x_2^\circ = \frac{(1 - \alpha_*)(1 - x_1^\circ - x_3^\circ) \pm \sqrt{D(x_1^\circ + x_3^\circ)}}{2(1 - \alpha_*)}.$$

In fact, we can choose $\epsilon > 0$ so that the coefficients of $q_1(x)$ and $\tilde{q}_1(x)$ are sufficiently close to each other to conclude that the inequality holds $0 < \Sigma_\circ^8 < 1$ for the point $\mathbf{x}_8^\circ \in \Delta^{7,8}$ where we declare $x_4^\circ = x_4^*$, $x_5^\circ = x_5^*$, $x_6^\circ = x_6^*$, and $x_7^\circ = x_7^*$. Since we have $\tilde{q}_1(x_2^\circ) = 0$, we get $\rho_\circ^1 \sigma_2^\circ = \alpha_*$. Moreover, using the continuity of ρ^1 and σ^8 , the number $\epsilon > 0$ can be chosen so that the inequalities $\rho_1^\circ \sigma_\circ^8 < \alpha_*$ and $\rho_\circ^1 \sigma_6^\circ < \alpha_*$ are satisfied. Note that the expressions $\Sigma_2^* = \Sigma_2^\circ$, $\Sigma_1^* < \Sigma_1^\circ$, $\sigma_7^* = \sigma_7^\circ$, and $\sigma_5^* = \sigma_5^\circ$ imply that $\rho_2^\circ \sigma_1^\circ < \rho_2^* \sigma_1^*$, $\rho_2^\circ \sigma_7^\circ = \rho_2^* \sigma_7^*$, $\rho_2^\circ \sigma_3^\circ < \rho_2^* \sigma_3^*$, $\rho_\circ^2 \sigma_5^\circ = \rho_*^2 \sigma_5^*$ and $\rho_1^\circ \sigma_4^\circ < \rho_1^* \sigma_4^*$. Therefore, if we consider the point $\mathbf{x}^\circ = \mathbf{p}_8^7(\mathbf{x}_8^\circ) \in \Delta^7$, we see that $f_2(\mathbf{x}^\circ) = \alpha_*$ and $f_j(\mathbf{x}^\circ) < \alpha_*$ for every $j \in I - \{2\}$. Then we use the parametrization $\mathbf{p}_3^7$. We apply the perturbation $\text{pert}_1^{2,3}$ at the point $\mathbf{x}_3^\circ = (\mathbf{p}_3^7)^{-1}(\mathbf{x}^\circ)$. We obtain a point $\mathbf{z} = \mathbf{p}_3^7(\text{pert}_1^{2,3}(\mathbf{x}_3^\circ)) \in \Delta^7$ so that $F^\dagger(\mathbf{z}) < \alpha_*$. This contradicts with the minimality of α_* . Hence, we conclude that we cannot have $D(p_1(\mathbf{x}_8^*)) > 0$.

Let us assume that $D(p_1(\mathbf{x}_8^*)) = 0$. Then we get $x_2^* = (1 - x_1^* - x_3^*)/2$ which implies that $x_2^* = \Sigma_{2,3}^*$. Therefore, we find that $x_6^* < x_2^*$. Using the equalities $\rho_1^* \sigma_2^* = \alpha_*$ and $\Sigma_1^* = 1 - \Sigma_{2,3}^*$, we obtain that $\rho_1^* = \sigma(\Sigma_{2,3}^*)$ and $\alpha_* = \sigma(\Sigma_{2,3}^*) \sigma_2^* < \sigma_6^* \sigma_2^*$. By the inequality $\rho_1^* \sigma_6^* < \alpha_*$, we also have $\sigma_2^* \sigma_6^* = \rho_1^* \sigma_6^* < \alpha_*$. This is a contradiction. Thus, Case I is not possible at the point $\mathbf{x}^*$. $\square$

Lemma 4.4 *Case II is not possible at the point $\mathbf{x}^*$.*

Proof We need to use two parametrization simultaneously in this case. We start with the parametrization $\mathfrak{p}_8^7$ where $J_8 = I - \{8\}$. We express Case II as

$$\begin{aligned} \rho_2^* \sigma_1^* &< \alpha_*, & \rho_1^* \sigma_2^* &< \alpha_*, & \rho_2^* \sigma_3^* &< \alpha_*, & \rho_1^* \sigma_4^* &< \alpha_*, \\ \rho_2^* \sigma_5^* &< \alpha_*, & \rho_1^* \sigma_6^* &< \alpha_*, & \rho_2^* \sigma_7^* &= \alpha_* & \rho_1^* \sigma_8^* &= \alpha_*. \end{aligned}$$

Note that $\Sigma_8^* = \Sigma_1^* + \Sigma_4^*$. We simplify the equation $\rho_1^* \sigma_8^* = \alpha_*$. We find that $(\alpha_* - 1)(\Sigma_1^*)^2 + (1 - \alpha_*)(1 - \Sigma_4^*)\Sigma_1^* + \Sigma_4^* = 0$. In other words, Σ_1^* is a solution for the quadratic equation $q_2(x) = 0$ where $q_2(x) = (\alpha_* - 1)x^2 + (1 - \alpha_*)(1 - \Sigma_4^*)x + \Sigma_4^*$. Let $p_2: \Delta^{7,8} \rightarrow \mathbb{R}^4$ denote the map $\mathbf{x}_8 \mapsto x_4 + x_5 + x_6 + x_7$. Then, we see that

$$\Sigma_1^* = \frac{-(1 - \alpha_*)(1 - \Sigma_4^*) \pm \sqrt{D(p_2(\mathbf{x}^*))}}{2(\alpha_* - 1)}.$$

Since $D(p_2(\mathbf{x}^*))$ corresponds to the discriminant of $q_2(x)$, we get $D(p_2(\mathbf{x}^*)) \geq 0$. Next, we express Case II using the parametrization $\mathfrak{p}_7^7$. Remember that $J_7 = I - \{7\}$. We get

$$\begin{aligned} \rho_2^* \sigma_1^* &< \alpha_*, & \rho_1^* \sigma_2^* &< \alpha_*, & \rho_2^* \sigma_3^* &< \alpha_*, & \rho_1^* \sigma_4^* &< \alpha_*, \\ \rho_2^* \sigma_5^* &< \alpha_*, & \rho_1^* \sigma_6^* &< \alpha_*, & \rho_2^* \sigma_7^* &= \alpha_* & \rho_1^* \sigma_8^* &= \alpha_*. \end{aligned}$$

Note that $\Sigma_7^* = \Sigma_2^* + \Sigma_5^*$. By the simplifications in the equation $\rho_2^* \sigma_7^* = \alpha_*$, we find that $(\alpha_* - 1)(\Sigma_2^*)^2 + (1 - \alpha_*)(1 - \Sigma_5^*)\Sigma_2^* + \Sigma_5^* = 0$ which means that Σ_2^* is a solution for $q_3(x) = 0$ where we define $q_3(x) = (\alpha_* - 1)x^2 + (1 - \alpha_*)(1 - \Sigma_5^*)x + \Sigma_5^*$. Let $p_3: \Delta^{7,7} \rightarrow \mathbb{R}^4$ denote the map $\mathbf{x}_7 \mapsto x_1 + x_2 + x_3 + x_8$. Then, we get that

$$\Sigma_2^* = \frac{(\alpha_* - 1)(1 - \Sigma_5^*) \pm \sqrt{D(p_3(\mathbf{x}^*))}}{2(\alpha_* - 1)}.$$

Note that $D(p_3(\mathbf{x}^*))$ corresponds to the discriminant of $q_3(x)$. Therefore, we have $D(p_3(\mathbf{x}^*)) \geq 0$. Thus, we obtain either:

- (i) $D(p_2(\mathbf{x}^*)) > 0$, (ii) $D(p_3(\mathbf{x}^*)) > 0$, (iii) $D(p_2(\mathbf{x}^*)) = 0$ and $D(p_3(\mathbf{x}^*)) = 0$.

We examine each of these cases: (i) Let us set $x_1^\circ = x_1^*$ and $x_3^\circ = x_3^*$. Since we assume that $D(p_2(\mathbf{x}^*)) > 0$ and D is continuous, there exists an $\epsilon > 0$ so that

$D(x_4^\circ + x_5^\circ + x_6^\circ + x_7^\circ) > 0$ for $x_4^\circ = x_4^*$, $x_5^\circ = x_5^*$, $x_6^\circ = x_6^*$ and $x_7^\circ = x_7^* + \epsilon$. We consider the quadratic function $\tilde{q}_2(x) = (\alpha_* - 1)x^2 + (1 - \alpha_*) \left(1 - \sum_{i \in I_4} x_i^\circ\right) x + \sum_{i \in I_4} x_i^\circ$. We see that $\tilde{q}_2(x) = 0$ has two real roots:

$$(16) \quad \sum_{i \in I_1} x_i^\circ = \frac{-(1 - \alpha_*) \left(1 - \sum_{i \in I_4} x_i^\circ\right) \pm \sqrt{D(x_4^\circ + x_5^\circ + x_6^\circ + x_7^\circ)}}{2(\alpha_* - 1)}.$$

In fact, we can choose $\epsilon > 0$ so that the coefficients of $q_2(x)$ and $\tilde{q}_2(x)$ are sufficiently close to each other to conclude that $0 < \Sigma_0^8 < 1$ for the point $\mathbf{x}_8^\circ \in \Delta^{7,8}$. Moreover, since ρ^1 and ρ_1 are continuous, the number $\epsilon > 0$ can be chosen so that the inequalities $\rho_1^\circ \sigma_4^\circ < \alpha_*$, $\rho_1^\circ \sigma_2^\circ < \alpha_*$ and $\rho_1^\circ \sigma_6^\circ < \alpha_*$ are satisfied. Using the equality $\tilde{q}_1(\Sigma_1^\circ) = 0$, we derive that $\rho_1^\circ \sigma_8^\circ = \alpha_*$. Equalities $\rho_2^* = \rho_2^\circ$ and $\sigma_7^\circ < \sigma_7^*$ imply that $\rho_2^\circ \sigma_1^\circ = \rho_2^* \sigma_1^*$, $\rho_2^\circ \sigma_5^\circ = \rho_2^* \sigma_5^*$, $\rho_2^\circ \sigma_3^\circ = \rho_2^* \sigma_3^*$, and $\rho_2^\circ \sigma_7^\circ < \rho_2^* \sigma_7^*$. Therefore, if we consider the point $\mathbf{x}^\circ = \mathbf{p}_8^7(\mathbf{x}_8^\circ) \in \Delta^7$, we see that $f_8(\mathbf{x}^\circ) = \alpha_*$ and $f_l(\mathbf{x}^\circ) < \alpha_*$ for every $l \in J_8$. Then we apply the perturbation $\text{pert}_7^{8,8}$ at $\bar{\mathbf{x}}_8^\circ = (\mathbf{p}_8^7)^{-1}(\mathbf{x}^\circ)$. We obtain a point $\mathbf{t} = \mathbf{p}_8^7(\text{pert}_7^{8,8}(\bar{\mathbf{x}}_8^\circ)) \in \Delta^7$ so that $F^\dagger(\mathbf{t}) < \alpha_*$. This contradicts with the minimality of α_* . Hence, we conclude that $D(p_2(\mathbf{x}^*)) = 0$.

(ii) We set $x_4^\circ = x_4^*$, and $x_6^\circ = x_6^*$. We assume that $D(p_3(\mathbf{x}^*)) > 0$. Since D is continuous, there exists an $\epsilon > 0$ so that $D(x_1^\circ + x_2^\circ + x_3^\circ + x_8^\circ) > 0$ for $x_1^\circ = x_1^*$, $x_2^\circ = x_2^*$, $x_3^\circ = x_3^*$, and $x_8^\circ = x_8^* + \epsilon$. Let $\tilde{q}_3(x) = (\alpha_* - 1)x^2 + (1 - \alpha_*) \left(1 - \sum_{i \in I_5} x_i^\circ\right) x + \sum_{i \in I_5} x_i^\circ$. Then we determine that the quadratic equation $\tilde{q}_3(x) = 0$ has the following real roots:

$$(17) \quad \sum_{i \in I_2} x_i^\circ = \frac{-(1 - \alpha_*) \left(1 - \sum_{i \in I_5} x_i^\circ\right) \pm \sqrt{D(x_1^\circ + x_2^\circ + x_3^\circ + x_8^\circ)}}{2(\alpha_* - 1)}.$$

In fact, we can choose $\epsilon > 0$ so that the coefficients of $q_3(x)$ and $\tilde{q}_3(x)$ are sufficiently close to each other to conclude that $0 < \Sigma_0^7 < 1$ for the point $\mathbf{x}_7^\circ \in \Delta^{7,7}$. The equality $\tilde{q}_3(\Sigma_2^\circ) = 0$ implies that $\rho_2^\circ \sigma_7^\circ = \alpha_*$. Since ρ_2 and ρ^2 are continuous, the number $\epsilon > 0$ can be chosen so that the inequalities $\rho_2^\circ \sigma_1^\circ < \alpha_*$, $\rho_2^\circ \sigma_3^\circ < \alpha_*$ and $\rho_2^\circ \sigma_5^\circ < \alpha_*$ are all satisfied. The expressions $\rho_1^* = \rho_1^\circ$ and $\sigma_8^\circ < \sigma_8^*$ show that $\rho_1^\circ \sigma_2^\circ = \rho_1^* \sigma_2^*$, $\rho_1^\circ \sigma_6^\circ = \rho_1^* \sigma_6^*$, $\rho_1^\circ \sigma_4^\circ = \rho_1^* \sigma_4^*$ and $\rho_1^\circ \sigma_8^\circ < \rho_1^* \sigma_8^*$. Therefore, if we consider the point $\mathbf{x}^\circ = \mathbf{p}_7^7(\mathbf{x}_7^\circ) \in \Delta^7$, we see that $f_7(\mathbf{x}^\circ) = \alpha_*$ and $f_l(\mathbf{x}^\circ) < \alpha_*$ for every $l \in J_7$. Then we apply the perturbation $\text{pert}_8^{7,7}$ at $\bar{\mathbf{x}}_7^\circ = (\mathbf{p}_7^7)^{-1}(\mathbf{x}^\circ)$. We obtain a point $\mathbf{t} = \mathbf{p}_7^7(\text{pert}_8^{7,7}(\bar{\mathbf{x}}_7^\circ)) \in \Delta^7$ so that $F^\dagger(\mathbf{t}) < \alpha_*$. This contradicts with the minimality of α_* . Hence, we conclude that $D(p_3(\mathbf{x}^*)) = 0$.

(iii) Assume that $D(p_2(\mathbf{x}^*)) = 0$ and $D(p_3(\mathbf{x}^*)) = 0$. Then we get $\Sigma_1^* = x_8^*$ and $\Sigma_2^* = x_7^*$ by (16) and (17). Using the equalities $\rho_1^* \sigma_8^* = \alpha_*$ and $\rho_2^* \sigma_7^* = \alpha_*$, we get $(\sigma_8^*)^2 = (\sigma_7^*)^2$ which implies that $x_7^* = x_8^*$. We know that $\Sigma_0^* = 1$. Therefore we have

$x_8^* = 1/4$ and $x_7^* = 1/4$. We obtain $\alpha_* = (\sigma_8^*)^2$ which implies that $\alpha_* = 9$. Since we have $x_4^* < x_7^*$, we get $x_4^* < 1/4$. We calculate that $\rho_1^* \sigma_4^* = \sigma_8^* \sigma_4^* > 9 = \alpha_*$. This contradicts with the assumption $\rho_1^* \sigma_4^* < \alpha_*$ of Case II. We obtain a contradiction in any of the cases (i), (ii), and (iii). Therefore, Case II is not possible at $\mathbf{x}^*$. $\square$

Lemma 4.5 *Case III is not possible at the point $\mathbf{x}^*$.*

Proof We use the parametrization $\mathbf{p}_8^7$. Remember that $J_8 = I - \{8\}$. In this case, we have

$$\begin{aligned} \rho_2^* \sigma_1^* &= \alpha_*, & \rho_1^* \sigma_2^* &= \alpha_*, & \rho_3^* \sigma_3^* &= \alpha_*, & \rho_1^* \sigma_4^* &= \alpha_*, \\ \rho_2^* \sigma_5^* &= \alpha_*, & \rho_1^* \sigma_6^* &= \alpha_*, & \rho_2^* \sigma_7^* &< \alpha_*, & \rho_1^* \sigma_8^* &< \alpha_*. \end{aligned}$$

Using the equalities $\rho_1^* \sigma_2^* = \alpha_*$ and $\rho_1^* \sigma_6^* = \alpha_*$, we obtain that $x_2^* = x_6^*$. Similarly, we derive that $x_3^* = x_5^*$ from the equalities $\rho_2^* \sigma_3^* = \alpha_*$ and $\rho_2^* \sigma_5^* = \alpha_*$. Then we consider the equations $\rho_1^* \sigma_4^* = \alpha_*$ and $\rho_2^* \sigma_1^* = \alpha_*$. We have $\rho_1^* \sigma_4^* = \rho_2^* \sigma_1^*$. We simplify the last equality using the facts $x_2^* = x_6^*$ and $x_3^* = x_5^*$. We see that $(x_1^* - x_4^*)(x_5^* + x_6^*)(1 - x_1^* - x_4^* - x_5^* - x_6^*) = 0$. Since we know that $x_5^* + x_6^* > 0$ and $1 - x_1^* - x_4^* - x_5^* - x_6^* > 0$ by the discussion given in §4.1, we obtain that $x_1^* = x_4^*$. Then we derive that $\rho_1^* = \rho_2^*$. We use the equalities $\rho_1^* \sigma_2^* = \alpha_*$ and $\rho_2^* \sigma_3^* = \alpha_*$ to determine the relation $x_2^* = x_3^*$. As a result, we obtain that $x_1^* = x_4^*$, $x_2^* = x_3^*$, $x_3^* = x_5^*$, $x_5^* = x_6^*$. Remember that $\Sigma_0^* = 1$. Therefore, we find the inequality $x_1^* + 2x_2^* < 1/2$ or $\sigma(x_1^* + 2x_2^*) > 1$. We know that $\sigma(x_1^* + 2x_2^*)\sigma_1^* = (1/\sigma(x_1^* + 2x_2^*))\sigma_2^*$ from the equalities $\rho_2^* \sigma_1^* = \alpha_*$ and $\rho_1^* \sigma_2^* = \alpha_*$. We solve the equation above for $\sigma(x_1^* + 2x_2^*)$. We get $\sigma(x_1^* + 2x_2^*) = (\sigma_2^* \sigma_1^*)^{1/2} > 1$. Then we find that $(\sigma_1^*)^{1/2} < (\sigma_2^*)^{1/2}$ which implies that $x_2^* < x_1^*$. Thus, we obtain that $x_2^* < 1/6$.

By the equation $\rho_1^* \sigma_4^* = \alpha_*$ in this case, we observe that x_1^* satisfies the equation $(1 - \alpha_*)(x_1^*)^2 + (-2 + 2x_2^* - 2\alpha_* x_2^*)x_1^* + 1 - 2x_2^* = 0$. In other words, x_1^* is a solution for the quadratic function $q_4(x) = (1 - \alpha_*)x^2 + (-2 + 2x_2^* - 2\alpha_* x_2^*)x + 1 - 2x_2^*$. Let $p_4: \Delta^{7,8} \rightarrow \mathbb{R}$ be the map $\mathbf{x}_8 \mapsto x_2$. Let $\bar{D}: \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by $\bar{D}(x) = (-2 + 2x - 2\alpha_* x)^2 - 4(1 - \alpha_*)(1 - 2x)$. Then we calculate that

$$x_1^* = \frac{-(-2 + 2x_2^* - 2\alpha_* x_2^*) \pm \sqrt{\bar{D}(p_4(\mathbf{x}^*))}}{2(1 - \alpha_*)}.$$

Since $\bar{D}(p_4(\mathbf{x}^*))$ corresponds to the discriminant of $q_4(x)$, we have $\bar{D}(p_4(\mathbf{x}^*)) \geq 0$. Assume that $\bar{D}(p_4(\mathbf{x}^*)) = 0$. We get the equation $(4 - 8\alpha_* + 4\alpha_*^2)(x_2^*)^2 + 4\alpha_* = 0$ in x_2^* where the discriminant is $-\alpha_* < 0$. Since x_2^* is a positive real number, we obtain a contradiction. Therefore, we conclude that $\bar{D}(p_4(\mathbf{x}^*)) > 0$. Because the expression $2(1 - \alpha_*)x_1^* - 2(1 + (\alpha_* - 1)x_2^*)$ is strictly less than 0, we have

$$x_1^* = \frac{-(-2 + 2x_2^* - 2\alpha_* x_2^*) - \sqrt{\bar{D}(p_4(\mathbf{x}^*))}}{2(1 - \alpha_*)}.$$

Since $\bar{D}$ is continuous and $\bar{D}(p_4(\mathbf{x}^*)) > 0$, there exists an $\epsilon > 0$ so that $\bar{D}(x_2^\circ) > 0$ for $x_2^\circ = x_2^* + \epsilon$ where $x_2^\circ \in (0, 1/6)$. Let us define the following quadratic function $\tilde{q}_4(x) = (1 - \alpha_*)x^2 + (-2 + 2x_2^\circ - 2\alpha_*x_2^\circ)x + 1 - 2x_2^\circ$. Then we see that

$$x_1^\circ = \frac{-(-2 + 2x_2^\circ - 2\alpha_*x_2^\circ) - \sqrt{D(x_2^\circ)}}{2(1 - \alpha_*)}$$

is a solution for the equation $\tilde{q}_4(x_1^\circ) = 0$. In fact, we can choose $\epsilon > 0$ so that the coefficients of $q_4(x)$ and $\tilde{q}_4(x)$ are sufficiently close to each other to conclude that $0 < \Sigma_o^8 < 1$ for the point $\mathbf{x}_8^\circ \in \Delta^{7,8}$ where we set $x_i^\circ = x_2^\circ$ for $i = 3, 5, 6$, $x_4^\circ = x_1^\circ$, and $x_7^\circ = x_7^*$. Note that we have $\Sigma_2^\circ = \Sigma_1^\circ$ and $x_1^\circ = x_4^\circ$. Therefore, the equality $\tilde{q}_4(x_1^\circ) = 0$ implies that $\rho_1^\circ \sigma_4^\circ = \alpha_*$ and $\rho_2^\circ \sigma_1^\circ = \alpha_*$. Moreover, since ρ_1 and ρ_2 are continuous, the number $\epsilon > 0$ can be chosen so that the inequalities $\rho_1^\circ \sigma_o^8 < \alpha_*$ and $\rho_2^\circ \sigma_7^\circ < \alpha_*$ are satisfied. Then we define the functions

$$h_\alpha(x) = \frac{(-1 + (1 - \alpha)x) + \sqrt{g_\alpha(x)}}{(\alpha - 1)} \text{ and } H_\alpha(x) = \frac{\sigma(x)}{\sigma(h_\alpha(x) + 2x)}$$

where $g_\alpha(x) = (1 - \alpha)^2 x^2 + \alpha$. Note that we have $g_{\alpha_*}(x) = \bar{D}(x)$, $h_{\alpha_*}(x_2^\circ) = x_1^\circ$, $h_{\alpha_*}(x_2^*) = x_1^*$, and $H_{\alpha_*}(x_2^*) = \rho_2^* \sigma_5^*$. We prove the following:

Lemma 4.6 $H_{\alpha_*}(x)$ is a strictly decreasing function at x_2^* .

Proof See Appendix A. □

Because we set $x_2^\circ > x_2^*$, we obtain that $\rho_o^2 \sigma_5^\circ = H_{\alpha_*}(x_2^\circ) < H_{\alpha_*}(x_2^*) = \alpha_*$ by Lemma 4.6. Since we have $\rho_o^1 \sigma_2^\circ = \rho_o^2 \sigma_3^\circ$, $\rho_o^2 \sigma_3^\circ = \rho_o^2 \sigma_5^\circ$ and $\rho_o^2 \sigma_5^\circ = \rho_o^1 \sigma_6^\circ$, when we introduce the point $\mathbf{x}^\circ = \mathbf{p}_8^7(\mathbf{x}_8^\circ) \in \Delta^7$, we see that $f_1(\mathbf{x}^\circ) = \alpha_*$, $f_4(\mathbf{x}^\circ) = \alpha_*$, and $f_l(\mathbf{x}^\circ) < \alpha_*$ for $l \in I - \{1, 4\}$. We use the parametrization $\mathbf{p}_2^7$ or $\mathbf{p}_3^7$. We apply the perturbation $\text{pert}_3^{\dagger,2}$ or $\text{pert}_2^{\dagger,3}$ at $\mathbf{x}_2^\circ = (\mathbf{p}_2^7)^{-1}(\mathbf{x}^\circ) \in \Delta^{7,2}$ or $\mathbf{x}_3^\circ = (\mathbf{p}_3^7)^{-1}(\mathbf{x}^\circ) \in \Delta^{7,3}$, respectively. We obtain a point $\tilde{\mathbf{x}}^\circ = \mathbf{p}_2^7(\text{pert}_3^{\dagger,2}(\mathbf{x}_2^\circ)) \in \Delta^7$ so that $f_4(\tilde{\mathbf{x}}^\circ) = \alpha_*$ and $f_l(\tilde{\mathbf{x}}^\circ) < \alpha_*$ for $l \in I - \{4\}$. Then we use the parametrization $\mathbf{p}_5^7$ or $\mathbf{p}_6^7$. We apply the perturbation $\text{pert}_6^{4,5}$ or $\text{pert}_5^{4,6}$ at $\mathbf{x}_5^\circ = (\mathbf{p}_5^7)^{-1}(\tilde{\mathbf{x}}^\circ) \in \Delta^{7,5}$ or $\mathbf{x}_6^\circ = (\mathbf{p}_6^7)^{-1}(\tilde{\mathbf{x}}^\circ) \in \Delta^{7,6}$, respectively. We obtain a point $\mathbf{w} = \mathbf{p}_5^7(\text{pert}_6^{4,5}(\mathbf{x}_5^\circ)) \in \Delta^7$ so that $F^1(\mathbf{w}) < \alpha_*$. This contradicts with the minimality of α_* . Therefore, Case III is not possible at $\mathbf{x}^*$. □

Before we proceed to Case IV, it is necessary to state two preliminary lemmas:

Lemma 4.7 Let $I_\alpha = [9, 5 + 3\sqrt{2}]$. Let us define the following continuous functions

$$g^\pm(\alpha) = \frac{(1 - \alpha^2)\sqrt{\alpha} + \alpha^3 + \alpha \pm \sqrt{d(\alpha)}}{2(-1 + (\alpha^3 - \alpha)\sqrt{\alpha} + \alpha^2)} \text{ and } d(\alpha) = b(\alpha) - c(\alpha)$$

over the interval I_α where we set $b(\alpha) = \alpha^6 + 6\alpha^3 + \alpha^2 + 4 + (4\alpha^4 + 6\alpha + 4)\alpha^{1/2}$, and $c(\alpha) = (3\alpha^5 + 6\alpha^4 + 3\alpha + (2\alpha^5 + 4\alpha^3 + 8\alpha^2)\alpha^{1/2})\alpha^{1/2}$. Then neither $G^+(\alpha)$ nor $G^-(\alpha)$ takes the value 1 over the interval I_α where we define

$$G^\pm(\alpha) = \frac{(\sqrt{\alpha} + 1)(\alpha - 1)(g^\pm(\alpha))^2 + 2(\alpha - 1)g^\pm(\alpha) + \sqrt{\alpha} + 3}{(\sqrt{\alpha} + 1)((\alpha - 1)g^\pm(\alpha) + 1)}.$$

Proof See Appendix A □

Lemma 4.8 Let $I_\alpha = [9, 5 + 3\sqrt{2}]$. Let us define the following continuous functions

$$h^\pm(\alpha) = \frac{(\alpha - 1)^2 \pm \sqrt{(\alpha - 1)(\alpha^3 - 3\alpha^2 - 5\alpha - 9)}}{2(\alpha^2 - 1)}$$

for every $\alpha \in I_\alpha$. Then neither $H^+(\alpha)$ nor $H^-(\alpha)$ takes the value $1/2$ over I_α where

$$H^\pm(\alpha) = \frac{(\alpha + 1)(\alpha^2 - 1)(h^\pm(\alpha))^2 + 5(\alpha^2 - 1)h^\pm(\alpha) + \alpha^2 + 9\alpha - 4}{(\alpha + 1)(\alpha^2 - 1)h^\pm(\alpha) + (\alpha + 1)(3\alpha - 1)}.$$

Proof See Appendix A □

Lemma 4.9 Case IV is not possible at the point $\mathbf{x}^*$.

Proof Let us assume that Case IV holds at the point $\mathbf{x}^*$. We have the expressions

$$\begin{aligned} \rho_2^* \sigma_1^* &= \alpha_*, & \rho_{2,3}^* \sigma_2^* &= \alpha_*, & \rho_{1,3}^* \sigma_3^* &= \alpha_*, & \rho_1^* \sigma_4^* &< \alpha_*, \\ \rho_{1,3}^* \sigma_5^* &< \alpha_*, & \rho_{2,3}^* \sigma_6^* &< \alpha_*, & \rho_2^* \sigma_7^* &= \alpha_* & \rho_1^* \sigma_8^* &= \alpha_*. \end{aligned}$$

Using the equalities $\rho_2^* \sigma_1^* = \alpha_*$ and $\rho_2^* \sigma_7^* = \alpha_*$, we derive that $x_1^* = x_7^*$. We use the parametrization $\mathbf{p}_7^7$. Then we see that

$$\begin{aligned} \rho_2^* \sigma_1^* &= \alpha_*, & \rho_1^* \sigma_2^* &= \alpha_*, & \rho_2^* \sigma_3^* &= \alpha_*, & \rho_1^* \sigma_4^* &< \alpha_*, \\ \rho_2^* \sigma_5^* &< \alpha_*, & \rho_1^* \sigma_6^* &< \alpha_*, & \rho_2^* \sigma_7^* &= \alpha_* & \rho_1^* \sigma_8^* &= \alpha_*. \end{aligned}$$

We know that Σ_1^* is a solution of the equation $q_2(x)$ from the equality $\rho_1^* \sigma_2^* = \alpha_*$. Hence, we get $D(p_3(\mathbf{x}^*)) \geq 0$. Let us assume that $D(p_3(\mathbf{x}^*)) = 0$. Then we have $\Sigma_1^* = x_8^*$. Using the equality $\rho_1^* \sigma_8^* = \alpha_*$, we see that $(\sigma_8^*)^2 = \alpha_*$. In other words, we get $x_8^* = 1/(\alpha_*^{1/2} + 1)$. By the equalities $\rho_1^* \sigma_2^* = \alpha_*$ and $\rho_1^* = 1/\sigma(x_8)$, we find that $\sigma_2^*/\sigma(x_8) = \alpha_*$. We solve the last equation for x_2^* . We obtain that $x_2^* = 1/(\alpha_*^{3/2} + 1)$. We multiply the equalities $\rho_2^* \sigma_1^* = \alpha_*$ and $\rho_2^* \sigma_3^* = \alpha_*$ side by side. We find that $\sigma_1^* \sigma_3^* = \alpha_*^2$ or

$$(18) \quad 1 - (x_1^* + x_3^*) = (\alpha_*^2 - 1)x_1^* x_3^*.$$

Since we know Σ_1^* and x_2^* in terms of α_* , we can calculate $x_1^* + x_3^*$ in terms of α_* . We get $x_1^* + x_3^* = (\alpha_* - \alpha_*^{1/2})/(\alpha_*^{3/2} + 1)$ or $x_3^* = (\alpha_* - \alpha_*^{1/2})/(\alpha_*^{3/2} + 1) - x_1^*$. We plug x_3^* in (18). Then we obtain the following quadratic equation in x_1^* where $(\alpha_*^{3/2} + 1)(\alpha_*^2 - 1)(x_1^*)^2 - \alpha_*^{1/2}(\alpha_*^2 - 1)(\alpha_*^{1/2} - 1)x_1^* + (\alpha_* + 1)\alpha_*^{1/2} + 1 - \alpha_* = 0$. Then we find that

$$x_1^* = \frac{(1 - \alpha_*^2)\sqrt{\alpha_*} + \alpha_*^3 - \alpha_* \pm \sqrt{d(\alpha_*)}}{2(-1 + (\alpha_*^3 - \alpha_*)\sqrt{\alpha_*} + \alpha_*^2)}$$

where $d(\alpha_*) = b(\alpha_*) - c(\alpha_*)$ for the functions $b(\alpha)$ and $c(\alpha)$ defined in Lemma 4.7. Note that we have either $x_1^* = g^+(\alpha_*)$ or $x_1^* = g^-(\alpha_*)$. Using the equality $\rho_2^*\sigma_1^* = \alpha_*$, we calculate that

$$\Sigma_2^* = \frac{1 - x_1^*}{(\alpha_* - 1)x_1^* + 1}.$$

We know that $x_1^* = x_7^*$ and $\Sigma_1^* = x_8^*$. Therefore, we have $1 = 2x_1^* + \Sigma_2^* + 2x_8^*$ which implies that

$$1 = \frac{(\sqrt{\alpha_*} + 1)(\alpha_* - 1)(x_1^*)^2 + 2(\alpha_* - 1)x_1^* + \sqrt{\alpha_*} + 3}{(\sqrt{\alpha_*} + 1)((\alpha_* - 1)x_1^* + 1)}.$$

Then we must have either $G^+(\alpha_*) = 1$ or $G^-(\alpha_*) = 1$. This is a contradiction. Because, in Lemma 4.7, we show that $G^\pm(\alpha) \neq 1$ for every $\alpha \in I_\alpha$. Therefore, we conclude that $D(p_3(\mathbf{x}^*)) > 0$. We need to consider two cases:

$$(i) \Sigma_1^* < x_8^* \text{ or, } (ii) \Sigma_1^* > x_8^*.$$

(i) Let us assume that $\Sigma_1^* < x_8^*$. Remember that $J_8 = I - \{8\}$. We use the parametrization $\mathfrak{p}_8^7$:

$$\begin{aligned} \rho_2^*\sigma_1^* &= \alpha_*, & \rho_*^1\sigma_2^* &= \alpha_*, & \rho_*^2\sigma_3^* &= \alpha_*, & \rho_*^1\sigma_4^* &< \alpha_*, \\ \rho_*^2\sigma_5^* &< \alpha_*, & \rho_*^1\sigma_6^* &< \alpha_*, & \rho_*^2\sigma_7^* &= \alpha_*, & \rho_*^1\sigma_8^* &= \alpha_*. \end{aligned}$$

From the equality $\rho_*^1\sigma_2^* = \alpha_*$, we see that x_2^* is a solution of the quadratic equation $q_1(x) = 0$. Since we also have the inequality $\rho_*^1\sigma_6^* < \alpha_*$, we know from Case I that $D(p_1(\mathbf{x}^*)) > 0$. There are two choices for x_2^* :

$$x_2^* = \frac{(1 - \alpha_*)(1 - x_1^* - x_3^*) \pm \sqrt{D(x_1^* + x_3^*)}}{2(1 - \alpha_*)}.$$

Note that the equality $(1 - \alpha_*)(x_1^* + 2x_2^* + x_3^* - 1) = -(D(x_1^* + x_3^*))^{1/2}$ implies that $x_1^* + 2x_2^* + x_3^* > 1$. In other words, we have $x_2^* > \Sigma_{2,3}^*$. Therefore, we get $x_2^* > x_6^*$. Then, we obtain that $\sigma_2^* < \sigma_6^*$. Using the expressions $\rho_*^1\sigma_2^* = \alpha_*$ and $\rho_*^1\sigma_6^* < \alpha_*$ in

Case IV, we find that $\alpha_* = \rho_*^1 \sigma_2^* < \rho_*^1 \sigma_6^* < \alpha_*$, a contradiction. Therefore, we get $x_1^* + 2x_2^* + x_3^* < 1$. In other words, we conclude that

$$(19) \quad x_2^* = \frac{(1 - \alpha_*)(1 - x_1^* - x_3^*) + \sqrt{D(x_1^* + x_3^*)}}{2(1 - \alpha_*)}.$$

Since we know that D is continuous and $D(x_1^* + x_3^*) > 0$, there exists an $\epsilon > 0$ so that $D(x_1^\circ + x_3^\circ) > 0$ where $x_1^\circ = x_1^* + \epsilon$ and $x_3^\circ = x_3^*$. We define the quadratic function $\tilde{q}_1(x) = (1 - \alpha)x^2 - (1 - \alpha)(1 - x_1^\circ - x_3^\circ)x - (x_1^\circ + x_3^\circ)$. Then we see that one of the roots of the quadratic equation $\tilde{q}_1(x) = 0$ is

$$x_2^\circ = \frac{(1 - \alpha_*)(1 - x_1^\circ - x_3^\circ) + \sqrt{D(x_1^\circ + x_3^\circ)}}{2(1 - \alpha_*)}.$$

Moreover, we can choose $\epsilon > 0$ so that the coefficients of $q_1(x)$ and $\tilde{q}_1(x)$ are sufficiently close each other to conclude that $0 < \Sigma_\circ^8 < 1$ for the point $\mathbf{x}_8^\circ \in \Delta^{7,8}$ where we set $x_4^\circ = x_4^*$, $x_5^\circ = x_5^*$, $x_6^\circ = x_6^*$, and $x_7^\circ = x_7^*$. Since we have $\tilde{q}_1(x_2^\circ) = 0$, we get $\rho_\circ^1 \sigma_2^\circ = \alpha_*$. Using the continuity of ρ_1 , the number $\epsilon > 0$ can be chosen so that the inequalities $\rho_1^\circ \sigma_4^\circ < \alpha_*$ and $\rho_\circ^1 \sigma_6^\circ < \alpha_*$ hold. Let $K_1 = x_3^*$ and $K_2 = \Sigma_4^*$. Let us define the following functions

$$h_\alpha(x) = \frac{(1 - \alpha)(1 - x - K_1) + \sqrt{g_\alpha(x)}}{2(1 - \alpha)} \text{ and } H_\alpha(x) = \frac{\sigma(x + h_\alpha(x) + K_1)}{\sigma(x + h_\alpha(x) + K_1 + K_2)}$$

where $g_\alpha(x) = (1 - \alpha)^2(1 - x - K_1)^2 + 4(1 - \alpha)(x + K_1)$. Notice that we have $g_{\alpha_*}(x) = D(x + K_1)$, $h_{\alpha_*}(x_1^*) = x_2^*$, $h_{\alpha_*}(x_1^\circ) = x_2^\circ$ and $H_{\alpha_*}(x_1^*) = \rho_1^* \sigma_*^8$. Then we prove

Lemma 4.10 $H_{\alpha_*}(x)$ is a strictly decreasing function at x_1^* .

Proof See Appendix A. □

Since we set $x_1^\circ + x_3^\circ > x_1^* + x_3^*$ and $H_{\alpha_*}(x)$ is a strictly decreasing function at $x_1^* + x_3^*$ by Lemma 4.10, we obtain that $\rho_1^\circ \sigma_\circ^8 = H_{\alpha_*}(x_1^\circ + x_3^\circ) < H_{\alpha_*}(x_1^* + x_3^*) = \alpha_*$. The expressions $\sigma_*^2 = \sigma_\circ^2$, $\rho_2^* = \rho_\circ^2$, $\sigma_1^\circ < \sigma_1^*$ and $\sigma_3^\circ = \sigma_3^*$ imply that $\rho_2^\circ \sigma_1^\circ < \rho_2^* \sigma_1^*$, $\rho_\circ^2 \sigma_5^\circ = \rho_2^* \sigma_5^*$, $\rho_\circ^2 \sigma_3^\circ = \rho_2^* \sigma_3^*$, and $\rho_2^\circ \sigma_7^\circ = \rho_2^* \sigma_7^*$. Therefore, if we consider the point $\mathbf{x}^\circ = \mathbf{p}_8^7(\mathbf{x}_8^\circ) \in \Delta^7$, we see that $f_l(\mathbf{x}^\circ) < \alpha_*$ for every $l \in I - \{2, 3, 7\}$ and $f_l(\mathbf{x}^\circ) = \alpha_*$ for $l = 2, 3, 7$. We use the parametrization $\mathbf{p}_3^7$. Then we apply the perturbation $\text{pert}_1^{2,3}$ at $\mathbf{x}_3 = (\mathbf{p}_3^7)^{-1}(\mathbf{x}^\circ) \in \Delta^{7,3}$. We obtain a point $\tilde{\mathbf{x}}^\circ = \mathbf{p}_3^7(\text{pert}_1^{2,3}(\mathbf{x}_3)) \in \Delta^7$ so that $f_l(\tilde{\mathbf{x}}^\circ) < \alpha_*$ for every $l \in I - \{3, 7\}$ and $f_l(\tilde{\mathbf{x}}^\circ) = \alpha_*$ for $l = 3, 7$. Next, we use the parametrization $\mathbf{p}_2^7$. We apply the perturbation $\text{pert}_1^{3,2}$ at $\mathbf{x}_2 = (\mathbf{p}_2^7)^{-1}(\tilde{\mathbf{x}}^\circ) \in \Delta^{7,2}$. We obtain a point $\bar{\mathbf{x}}^\circ = \mathbf{p}_2^7(\text{pert}_1^{3,2}(\mathbf{x}_2)) \in \Delta^7$ so that $f_l(\bar{\mathbf{x}}^\circ) < \alpha_*$ for every $l \in J_7$

and $f_7(\tilde{\mathbf{x}}^\circ) = \alpha_*$. Finally, we use the parametrization $\mathbf{p}_7^7$. We apply the perturbation $\text{pert}_8^{7,7}$ at $\mathbf{x}_7 = (\mathbf{p}_7^7)^{-1}(\tilde{\mathbf{x}}^\circ) \in \Delta^{7,7}$. We obtain a point $\mathbf{t} = \mathbf{p}_7^7(\text{pert}_8^{7,7}(\mathbf{x}_7)) \in \Delta^7$ so that $f_l(\mathbf{t}) < \alpha_*$ for every $l \in I$, i.e., $F^1(\mathbf{t}) < \alpha_*$. This contradicts with the minimality of α_* . Therefore, we cannot have $\Sigma_1^* < x_8^*$.

(ii) Let us assume that $\Sigma_1^* > x_8^*$. We need to divide this case into three parts:

$$(iii) x_1^* > x_8^*, \quad (iv) x_1^* = x_8^*, \quad (v) x_1^* < x_8^*.$$

(iii) Let us assume that $x_8^* < x_1^*$. Remember that we have $x_1^* = x_7^*$. Therefore, we obtain that $x_8^* < x_7^*$. We know that $q_2(\Sigma_1^*) = 0$ by the equality $\sigma_1^* \sigma_8^* = \alpha_*$. The assumption $\Sigma_1^* > x_8^*$ implies that $D(p_3(\mathbf{x}^*)) > 0$. Let us denote the second root of $q_2(x) = 0$ with t_1 . The formula for the addition of the roots of a quadratic equation for $q_2(x) = 0$ gives that $\Sigma_1^* + t_1 = 1 - \Sigma_4^*$ or $t_1 = x_8^*$. Therefore, we get $q_2(x_8^*) = 0$.

Similarly, by the equality $\rho_2^* \sigma_7^* = \alpha_*$, we know that $q_2(\Sigma_2^*) = 0$ which implies that $D(p_3(\mathbf{x}^*)) \geq 0$. If we assume that $D(p_3(\mathbf{x}^*)) = 0$, we obtain that $\Sigma_2^* = x_7^*$. If we assume that $D(p_3(\mathbf{x}^*)) > 0$, the formula for the addition of the roots of a quadratic equation for $q_3(x) = 0$ gives that $\Sigma_2^* + t_2 = 1 - \Sigma_5^*$ or $t_2 = x_7^*$ where t_2 denote the second root of $q_3(x)$. In either case, we see that $q_3(x_1^*) = 0$. Let us define the linear function $L(x) = q_2(x) - q_3(x)$ where

$$q_2(x) - q_3(x) = (\Sigma_4^* - \Sigma_5^*)((\alpha_* - 1)x + 1).$$

Using the expressions $\rho_1^* \sigma_8^* = \alpha_*$ and $\rho_1^* \sigma_4^* < \alpha_*$, we conclude that $x_8^* < x_4^*$. Similarly, we have $x_2^* < x_6^*$ and $x_3^* < x_5^*$ by $\rho_*^1 \sigma_2^* = \alpha_*$, $\rho_*^1 \sigma_6^* < \alpha_*$, $\rho_*^2 \sigma_5^* < \alpha_*$ and $\rho_*^2 \sigma_3^* = \alpha_*$. Then, we obtain that $x_2^* + x_3^* + x_8^* < \Sigma_2^*$ or $\Sigma_1^* + x_8^* < \Sigma_2^* + x_7^*$ which implies that $\Sigma_5^* < \Sigma_4^*$. Therefore, L has only one root $x_0 = 1/(1 - \alpha_*) < 0$.

We calculate that $L(0) = q_2(0) - q_3(0)$ where $q_2(0) - q_3(0) = \Sigma_4^* - \Sigma_5^* > 0$. It is clear that $x_1^* < \Sigma_1^*$. Since we have $x_8^* < x_1^* < \Sigma_1^*$ and q_2 is concave up, we derive that $q_2(x_1^*) < 0$. Then, we calculate $L(x)$ at the point x_1^* . We see that $L(x_1^*) = q_2(x_1^*) - q_3(x_1^*)$ where $q_2(x_1^*) - q_3(x_1^*) = q_2(x_1^*) < 0$. By the Intermediate Value Theorem, $L(x)$ must have a root in the interval $(0, 1)$. This is a contradiction. Hence, we cannot have $x_8^* < x_1^*$.

(iv) Assume that $x_1^* = x_8^*$. We use the equalities $\rho_2^* \sigma_1^* = \alpha_*$ and $\rho_1^* \sigma_8^* = \alpha_*$. We find that $\rho_2^* = \rho_1^*$. This means that $\Sigma_1^* = \Sigma_2^*$. Using the expressions $\rho_1^* \sigma_4^* < \alpha_*$ and $\rho_2^* \sigma_1^* = \alpha_*$, we conclude that $x_1^* < x_4^*$. We know that $x_2^* < x_6^*$ and $x_3^* < x_5^*$ from the expressions $\rho_*^1 \sigma_6^* < \alpha_*$, $\rho_*^1 \sigma_2^* = \alpha_*$, $\rho_*^2 \sigma_3^* = \alpha_*$ and $\rho_*^2 \sigma_5^* < \alpha_*$. Then we find that $\Sigma_1^* = \Sigma_2^* > \Sigma_1^*$, a contradiction. Therefore, we cannot have $x_1^* = x_8^*$.

(v) We need to divide the case $x_1^* < x_8^*$ into the following three parts below:

$$(vi) \Sigma_2^* < \frac{1}{2}, \quad (vii) \Sigma_2^* = \frac{1}{2}, \quad (viii) \Sigma_2^* > \frac{1}{2}.$$

(vi) Assume that $\Sigma_2^* < 1/2$. We consider the equalities $\rho_3^2 \sigma_3^* = \alpha_*$ and $\rho_2^* \sigma_1^* = \alpha_*$. We obtain that $\rho_3^2 \sigma_3^* = \rho_2^* \sigma_1^*$ or $\rho_2^* = (\sigma_3^* / \sigma_1^*)^{1/2}$ which implies that

$$\Sigma_2^* = \frac{1}{\sqrt{\frac{\sigma_3^*}{\sigma_1^*} + 1}}.$$

Let us define the function $U: \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $U(x, y) = 1/((\sigma(y)/\sigma(x))^{1/2} + 1)$. Note that we have $\Sigma_2^* = U(x_1^*, x_3^*)$. We already know that Σ_1^* is a root of the quadratic equation $q_2(x) = 0$, i.e., $(\alpha_* - 1)(\Sigma_1^*)^2 + (1 - \alpha_*)(1 - \Sigma_4^*)\Sigma_1^* + \Sigma_4^* = 0$. Note that the previous equation can be expanded and the terms can be rearranged so that x_2^* is a solution of the quadratic equation $\bar{q}_2(x) = 0$ where

$$\bar{q}_2 = \delta_* x^2 - \delta_*(1 - 2K - \Sigma_4)x + \delta_* K^2 - \delta_* K(1 - \Sigma_4) + \Sigma_4,$$

$K = x_1^* + x_3^*$ and $\delta_* = \alpha_* - 1$. Let us define the function $d_2: \mathbb{R}^2 \rightarrow \mathbb{R}$ so that $d_2(x, y) = (1 - \alpha_*)^2(1 - x - U(x, y))^2 - 4(\alpha_* - 1)(x + U(x, y))$. Since we know that $x_1^* = x_7^*$ from the equalities $\rho_2^* \sigma_1^* = \alpha_*$ and $\rho_2^* \sigma_7^* = \alpha_*$, we derive the expression $d_2(x_1^*, x_3^*) = (1 - \alpha)^2(1 - x_1^* - U(x_1^*, x_3^*))^2 - 4(\alpha - 1)(x_1^* + U(x_1^*, x_3^*))$. Note that the discriminant of $\bar{q}_2(x)$ is $d_2(x_1^*, x_3^*)$. The condition of (ii) implies that

$$(20) \quad x_2^* = \frac{(\alpha_* - 1)(1 - 3x_1^* - 2x_3^* - U(x_1^*, x_3^*)) + \sqrt{d_2(x_1^*, x_3^*)}}{2(\alpha_* - 1)}.$$

Equations $q_1(x) = 0$ and $\bar{q}_2(x) = 0$ have x_2^* as a common solution. Let $D_1: \mathbb{R}^2 \rightarrow \mathbb{R}$ be the function defined by $D_1(x, y) = (1 - \alpha)^2(1 - x - y)^2 - 4(\alpha - 1)(x + y)$. Then we have

$$(21) \quad x_2^* = \frac{(1 - \alpha_*)(1 - x_1^* - x_3^*) \pm \sqrt{D_1(x_1^*, x_3^*)}}{2(1 - \alpha_*)}.$$

We prove the following

Lemma 4.11 *For any $\alpha \in I_\alpha = [9, 5 + 3\sqrt{2}]$, let F_α^1 and F_α^2 be the functions defined as*

$$F_\alpha^1(x, y) = \frac{(\alpha - 1)(1 - 3x - 2y - U(x, y)) + \sqrt{d_2(x, y)}}{2(\alpha - 1)},$$

$$F_\alpha^2(x, y) = \frac{(1 - \alpha)(1 - x - y) + \sqrt{D_1(x, y)}}{2(1 - \alpha)}$$

where we have $d_2(x, y) = (1 - \alpha)^2(1 - x - U(x, y))^2 - 4(\alpha - 1)(x + U(x, y))$ and $D_1(x, y) = (1 - \alpha)^2(1 - x - y)^2 - 4(\alpha - 1)(x + y)$. Then there exists a neighborhood of (x_1^*, x_3^*) so that the equation $F_{\alpha_*}^1(x, y) - F_{\alpha_*}^2(x, y) = 0$ defines y as a function of x over that neighborhood.

Proof See Appendix A. □

Note that we have $x_2^* = F_{\alpha_*}^1(x_1^*, x_3^*)$ by (20) and $x_2^* = F_{\alpha_*}^2(x_1^*, x_3^*)$ by (21). In addition, we also have $d_2(x_1^*, x_3^*) = D(x_4^* + x_5^* + x_6^* + x_7^*)$. We know that $\rho_2^* \sigma_1^* = \rho_2^* \sigma_7^*$ where we rewrite $\rho_2^* \sigma_7^* = \sigma(U(x_1^*, x_2^*)) \sigma_1^*$ or $\rho_2^* \sigma_7^* = (\sigma_1^* \sigma_3^*)^{1/2}$. Similarly, we observe that $\rho_2^* \sigma_3^* = \sigma_3^* / \sigma(U(x_1^*, x_2^*))$ or $\rho_2^* \sigma_3^* = (\sigma_1^* \sigma_3^*)^{1/2}$. By Lemma 4.11, we also know that the equation $F_{\alpha_*}^1(x, y) = F_{\alpha_*}^2(x, y)$ defines y as a function of x around a neighborhood of x_1^* . Then we define the functions

$$h_{\alpha}(x) = y(x) \text{ and } H_{\alpha}(x) = \sqrt{\sigma(x)\sigma(h_{\alpha}(x))}$$

where we have the equalities $h_{\alpha_*}(x_1^*) = x_3^*$, $H_{\alpha_*}(x_1^*) = \rho_2^* \sigma_1^*$, $H_{\alpha_*}(x_1^*) = \rho_2^* \sigma_7^*$, and $H_{\alpha_*}(x_1^*) = \rho_2^* \sigma_3^*$. We prove the following lemma:

Lemma 4.12 $H_{\alpha_*}(x)$ is a strictly monotone function at the point x_1^* .

Proof See Appendix A. □

We examine each of the cases where H_{α_*} is strictly decreasing or strictly increasing at x_1^* . If $H_{\alpha_*}(x)$ is a strictly decreasing function at x_1^* , then, for a sufficiently small $\epsilon > 0$, the inequality $0 < \Sigma_{\circ}^8 < 1$ holds for the point $\mathbf{x}^{\circ} = \mathbf{p}_8^7(\mathbf{x}_8^{\circ}) \in \Delta^7$ where we set $x_4^{\circ} = x_4^*$, $x_5^{\circ} = x_5^*$, $x_7^{\circ} = x_1^*$, $x_1^{\circ} = x_1^* + \epsilon$, $x_3^{\circ} = h_{\alpha_*}(x_1^{\circ})$, $x_2^{\circ} = F^1(x_1^{\circ}, x_3^{\circ})$, and $x_6^{\circ} = U(x_1^{\circ}, x_3^{\circ}) - x_4^{\circ} - x_5^{\circ}$. If $H_{\alpha_*}(x)$ is a strictly increasing function at x_1^* , then, for a sufficiently small $\epsilon > 0$, the inequality $0 < \Sigma_{\circ}^8 < 1$ holds for the point $\mathbf{x}^{\circ} = \mathbf{p}_8^7(\mathbf{x}_8^{\circ}) \in \Delta^7$ where we set $x_4^{\circ} = x_4^*$, $x_5^{\circ} = x_5^*$, $x_7^{\circ} = x_1^*$, $x_1^{\circ} = x_1^* - \epsilon$, $x_3^{\circ} = h_{\alpha_*}(x_1^{\circ})$, $x_2^{\circ} = F^1(x_1^{\circ}, x_3^{\circ})$, and $x_6^{\circ} = U(x_1^{\circ}, x_3^{\circ}) - x_4^{\circ} - x_5^{\circ}$. In either case, we obtain that $\rho_2^{\circ} \sigma_1^{\circ} = \rho_2^{\circ} \sigma_7^{\circ}$, $\rho_2^{\circ} \sigma_7^{\circ} = \rho_2^{\circ} \sigma_3^{\circ}$ and $\rho_2^{\circ} \sigma_3^{\circ} = H_{\alpha_*}(x_1^{\circ}) < H_{\alpha_*}(x_1^*) = \alpha_*$ by Lemma 4.12. Moreover, because ρ^2 , ρ^1 , and ρ_2 are continuous, the number $\epsilon > 0$ can be chosen so that the inequalities $\rho_1^{\circ} \sigma_4^{\circ} < \alpha_*$, $\rho_2^{\circ} \sigma_5^{\circ} < \alpha_*$ and $\rho_1^{\circ} \sigma_6^{\circ} < \alpha_*$ hold. Since x_2° is on the curve defined by the equality $F_{\alpha_*}^1(x, h_{\alpha_*}(x)) = F_{\alpha_*}^2(x, h_{\alpha_*}(x))$ on a neighborhood of x_1^* such that $x_2^{\circ} = F_{\alpha_*}^1(x_1^{\circ}, h_{\alpha_*}(x_1^{\circ}))$ and $x_2^{\circ} = F_{\alpha_*}^2(x_1^{\circ}, h_{\alpha_*}(x_1^{\circ}))$, we get $\rho_2^{\circ} \sigma_2^{\circ} = \alpha_*$ and $\rho_1^{\circ} \sigma_8^{\circ} = \alpha_*$. In other words, we get $f_l(\mathbf{x}^{\circ}) < \alpha_*$ for every $l \in I - \{2, 8\}$ and $f_l(\mathbf{x}^{\circ}) = \alpha_*$ for $l \in \{2, 8\}$. Then we use the parametrization $\mathbf{p}_8^7$. We apply the perturbation $\text{pert}_{\gamma}^{8,8}$ at $\mathbf{x}_8 = (\mathbf{p}_8^7)^{-1}(\mathbf{x}^{\circ}) \in \Delta^{7,8}$. We obtain a point

$\tilde{\mathbf{x}}^\circ = \mathbf{p}_8^7(\text{pert}_7^{8,8}(\mathbf{x}_8^\circ)) \in \Delta^7$ so that $f_l(\tilde{\mathbf{x}}^\circ) < \alpha_*$ for every $l \in I - \{2\}$ and $f_l(\tilde{\mathbf{x}}^\circ) = \alpha_*$ for $l = 2$. Then we use the parametrization $\mathbf{p}_1^7$ or $\mathbf{p}_3^7$. We apply the perturbation $\text{pert}_3^{2,1}$ or $\text{pert}_1^{2,3}$ at $\mathbf{x}_1 = (\mathbf{p}_1^7)^{-1}(\tilde{\mathbf{x}}^\circ) \in \Delta^{7,1}$ or $\mathbf{x}_3 = (\mathbf{p}_3^7)^{-1}(\tilde{\mathbf{x}}^\circ) \in \Delta^{7,3}$. We obtain a point $\mathbf{z} = \mathbf{p}_3^7(\text{pert}_1^{2,3}(\mathbf{x}_3)) \in \Delta^7$ so that $f_l(\mathbf{z}) < \alpha_*$ for every $l \in I$, i.e., $F^1(\mathbf{z}) < \alpha_*$. This contradicts with the minimality of α_* . Hence, we cannot have $\Sigma_2^* < 1/2$.

(vi) Let us assume that $\Sigma_2^* = 1/2$. Then we get $\rho_2^* = 1$ and $x_1^* = x_3^*$. We solve the equality $\rho_2^* \sigma_1^* = \alpha_*$ for x_1^* , we see that $x_1^* = 1/(\alpha_* + 1)$. Using the expressions $\rho_1^* \sigma_2^* = \alpha_*$ and $x_1^* = x_3^*$, we get $(\alpha_*^2 - 1)(x_2^*)^2 - (\alpha_* - 1)^2 x_2^* + 2 = 0$. We solve the previous quadratic equation for x_2^* . We find that

$$x_2^* = \frac{(\alpha_* - 1)^2 \pm \sqrt{(\alpha_* - 1)(\alpha_*^3 - 3\alpha_*^2 - 5\alpha_* - 9)}}{2(\alpha_*^2 - 1)}.$$

Then we see that either $x_2^* = h^+(\alpha_*)$ or $x_2^* = h^-(\alpha_*)$ where the functions h^+ and h^- are defined in Lemma 4.7. We plug x_1^* into the equation $\rho_1^* \sigma_8^* = \alpha_*$. Then we solve for x_8^* . We find that

$$x_8^* = -\frac{1 - \alpha_* + (1 + \alpha_*)x_2^*}{3\alpha_* - 1 + (\alpha_*^2 - 1)x_2^*}.$$

Remember that we have $x_1^* = x_7^*$, $x_1^* = x_3^*$ and $\Sigma_2^* = 1/2$. Thus, using the fact that $\Sigma_0^* = 1$, we see that

$$\frac{1}{2} = \frac{(\alpha_* + 1)(\alpha_*^2 - 1)(x_2^*)^2 + (3(\alpha_*^2 - 1) + 2(\alpha_*^2 - 1))x_2^* + \alpha_*^2 + 9\alpha_* - 4}{(\alpha_* + 1)(\alpha_*^2 - 1)x_2^* + (\alpha_* + 1)(3\alpha_* - 1)}$$

for $\alpha_* \in I_\alpha$. In other words, we get either $H^+(\alpha_*) = 1/2$ or $H^-(\alpha_*) = 1/2$ which is a contradiction. Because, in Lemma 4.8, we showed that $H^\pm(\alpha) \neq 1/2$ for every $\alpha \in I_\alpha$. Hence, we conclude that $\Sigma_2^* \neq 1/2$.

(vii) Let us assume that $\Sigma_2^* > 1/2$. We will show that $x_8^* < 1/4 < \Sigma_1^*$. Assume that $1/4 \leq x_8^*$. Note that if we have $x_8^* = 1/4$, then we get $\Sigma_1^* > 1/4$. Thus, we see that $\Sigma_1^* + \Sigma_2^* + x_8^* > 1/2 + 1/4 + 1/4 = 1$. This is a contradiction. If we have $1/4 < x_8^* < \Sigma_1^*$, then we calculate that $\sigma_8^* < 3$ and $\rho_1^* < 3$. In other words, we get $\alpha_* = f_3(\mathbf{x}^*) < 9$. We know that $\alpha_* \in I_\alpha$. Therefore, we get a contradiction.

Let us assume that $\Sigma_1^* \leq 1/4$. Then we see that $\rho_1^* \geq 3$. We get $3\sigma_8^* \leq \rho_1^* \sigma_8^* = \alpha_*$ which implies that $3/(\alpha_* + 3) \leq x_8^*$ so that we derive $(24 - 9\sqrt{2})/46 \leq x_8^* < \Sigma_1^*$. Therefore, we have $(24 - 9\sqrt{2})/23 \leq \Sigma_1^* + x_8^*$. Since we assumed that $1/2 < \Sigma_2^*$, we obtain $(71 - 18\sqrt{2})/46 < \Sigma_1^* + \Sigma_2^* + x_8^*$. Thus, we get $x_7^* < (-25 + 18\sqrt{2})/46$. Then we consider the inequality $\rho_2^*(49 + 36\sqrt{2}) < \rho_2^* \sigma_7^* < \alpha_*$ and solve for Σ_2^* . We find that $\Sigma_2^* > (54 - 11\sqrt{2})/42$. Finally, we conclude that $\Sigma_1^* + \Sigma_2^* = (873 - 221\sqrt{2})/483 > 1$, which is a contradiction. Therefore, we have $x_8^* < 1/4 < \Sigma_1^*$. This implies that

$q_2(1/4) < 0$ or $(1/16)(\alpha_* - 1) + (1/4)(1 - \alpha_*)(1 - \Sigma_4^*) + \Sigma_4^* < 0$. Using the fact that $\alpha_* \in I_\alpha$, we obtain that $\Sigma_4^* < (21 + 18\sqrt{2})/92$.

We assume that $1/2 < \Sigma_2^*$. Therefore, we see that $q_3(1/2) < 0$. In other words, we have $(1/4)(\alpha_* - 1) + (1/2)(1 - \alpha_*)(1 - \Sigma_5^*) + \Sigma_5^* < 0$. Using the fact that $\alpha_* \in I_\alpha$, we find that $\Sigma_5^* < (1 + \sqrt{2})/6$. Then we obtain the following inequality $\Sigma_0^* = \Sigma_4^* + \Sigma_5^* < (109 + 100\sqrt{2})/276 < 1$, a contradiction. Since we have a contradiction in any of the cases (v), (vi), and (vii), Case IV is not possible at $\mathbf{x}^*$. $\square$

We showed that Cases I, II, III and IV are not possible at $\mathbf{x}^*$. We are ready to prove Lemma 4.2.

Proof of Lemma 4.2 Assume that $\mathbf{y} \notin \Delta_7$ where $F^\dagger(\mathbf{y}) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$. Then there exists a $j_0 \in I$ such that $f_{j_0}(\mathbf{y}) < \alpha_*$. Let $N_i = \{j \in I_i : f_j(\mathbf{y}) < \alpha_*\}$ for every $i \in \{1, 2, 3\}$. Since we assume that $f_{j_0}(\mathbf{y}) < \alpha_*$ and $f_j(\mathbf{y}) = \alpha_*$ for some $j \in I - \{j_0\}$, there is a nonempty set $N \subset I$ so that $|N \cap N_i| = 1$ for every i provided that $N_i \neq \emptyset$. Without loss of generality, we can assume that $j_0 \in N$. Assume that $|N| = 1$. Then we see that $N \subset N_i$ for some i and $N \cap N_j = \emptyset$ for every $j \in \{1, 2, 3\} - \{i\}$. Note that we cannot have Case I, I_1 or II. We apply one or two perturbations from the list $\mathcal{L}_i$. We find a point $\mathbf{x}^* \in \Delta^7$ such that $f_j(\mathbf{x}^*) < \alpha_*$ for every $j \in I_i$ and $f_j(\mathbf{x}^*) = \alpha_*$ for every $j \in \mathcal{I}_i$, i.e., $F^\dagger(\mathbf{x}^*) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$. This means that we have either Case III, IV or IV_1 at the point $\mathbf{x}^*$. By Lemma 4.9 and Lemma 4.8, this is a contradiction.

Assume that $|N| = 2$. Then we conclude that there are two different indices i, j so that $N \cap N_i \neq \emptyset$ and $N \cap N_j \neq \emptyset$. We apply one or two perturbations from each list $\mathcal{L}_i$ and $\mathcal{L}_j$. We obtain a point $\mathbf{x}^* \in \Delta^7$ so that $f_k(\mathbf{x}^*) < \alpha_*$ for every $k \in \mathcal{I}_r$ and $f_k(\mathbf{x}^*) = \alpha_*$ for every $k \in I_r$ where $r \in \{1, 2, 3\} - \{i, j\}$. We get either Case I, I_1 or II which is a contradiction by Lemma 4.14.

Let us assume that $|N| = 3$. Then we apply one or two perturbations from each list $\mathcal{L}_1, \mathcal{L}_2$, and $\mathcal{L}_3$. In this way, we obtain a point $\mathbf{x}^* \in \Delta^7$ such that $f_j(\mathbf{x}^*) < \alpha_*$ for every $j \in I$. This contradicts with the minimality of α_* . By the argument above, we conclude that $|N| = 0$. But this contradicts with the statement that $j_0 \in N$ which shows that there is no such j_0 . Therefore, if we have $F^\dagger(\mathbf{y}) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$ for some $\mathbf{y} \in \Delta^7$, then we conclude that $\mathbf{y} \in \Delta_7$. In particular, we must have $\mathbf{x}^* \in \Delta_7$. $\square$

4.1.6 Calculations of the infimums

In this section, we use Lemma 4.2 to calculate the infimum of $G^\dagger$ over the simplex Δ^7 .

Lemma 4.13 *If $F^\dagger$ is the function defined in §4.1, then $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = 5 + 3\sqrt{2}$.*

Proof By Lemma 4.2, we know that $\mathbf{x}^* \in \Delta_7$. This means that we have

$$\begin{aligned} \rho_2^* \sigma_1^* &= \alpha_*, & \rho_{2,3}^* \sigma_2^* &= \alpha_*, & \rho_{1,3}^* \sigma_3^* &= \alpha_*, & \rho_1^* \sigma_4^* &= \alpha_*, \\ \rho_{1,3}^* \sigma_5^* &= \alpha_*, & \rho_{2,3}^* \sigma_6^* &= \alpha_*, & \rho_2^* \sigma_7^* &= \alpha_* & \rho_1^* \sigma_8^* &= \alpha_*. \end{aligned}$$

Using $\rho_1^* \sigma_8^* = \rho_1^* \sigma_4^*$, $\rho_2^* \sigma_7^* = \rho_2^* \sigma_1^*$, $\rho_{2,3}^* \sigma_2^* = \rho_{2,3}^* \sigma_6^*$ and $\rho_{1,3}^* \sigma_3^* = \rho_{1,3}^* \sigma_5^*$, we get $x_4^* = x_8^*$, $x_1^* = x_7^*$, $x_6^* = x_2^*$, and $x_3^* = x_5^*$. Using $\rho_1^* \sigma_4^* = \alpha_*$ and $\rho_2^* \sigma_1^* = \alpha_*$, we see that

$$\sigma(x_1^* + x_2^* + x_3^*)\sigma(x_4^*) = \sigma(x_4^* + x_2^* + x_3^*)\sigma(x_1^*).$$

We simplify the equality above. We obtain $(x_1^* - x_4^*)(x_2^* + x_3^*)(1 - x_1^* - x_2^* - x_3^* - x_4^*) = 0$, which implies that $x_1^* = x_4^*$. Therefore, we get that $\rho_{2,3}^* = \rho_{1,3}^*$. Then by the equalities $\rho_{2,3}^* \sigma_2^* = \alpha_*$ and $\rho_{1,3}^* \sigma_3^* = \alpha_*$, we see that $x_2^* = x_3^*$. In other words, we obtain $x_1^* = x_4^*$, $x_4^* = x_7^*$, $x_7^* = x_8^*$, $x_2^* = x_3^*$, $x_3^* = x_5^*$, and $x_5^* = x_6^*$. Since we know that $\Sigma_0^* = 1$, we have $x_1^* + x_2^* = 1/4$ or $x_2^* = 1/4 - x_1^*$. Using the equalities $\rho_2^* \sigma_1^* = \alpha_*$ and $\rho_{2,3}^* \sigma_2^* = \alpha_*$, we see that

$$\sigma(x_1^* + 2x_2^*)\sigma(x_1^*) = \frac{\sigma(x_2^*)}{\sigma(x_1^* + 2x_2^*)}.$$

We simplify the equation above. We get the following relation between x_1^* and x_2^* :

$$x_2^* - 3x_2^*x_1^* - (x_1^*)^2x_2^* - 4(x_2^*)^2 + 4(x_2^*)^2x_1^* + 4(x_2^*)^3 - (x_1^*)^3 = 0.$$

Then we substitute $x_2^* = 1/4 - x_1^*$. We obtain the equation $1 - 4x_1^* - 4(x_1^*)^2 = 0$. We solve this quadratic equation for x_1^* . We find that $x_1^* = (-1 \pm \sqrt{2})/2$. We need to choose the positive solution for x_1^* . Then we calculate that $x_2^* = (3 - 2\sqrt{2})/4$. In particular, we obtain $\mathbf{x}^* = (x_i^*)$ where $x_i^* = x_j^*$ for every $i, j \in \{1, 4, 7, 8\}$ and $x_r^* = x_s^*$ for every $r, s \in \{2, 3, 5, 6\}$. We calculate that $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = F^\dagger(\mathbf{x}^*)$ so that $F^\dagger(\mathbf{x}^*) = f_i(\mathbf{x}^*)$ for every $i \in I$ where $f_1(\mathbf{x}^*) = 5 + 3\sqrt{2}$. $\square$

Lemma 4.14 *If $G^\dagger$ is the function defined in §3.3, then $\inf_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x}) = 5 + 3\sqrt{2}$.*

Proof The argument we used in §4.1 to show that $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$ can be repeated to show that $\inf_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x})$. Let $\mathbf{z}^* \in \Delta^7$ be a point at which $G^\dagger$ takes its minimum value. Then we get $G^\dagger(\mathbf{x}^*) \geq G^\dagger(\mathbf{z}^*)$. We know that $f_i(\mathbf{x}^*) = 5 + 3\sqrt{2}$ for every $i \in I$ because we have $x_k^* = x_j^*$ for every $k, j \in \{1, 4, 7, 8\}$ and $x_r^* = x_s^*$ for every $r, s \in \{2, 3, 5, 6\}$ where $x_1^* = (\sqrt{2} - 1)/2$ and

$x_2^* = (3 - 2\sqrt{2})/4$. We use the same bijection $\sigma^\dagger: \Psi^\dagger \rightarrow I$ with which we introduced the set of functions $\{f_i\}_{i \in I}$ to express the functions $\{g_j\}$ for $j \in I - \{7, 8\}$. We get

$$g_1 = \rho^7 \sigma_1, g_2 = \rho^6 \sigma_2, g_3 = \rho^5 \sigma_3, g_4 = \rho^8 \sigma_4, g_5 = \rho^3 \sigma_5, g_6 = \rho^2 \sigma_6.$$

Then we calculate that $g_i(\mathbf{x}^*) = 1$ for every $i \in I - \{7, 8\}$. Therefore, we see that $5 + 3\sqrt{2} = G^\dagger(\mathbf{x}^*) \geq G^\dagger(\mathbf{z}^*)$. We know that $G^\dagger(\mathbf{z}^*) \geq 5 + 3\sqrt{2}$ because $\{f_i\}_{i=1}^8 \subset \mathcal{F}^\dagger$. Hence, we conclude that $G^\dagger(\mathbf{z}^*) = 5 + 3\sqrt{2}$. $\square$

4.2 Proof of Proposition 4.1

In this section, we give the proof of Proposition 4.1 which requires the inequalities similar to the ones used in the proof of Proposition 1.5.

Proof We know that $\text{dist}(z_0, \gamma \cdot z_0) = \text{dist}(z_0, \gamma^{-1} \cdot z_0)$ for $\gamma \in \Gamma_\dagger$. Let us say $\mathbf{m}_\dagger = (m_{\sigma^\dagger(\psi)})_{\psi \in \Psi^\dagger} \in \Delta^7$ where $m_{\sigma^\dagger(\psi)}$ is the total measure defined in §3.3. Since the function $(1/2) \log t$ is strictly increasing, we see that

$$\max_{\gamma \in \Gamma_\dagger} (\text{dist}(z_0, \gamma \cdot z_0)) \geq \frac{1}{2} \log G^\dagger(\mathbf{m}_\dagger) \geq \frac{1}{2} \log \left(\inf_{\mathbf{x} \in \Delta^7} G^\dagger(\mathbf{x}) \right) = \frac{1}{2} \log (5 + 3\sqrt{2}),$$

by the discussion given in §3.3 and Lemma 4.14. As a result, we find that

$$e^{2 \max_{\gamma \in \Gamma_\dagger} \{\text{dist}(z_0, \gamma \cdot z_0)\}} \geq \alpha_\dagger = 5 + 3\sqrt{2}$$

which completes the proof of Proposition 4.1. $\square$

We would like to emphasize the similarities between the following two statements

- C. $\inf_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$,
- D. There exists $\mathbf{x}^* \in \Delta^7 \subset \Delta^7$ such that $\min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = f_i^\dagger(\mathbf{x}^*)$ for $i \in I$,

used in the proof of Lemma 4.13 to calculate the number $(1/2) \log(5 + 3\sqrt{2})$ and the statements (A) and (B) listed in §2.3. Although it is straight forward to observe the fact in (B), it takes considerable amount of calculations to prove the statement given in (D). Analogous to Lemma 2.1, Lemma 4.13 shows that the point $\mathbf{x}^*$ is unique. Assuming the uniqueness of the point $\mathbf{x}^*$ a priori together with (C) suggests an alternative way of finding the coordinates of the point $\mathbf{x}^* \in \Delta^7$.

Let $T_1: \mathbb{R}^8 \rightarrow \mathbb{R}^8$ be the transformation defined by $x_1 \leftrightarrow x_4$, $x_2 \leftrightarrow x_5$, $x_3 \leftrightarrow x_6$ and $x_7 \leftrightarrow x_8$. Note that the simplex Δ^7 is preserved by the transformation T_1 . By the definition of T_1 , we have $f_i(T_1(\mathbf{x})) = f_j(\mathbf{x})$ for every $\mathbf{x} \in \Delta^7$ for every pair

$(i, j) \in \{(1, 4), (4, 1), (3, 5), (5, 3), (7, 8), (8, 7), (2, 6), (6, 2)\}$. In other words, we may conclude that $\{f_i : i \in I\} = \{f_i \circ T_1 : i \in I\}$. Let us define $H_1 : \Delta^7 \rightarrow \mathbb{R}$ where $H_1(\mathbf{x}) = \max\{(f_i \circ T_1)(\mathbf{x}) : i \in I\}$. We see that $F^\dagger(\mathbf{x}) = H_1(\mathbf{x})$ for every $\mathbf{x} \in \Delta^7$ and $\min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^7} H_1(\mathbf{x})$. Since $F^\dagger$ takes its minimum value at the point $\mathbf{x}^*$ and $\{f_i(\mathbf{x}^*) : i \in I\} = \{(f_i \circ T_1)(\mathbf{x}^*) : i \in I\}$, H_1 takes its minimum value at the point $T_1^{-1}(\mathbf{x}^*)$. We assumed that $\mathbf{x}^*$ is unique. Hence, we obtain that $T_1^{-1}(\mathbf{x}^*) = \mathbf{x}^*$. This means that $x_1^* = x_4^*, x_2^* = x_5^*, x_3^* = x_6^*$ and $x_7^* = x_8^*$. Let us introduce $\Delta_1^7 = \{\mathbf{x} \in \Delta^7 : x_1 = x_4, x_2 = x_5, x_3 = x_6, x_7 = x_8\}$. Note that we have $f_1(\mathbf{x}) = f_4(\mathbf{x})$, $f_2(\mathbf{x}) = f_5(\mathbf{x})$, $f_3(\mathbf{x}) = f_6(\mathbf{x})$ and $f_7(\mathbf{x}) = f_8(\mathbf{x})$ for every $\mathbf{x} \in \Delta_1^7$. Let us define the continuous function

$$\begin{aligned} F_1^\dagger : \Delta_1^7 &\rightarrow \mathbb{R} \\ \mathbf{x} &\mapsto \max(\bar{f}_1(\mathbf{x}), \bar{f}_2(\mathbf{x}), \bar{f}_3(\mathbf{x}), \bar{f}_7(\mathbf{x})). \end{aligned}$$

where $\bar{f}_i = f_i|_{\Delta_1^7}$ for $i = 1, 2, 3, 7$. Then we have $\min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta_1^7} F_1^\dagger(\mathbf{x})$. Therefore, it is enough to calculate the minimum of $F_1^\dagger$ over the simplex Δ_1^7 .

Let us consider the transformation $T_2 : \mathbb{R}^8 \rightarrow \mathbb{R}^8$ defined by $x_2 \leftrightarrow x_3$ and $x_i \leftrightarrow x_i$ for every $i \in I - \{2, 3\}$. Note that the two sets Δ^7 and Δ_1^7 are preserved by T_2 . By definition of T_2 , we have $\bar{f}_i(T_2(\mathbf{x})) = \bar{f}_j(\mathbf{x})$ for every $\mathbf{x} \in \Delta_1^7$ for every pair $(i, j) \in \{(1, 1), (2, 3), (3, 2), (7, 7)\}$. Let us define the function $H_2 : \Delta^7 \rightarrow \mathbb{R}$ where $H_2(\mathbf{x}) = \max\{(f_i \circ T_2)(\mathbf{x}) : i = 1, 2, 3, 7\}$. We see that $F_1^\dagger(\mathbf{x}) = H_2(\mathbf{x})$ for every $\mathbf{x} \in \Delta_1^7$ and $\min_{\mathbf{x} \in \Delta_1^7} F_1^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta_1^7} H_2(\mathbf{x})$. Since $F_1^\dagger$ takes its minimum value at the point $\mathbf{x}^*$ and $\{f_i(\mathbf{x}^*) : i = 1, 2, 3, 7\} = \{(f_i \circ T_2)(\mathbf{x}^*) : i = 1, 2, 3, 7\}$, H_2 takes its minimum value at the point $T_2^{-1}(\mathbf{x}^*)$. We assumed that $\mathbf{x}^*$ is unique. Hence, we obtain that $T_2^{-1}(\mathbf{x}^*) = \mathbf{x}^*$. This means that $x_2^* = x_3^*$. In other words, we find that $\mathbf{x}^* = (x_1^*, x_2^*, x_2^*, x_1^*, x_2^*, x_2^*, x_7^*, x_7^*)$ where $2x_1^* + 4x_2^* + 2x_7^* = 1$.

Let us define $\Delta_2^7 = \{\mathbf{x} \in \Delta_1^7 : x_2 = x_3\}$. Note that $\bar{f}_2(\mathbf{x}) = \bar{f}_3(\mathbf{x})$ for every $\mathbf{x} \in \Delta_2^7$. Let us define the functions $\tilde{f}_i : \Delta_2^7 \rightarrow \mathbb{R}$ for $i = 1, 2, 7$ where $\tilde{f}_i = \bar{f}_i|_{\Delta_2^7}$ for $i = 1, 2, 7$. Then we introduce the following continuous function

$$\begin{aligned} F_2^\dagger : \Delta_2^7 &\rightarrow \mathbb{R} \\ \mathbf{x} &\mapsto \max(\tilde{f}_1(\mathbf{x}), \tilde{f}_2(\mathbf{x}), \tilde{f}_7(\mathbf{x})). \end{aligned}$$

Since we have $\min_{\mathbf{x} \in \Delta_1^7} F_1^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta_2^7} F_2^\dagger(\mathbf{x})$, it is enough to calculate the minimum of $F_2^\dagger$ over the simplex Δ_2^7 .

Let us define $\Delta = \{(x_1, x_2) \in \mathbb{R}^2 : 0 < x_1 < 1/2, 0 < x_2 < 1/2(1/2 - x_1)\}$. Note that we have $(x_1^*, x_2^*) \in \Delta$. We introduce the parametrization $\mathbf{p} : \Delta \rightarrow \Delta_2^7$ defined by

$(x_1, x_2) \mapsto (x_1, x_2, x_2, x_1, x_2, x_2, 1/2 - x_1 - 2x_2, 1/2 - x_1 - 2x_2)$. Then we obtain the commuting diagram below

$$\begin{array}{ccc} \Delta & & \\ \downarrow \mathfrak{p} & \searrow F & \\ \Delta_2^7 & \xrightarrow{F_2^\dagger} & \mathbb{R} \end{array}$$

where $F(x_1, x_2) = \max_{(x_1, x_2) \in \Delta} \{f_i(x_1, x_2) : i = 1, 2, 7\}$ for the following functions $f_1 = \sigma(x_1 + 2x_2)\sigma(x_1)$, $f_2 = \sigma(x_2)/\sigma(x_1 + 2x_2)$ and $f_7 = \sigma(x_1 + 2x_2)\sigma(1/2 - x_1 - 2x_2)$. By the diagram above, we derive that $F(x^*) = F_2^\dagger(\mathbf{x}^*)$ such that $x^* = \mathfrak{p}^{-1}(\mathbf{x}^*)$. Note that we reduced the calculation of the minimum of $F^\dagger$ over the simplex Δ^7 to the calculation of the minimum of F over a two-dimensional region Δ .

Let us calculate the partial derivatives of the functions f_1 and f_2 . We find the followings

$$\begin{aligned} (f_1)_1 &= 2\tau(x_1)\tau(x_1 + 2x_2)(x_1^2 + 2x_1x_2 - x_1 - x_2 + 2x_2^2), \\ (f_1)_2 &= -2\tau(x_1 + 2x_2)\sigma(x_1), \\ (f_2)_1 &= -\tau(1 - x_1 - 2x_2)\sigma(x_2), \\ (f_2)_2 &= \tau(x_2)\tau(1 - x_1 - 2x_2)(4x_1x_2 + 2x_2^2 - x_1 + x_1^2). \end{aligned}$$

Note that we have $(f_1)_2(x_1, x_2) < 0$ for every $(x_1, x_2) \in \Delta$. Therefore, f_1 has no critical point in Δ . Similarly, f_2 has no critical point in Δ since we have $(f_2)_1(x_1, x_2) > 0$ for every $(x_1, x_2) \in \Delta$. On the contrary, every point on the set $L = \{(x_1, x_2) \in \Delta : -1 + 4x_1 + 8x_2 = 0\}$ is a critical point for the function f_7 because we have $(f_7)_1 = \tau(x_1 + 2x_2)\tau(-1 + 2x_1 + 4x_2)(-1 + 4x_1 + 8x_2)$, $(f_7)_2 = -2(f_7)_1$. Let $g_i|_L = f_i$ for $i = 1, 2, 7$ where $L \subset \Delta$. Then we get

$$g_1(x_2) = 3 \cdot \frac{3 + 8x_2}{1 - 8x_2}, \quad g_2(x_2) = \frac{1}{3} \cdot \frac{1 - x_2}{x_2}, \quad g_7(x_2) = 9$$

for $0 < x_2 < 1/8$. Let us say $I = (0, 1/8)$. Then g_1 and g_2 have no critical points in I since $g_1'(x_2) = 96\tau(-1 + 8x_2) > 0$ and $g_2'(x_2) = -\tau(\sqrt{3}x_2) < 0$ for every $x_2 \in I$. We see that $g_1(x_2) > g_7(x_2)$ for every x_2 in the interval I because of the following limits

$$\lim_{x_2 \rightarrow 0^+} 3 \cdot \frac{3 + 8x_2}{1 - 8x_2} = 9^+ \quad \text{and} \quad \lim_{x_2 \rightarrow \frac{1}{8}^-} 3 \cdot \frac{3 + 8x_2}{1 - 8x_2} = \infty.$$

Therefore, it is enough to calculate the infimum of the maximum of g_1 and g_2 over the interval I to calculate the infimum of the maximum of f_1 , f_2 and f_7 over the set L . Since g_1 and g_2 have no critical point in I , the infimum of the maximum of g_1 and g_2 is attained at a point t^* such that $g_1(t^*) = g_2(t^*)$. In other words, we need to solve the equation $64x_2^2 + 36x_2 - 1 = 0$. Then, we get that $t^* = (-9 \pm \sqrt{97})/32$. We

need to pick the positive solution. We calculate that $g_1(t^*) = (17 + 2\sqrt{97})/3$ where $g_1(t^*) = \inf_{x_2 \in I} \{\max(g_1(x_2), g_2(x_2))\}$. Note that the point t^* at which the infimum of the maximum of g_1 and g_2 over L is unique.

We claim that there exists $i, j \in \{1, 2, 7\}$ where $i \neq j$ such that $f_i(x^*) = f_j(x^*)$. Let us assume otherwise that $f_i(x^*) \neq f_j(x^*)$ for every $i, j \in \{1, 2, 7\}$ where $i \neq j$. Then we have either $f_1(x^*) > f_i(x^*)$ for $i = 2, 7$ or $f_2(x^*) > f_i(x^*)$ for $i = 1, 7$ or $f_7(x^*) > f_i(x^*)$ for $i = 1, 2$. Since f_1 and f_2 have no critical points in Δ , we cannot have $f_1(x^*) > f_i(x^*)$ for $i = 2, 7$ and $f_2(x^*) > f_i(x^*)$ for $i = 1, 7$. If we assume that $f_7(x^*) > f_i(x^*)$ for $i = 1, 2$, then there exists a neighborhood U of x^* in Δ so that $f_7(x_1, x_2) > f_i(x_1, x_2)$ for every $(x_1, x_2) \in U$ for $i = 1, 2$ because f_7 is continuous on Δ . Hence, we get $F(x_1, x_2) = f_7(x_1, x_2)$ for every $(x_1, x_2) \in U$. Since we have $f_7(x^*) = F(x^*)$, the function f_7 has a local minimum at $x^* \in U$. This means that $x^* \in L$. In other words, we find that $x^* = ((13 - \sqrt{97})/16, -(9 + \sqrt{97})/32)$. This is a contradiction. Because we know that $F(\bar{x}) = 5 + 3\sqrt{2} < \tilde{F}(x^*)$ where we have $\bar{x} = ((\sqrt{2} - 1)/2, (3 - 2\sqrt{2})/4) \in \Delta$. Hence, there exists $i, j \in \{1, 2, 3\}$ where $i \neq j$ such that $f_i(x^*) = f_j(x^*)$. We need to consider the cases $f_1(x^*) = f_7(x^*) > f_2(x^*)$, $f_1(x^*) = f_2(x^*) > f_7(x^*)$, $f_7(x^*) = f_2(x^*) > f_1(x^*)$ and $f_1(x^*) = f_2(x^*) = f_7(x^*)$.

Let us assume that $f_1(x^*) = f_7(x^*) > f_2(x^*)$. Let $\vec{v}_1 = \langle 1, -1/2 \rangle$. The first order partial derivatives of f_1 , f_2 and f_7 show that

$$D_{\vec{v}_1}f_1(x_1, x_2) < 0, \quad D_{\vec{v}_1}f_7(x_1, x_2) = 0, \quad D_{\vec{v}_1}f_2(x_1, x_2) > 0$$

for every $(x_1, x_2) \in \Delta$. Therefore, we conclude that f_1 is decreasing, f_2 is increasing and f_7 is constant on the line segment $\ell_1 \cap \Delta$ where $\ell_1 = \{\vec{x}^* + t\vec{v}_1 : t \in \mathbb{R}\}$. Since f_2 is continuous on $\ell_1 \cap \Delta$ and $f_2(x^*) < \alpha_*$, there exists an $\epsilon_1 > 0$ so that $f_2(x_1, x_2) < \alpha_*$ for every $(x_1, x_2) \in \ell_1^{\epsilon_1} = \{\vec{x}_0^* + t\vec{v}_1 : t \in (0, \epsilon_1)\}$. Let (y_1, y_2) be a point on $\ell_1^{\epsilon_1}$. Then we get $f_2(y_1, y_2) < \alpha_*$. Since f_1 is decreasing on the line segment $\ell_1^{\epsilon_1} \subset \ell_1$ and $f_1(x^*) = \alpha_*$, we see that $f_1(y_1, y_2) < \alpha_*$. This means that if we move along the line segment starting at x^* in the direction of the vector $\vec{v}_1$ for a sufficiently small amount, we obtain a point $(y_1, y_2) \in \Delta$ such that $f_i(y_1, y_2) < \alpha_*$ for $i = 1, 2$ and $f_7(y_1, y_2) = \alpha_*$. This is a contradiction. Therefore, we cannot have $f_1(x^*) = f_7(x^*) > f_2(x^*)$.

Let us assume that $f_2(x^*) = f_7(x^*) > f_1(x^*)$. Let $\vec{v}_2 = -\vec{v}_1$. The first order partial derivatives of f_1 , f_2 and f_7 show that

$$D_{\vec{v}_2}f_1(x_1, x_2) > 0, \quad D_{\vec{v}_2}f_7(x_1, x_2) = 0, \quad D_{\vec{v}_2}f_2(x_1, x_2) < 0$$

for every $(x_1, x_2) \in \Delta$. Therefore, we conclude that f_2 is decreasing, f_1 is increasing and f_7 is constant on the line segment $\ell_2 \cap \Delta$ where $\ell_2 = \{\vec{x}^* + t\vec{v}_2 : t \in \mathbb{R}\}$. Since f_1 is continuous on $\ell_2 \cap \Delta$ and $f_1(x^*) < \alpha_*$, there exists an $\epsilon_2 > 0$ so that $f_1(x_1, x_2) < \alpha_*$

for every $(x_1, x_2) \in \ell_2^{\epsilon_2} = \{\vec{x}^* + t\vec{v}_2 : t \in (0, \epsilon_2)\}$. Let (y_1, y_2) be a point on $\ell_2^{\epsilon_2}$. Then we get $f_1(y_1, y_2) < \alpha_*$. Since f_2 is decreasing on the line segment $\ell_2^{\epsilon_2} \subset \ell_2$ and $f_2(x^*) = \alpha_*$, we see that $f_2(y_1, y_2) < \alpha_*$. This means that if we move along the line segment starting at x^* in the direction of the vector $\vec{v}_2$ for a sufficiently small amount, we obtain a point $(y_1, y_2) \in \Delta$ such that $f_i(y_1, y_2) < \alpha_*$ for $i = 1, 2$ and $f_7(y_1, y_2) = \alpha_*$. This is a contradiction. Therefore, we cannot have $f_2(x^*) = f_7(x^*) > f_1(x^*)$. There are only two cases left to consider:

$$\text{I. } f_1(x^*) = f_2(x^*) > f_7(x^*), \quad \text{II. } f_2(x^*) = f_7(x^*), \quad f_7(x^*) = f_1(x^*).$$

We know that Case I is not possible at the point x^* . Because this case corresponds to Case III in Lemma 4.5. Therefore, we have $f_2(x^*) = f_7(x^*)$ and $f_7(x^*) = f_1(x^*)$. Using the equality $f_1(x^*) = f_2(x^*)$, we see that $\sigma(x_1^*) = \sigma(1/2 - x_1^* - 2x_2^*)$. When we simplify the previous equality, we find that $x_1^* + x_2^* = 1/4$ or $x_1^* = 1/4 - x_2^*$. We use the equation $f_2(x^*) = f_7(x^*)$ together with $x_1^* = 1/4 - x_2^*$. We obtain that

$$\frac{1}{4}(x_2^*)^2 - \frac{3}{8}(x_2^*) + \frac{1}{64} = 0 \quad \text{or} \quad x_2^* = \frac{3 \pm 2\sqrt{2}}{4}.$$

We need to choose x_2^* as $(3 - 2\sqrt{2})/4$ because we have $(3 + 2\sqrt{2})/4 > 1$. Then we find that $x_1^* = (2 - \sqrt{2})/2$. In other words, we get the point $x^* = ((\sqrt{2} - 1)/2, (3 - 2\sqrt{2})/4)$. Finally, we calculate that $f_i(x^*) = 5 + 3\sqrt{2}$ for $i = 1, 2, 7$. In particular, we obtain that $\min_{(x_1, x_2) \in \Delta} F(x_1, x_2) = 5 + 3\sqrt{2}$ which implies that

$$\min_{\mathbf{x} \in \Delta_2^7} F_2^\dagger(\mathbf{x}) = \min_{(x_1, x_2) \in \Delta} F(x_1, x_2).$$

We know that $\min_{\mathbf{x} \in \Delta_1^7} F_2^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta_2^7} F_1^\dagger(\mathbf{x})$ and $\min_{\mathbf{x} \in \Delta_1^7} F_1^\dagger(\mathbf{x}) = \min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x})$. Thus we conclude that

$$\min_{\mathbf{x} \in \Delta^7} F^\dagger(\mathbf{x}) = 5 + 3\sqrt{2}$$

which is attained at the point $\mathbf{x}^* = (x_i^*)$ where $x_i^* = x_j^*$ for $i, j \in \{1, 4, 7, 8\}$ and $i, j \in \{2, 3, 5, 6\}$.

In the discussion above, we don't refer to the statement $\mathbf{x}^* \in \Delta_7$ given in (D). The assumption that the point $\mathbf{x}^*$ is unique reduces the necessary calculations to obtain α_* considerably. We would like to emphasize the fact that $\Gamma_{\mathcal{D}^\dagger}$ is not a symmetric decomposition of Γ . We will leave the investigation of the likely conditions such as the convexity properties of the displacement functions for the decompositions $\Gamma_{\mathcal{D}^\dagger}$ and $\Gamma_{\mathcal{D}^\dagger}$ that might lead to a proof of the uniqueness of the points $\mathbf{x}^*$ to future studies.

5 Proof of The Main Theorem

In this final section, we present a detailed proof of the main result of this paper.

Theorem 5.1 *Let ξ and η be non-commuting isometries of $\mathbb{H}^3$. Suppose that ξ and η generate a torsion-free discrete group which is not co-compact and contains no parabolic. Let $\Gamma_{\dagger}$ and $\alpha_{\dagger}$ denote the set of isometries $\{\xi, \eta, \xi\eta\}$ and the real number $5 + 3\sqrt{2}$, respectively. Then for any $z_0 \in \mathbb{H}^3$ we have*

$$e^{(2 \max_{\gamma \in \Gamma_{\dagger}} \{\text{dist}(z_0, \gamma \cdot z_0)\})} \geq \alpha_{\dagger}.$$

Proof Let $\Gamma = \langle \xi, \eta \rangle$. Since Γ is a torsion-free discrete group which is not co-compact and contains no parabolic, it is a free group on the generators ξ and η ([9], Proposition 9.2). Then there are two cases to consider $\Lambda_{\Gamma} = S_{\infty}$ or $\Lambda_{\Gamma} \neq S_{\infty}$.

Let us assume that $\Lambda_{\Gamma} = S_{\infty}$. Since Γ is topologically tame and $\Lambda_{\Gamma} = S_{\infty}$, every superharmonic function on the manifold $M = \mathbb{H}^3/\Gamma$ is constant ([9], Proposition 6.8 and [4], Theorem 7.2) which implies that every Γ -invariant superharmonic function is constant on $\mathbb{H}^3$. Then every Γ -invariant D -conformal density $\mathcal{M}$ is a constant multiple of the area density $\mathcal{A}$ ([9], Proposition 3.9). Hence, Proposition 4.1 implies that

$$e^{(2 \max_{\gamma \in \Gamma_{\dagger}} \{\text{dist}(z_0, \gamma \cdot z_0)\})} \geq \alpha_{\dagger},$$

which gives the conclusion of Theorem 5.1 in the case $\Lambda_{\Gamma} = S_{\infty}$.

Let us assume that $\Lambda_{\Gamma} \neq S_{\infty}$. Let V denote the variety $PSL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$ with the complex topology. Remember that $\mathfrak{D}$ is the set of all points $(\xi', \eta') \in V$ such that the group $\langle \xi', \eta' \rangle$ generated by ξ' and η' is a free Kleinian group of rank 2. We know that $\mathfrak{D}$ is a closed subset of V ([2]). The set of all points $(\xi', \eta') \in \mathfrak{D}$ such that $\Lambda_{\Gamma'} \neq S_{\infty}$ where $\Gamma' = \langle \xi', \eta' \rangle$ is denoted by $\mathfrak{G}\mathfrak{F}$ which is an open subset of V ([5], Theorem 8.1). The frontier $\mathfrak{B}$ of $\mathfrak{G}\mathfrak{F}$ contains a dense subset $\mathfrak{C}$ of elements $(\xi', \eta') \in \mathfrak{B}$ so that $\Lambda_{\Gamma'} = S_{\infty}$ ([9], Theorem 8.2). Let z_0 be a given point in $\mathbb{H}^3$. Let us consider the function

$$\begin{aligned} f_{z_0}^{\dagger} : \quad \mathfrak{D} &\longrightarrow \mathbb{R} \\ (\xi', \eta') &\longmapsto \max_{\gamma \in \Gamma'_{\dagger}} \{\text{dist}(z_0, \gamma \cdot z_0)\} \end{aligned}$$

where $\Gamma'_{\dagger} = \{\pi(\gamma) : \gamma \in \Gamma_{\dagger}\}$ defined in §1.4.

Since $(f_{z_0}^{\dagger})^{-1}([0, B]) = \{(\xi', \eta') \in V : \max_{\gamma \in \{\xi', \eta'\}} \{\text{dist}(z_0, \gamma \cdot z_0)\} \leq B\}$ is compact for every $B \geq 0$, the function $f_{z_0}^{\dagger}$ is a continuous, proper, non-negative valued function on V . Therefore, $f_{z_0}^{\dagger}$ must take a minimum value on $\overline{\mathfrak{G}\mathfrak{F}}$. Let $(\xi_0, \eta_0) \in \overline{\mathfrak{G}\mathfrak{F}}$ be a point at which $f_{z_0}^{\dagger}$ takes a minimum value. The fact that every superharmonic function on $M = \mathbb{H}^3/\Gamma'$ is constant for every $(\xi', \eta') \in \mathfrak{C}$ implies that $\mathcal{M}_{\Gamma', z_0} = \mathcal{A}$ for every $(\xi', \eta') \in \mathfrak{C}$ where $\Gamma' = \langle \xi', \eta' \rangle$. Then Proposition 4.1 gives the inequality

$f_{z_0}^\dagger(\xi', \eta') \geq (1/2) \log(5 + 3\sqrt{2})$ for every $(\xi', \eta') \in \mathfrak{C}$. If (ξ_0, η_0) is in $\mathfrak{B}$, then we also have $f_{z_0}^\dagger(\xi_0, \eta_0) \geq (1/2) \log(5 + 3\sqrt{2})$ since $\mathfrak{C}$ is a dense subset of $\mathfrak{B}$. Hence, it is enough to show that $f_{z_0}^\dagger$ has no local minimum on the open set $\mathfrak{GF}$.

Let us assume that (ξ_0, η_0) is in $\mathfrak{GF}$. Then $f_{z_0}^\dagger$ has a local minimum at (ξ_0, η_0) . Since ξ_0, η_0 and $\xi_0\eta_0$ have infinite orders in $\langle \xi_0, \eta_0 \rangle$, we have $\xi_0 \cdot z \neq z$, $\eta_0 \cdot z \neq z$ and $\xi_0\eta_0 \cdot z \neq z$ for every $z \in \mathbb{H}^3$. In particular, we get that $\xi_0 \cdot z_0 \neq z_0$, $\eta_0 \cdot z_0 \neq z_0$ and $\xi_0\eta_0 \cdot z_0 \neq z_0$. Therefore, there exists hyperbolic geodesic segments joining z_0 to $\xi_0 \cdot z_0$, z_0 to $\eta_0 \cdot z_0$ and z_0 to $\xi_0\eta_0 \cdot z_0$. Note that we have the equalities $\text{dist}(z_0, \xi_0\eta_0 \cdot z_0) = \text{dist}(\xi_0^{-1} \cdot z_0, \eta_0 \cdot z_0)$ and $\text{dist}(z_0, \xi \cdot z_0) = \text{dist}(z_0, \xi^{-1} \cdot z_0)$. We consider the geodesic triangle $\triangle_{P_2P_0P_1}$ where $P_1 = \xi_0^{-1} \cdot z_0$, $P_0 = z_0$ and $P_2 = \eta_0 \cdot z_0$. Let $d_1 = \text{dist}(P_1, P_0)$, $d_2 = \text{dist}(P_2, P_0)$, $d_{1,2} = \text{dist}(P_1, P_2)$ and $\theta_{1,2}$ be the angle between the line segments P_0P_1 and P_0P_2 .

Let $G_{z_0}^\dagger: \mathbb{H}^3 \times \mathbb{H}^3 \rightarrow \mathbb{R}_+^2$ be the map defined by $(P, Q) \mapsto (\text{dist}(P, z_0), \text{dist}(Q, z_0))$. We examine the restriction $H_{z_0}^\dagger$ of $G_{z_0}^\dagger$ onto disjoint union of the geodesics C_i starting at z_0 and passing through the points P_i for $i = 1, 2$. If P'_i denotes a point on the geodesic C_i for $i = 1, 2$, then the formula for $H_{z_0}^\dagger$ is given with the following formula $(P'_1, P'_2) \mapsto (\text{dist}(P'_1, P_0), \text{dist}(P'_2, P_0))$. Note that $H_{z_0}^\dagger$ is continuous, one-to-one and onto. Let us consider the map $F_1: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined such that if $\mathbf{x} = (x_1, x_2)$ is a point in $\mathbb{R}^2$, then the restriction of F_1 onto each and all of the coordinate axis in $\mathbb{R}^2$ is given by one and only one of the functions with the formula $\mathbf{x} \mapsto f_{i,j}(\mathbf{x})$ for $(i, j) \in \{(1, 2), (2, 1)\}$ where $f_{i,j}(x_1, x_2) = \cosh x_i \cosh x_j - \sinh x_i \sinh x_j \cos \theta_{i,j}$. Note that we have $f_{1,2}(\mathbf{x}) = f_{2,1}(\mathbf{x})$ and $f_{1,2}(\mathbf{d}) = d_{1,2}$ for every $\mathbf{x} \in \mathbb{R}^2$ where $\mathbf{d} = (d_1, d_2)$. We derive that

$$\begin{aligned} (f_{1,2})_1(\mathbf{x}) &= \sinh x_1 \cosh x_2 - \cosh x_1 \sinh x_2 \cos \theta_{1,2}, \\ (f_{1,2})_2(\mathbf{x}) &= \cosh x_1 \sinh x_2 - \sinh x_1 \cosh x_2 \cos \theta_{1,2}. \end{aligned}$$

Let us assume that $d_1 = d_2$. Then we derive that $0 < \theta_{1,2} < \pi$ because we have $\xi_0^{-1} \cdot z_0 \neq \eta_0 \cdot z_0$ and $\xi_0 \cdot z_0 \neq \eta_0 \cdot z_0$. We know that $\sinh 2d_1 > 0$ and $1 - \cos \theta_{1,2} > 0$. As a consequence, we find that $(f_{1,2})_1(\mathbf{d}) + (f_{1,2})_2(\mathbf{d}) = \sinh 2d_1(1 - \cos \theta_{1,2}) > 0$. Let $\vec{v}_1 = \langle -1, -1 \rangle$. We calculate the directional derivative $D_{\vec{v}_1} F_1(\mathbf{d}) = \langle v_{i,j}^1 \rangle$ of F_1 in the direction of $\vec{v}_1$ at the point $\mathbf{d}$ where we determine the coordinates of the vector $v_{i,j}^1$ as

$$v_{i,j}^1 = \begin{cases} -(f_{i,j})_1(\mathbf{d}) - (f_{i,j})_2(\mathbf{d}) & \text{if } (i, j) = (1, 2) \\ -(f_{i,j})_1(\mathbf{d}) - (f_{i,j})_2(\mathbf{d}) & \text{if } (i, j) = (2, 1). \end{cases}$$

The directional derivative $D_{\vec{v}_1} F_1(\mathbf{d})$ shows that, for $\epsilon > 0$ however small, there is a point $\mathbf{d}_1 = (d'_1, d'_2)$ on the half line given as $\ell_1 = \{\vec{\mathbf{d}} + t\vec{v}_1: t \in \mathbb{R}_+\} \subset \mathbb{R}_+^2$ such that

$0 < d_i - d'_i < \epsilon$ for $i = 1, 2$ and $f_{i,j}(\mathbf{d}_1) < f_{i,j}(\mathbf{d})$ for every $(i, j) \in \{(1, 2), (2, 1)\}$ as the functions $f_{i,j}$ are increasing in the direction of $\vec{v}_1$. As a result, we obtain a point $(P'_1, P'_2) \in \mathbb{H}^3 \times \mathbb{H}^3$ so that $H_{z_0}^\dagger(P'_1, P'_2) = \mathbf{d}_1$ since $H_{z_0}^\dagger$ is a continuous bijection. In other words, we see that the inequalities $\text{dist}(P_1, P'_1) < \epsilon$, $\text{dist}(P'_2, P_2) < \epsilon$ and $\text{dist}(P'_1, P'_2) < \text{dist}(P_1, P_2)$ are all satisfied with $\text{dist}(P_0, P'_1) = \text{dist}(P_0, P'_2)$.

Let us consider the case $d_1 > d_2$. Then we have $0 \leq \theta_{1,2} \leq \pi$. We study the partial derivative $(f_{1,2})_1$. We observe that

$$(22) \quad -\sinh(x_2 - x_1) = \frac{e^{x_2} + e^{-x_2}}{2} \cdot \frac{e^{x_1} - e^{-x_1}}{2} - \frac{e^{x_2} - e^{-x_2}}{2} \cdot \frac{e^{x_1} + e^{-x_1}}{2} \leq (f_{1,2})_1.$$

Since we set $d_1 > d_2$, we find that $-\sinh(d_2 - d_1) > 0$. Using the inequality in (22), we obtain that $(f_{1,2})_1(\mathbf{d}) > 0$. Then we calculate the directional derivative $D_{\vec{v}_2} F_1(\mathbf{d}) = \langle v_{i,j}^2 \rangle$ of F_1 in the direction of the vector $\vec{v}_2 = \langle -1, 0 \rangle$ at the point $\mathbf{d} = (d_1, d_2)$ where the entries $v_{i,j}^2$ are given as follows

$$v_{i,j}^2 = \begin{cases} -(f_{i,j})_1(\mathbf{d}) & \text{if } (i, j) = (1, 2), \\ -(f_{i,j})_1(\mathbf{d}) & \text{if } (i, j) = (2, 1). \end{cases}$$

The directional derivative $D_{\vec{v}_2} F_1(\mathbf{d})$ shows that, for $\epsilon_1 > 0$ however small, there is a point $\mathbf{d}_1 = (d'_1, d_2)$ on the half line $\ell_2 = \{\vec{\mathbf{d}} + t\vec{v}_2 : t \in \mathbb{R}_+\} \subset \mathbb{R}_+^2$ such that $0 < d_1 - d'_1 < \epsilon_1$ and $f_{i,j}(\mathbf{d}_1) < f_{i,j}(\mathbf{d})$ for every $(i, j) \in \{(1, 2), (2, 1)\}$ as the functions $f_{i,j}$ are increasing in the direction of $\vec{v}_2$. As a result, we obtain a point $(P'_1, P_2) \in \mathbb{H}^3 \times \mathbb{H}^3$ so that $H_{z_0}^\dagger(P'_1, P_2) = \mathbf{d}_1$ because $H_{z_0}^\dagger$ is a continuous bijection. In other words, we see that $\text{dist}(P'_1, P_2) < \text{dist}(P_1, P_2)$ and $\text{dist}(P_1, P'_1) < \epsilon_1$. We can choose $\epsilon_1 > 0$ so that $d'_1 > d_2$.

Let $\vec{v}_3 = \langle 0, -1 \rangle$. We calculate the directional derivative $D_{\vec{v}_3} F_1(\mathbf{d}_1) = \langle v_{i,j}^3 \rangle$ of F_1 in the direction $\vec{v}_3$ at the point $\mathbf{d}_1$ where

$$v_{i,j}^3 = \begin{cases} -(f_{i,j})_2(\mathbf{d}_1) & \text{if } (i, j) = (1, 2), \\ -(f_{i,j})_2(\mathbf{d}_1) & \text{if } (i, j) = (2, 1). \end{cases}$$

Note that it may be the case that $(f_{i,j})_2(\mathbf{d}_1) < 0$. We know by the previous step that $f_{i,j}(\mathbf{d}_1) < f_{i,j}(\mathbf{d})$ for $(i, j) \in \{(1, 2), (2, 1)\}$. Since $f_{1,2}$ and $f_{2,1}$ are continuous functions, for $\epsilon_2 > 0$ however small, there is a point $\mathbf{d}_2 = (d'_1, d'_2)$ on the half line $\ell_3 = \{\vec{\mathbf{d}}_1 + t\vec{v}_3 : t \in \mathbb{R}_+\} \subset \mathbb{R}_+^2$ so that $0 < d_2 - d'_2 < \epsilon_2$ and $f_{i,j}(\mathbf{d}_2) < f_{i,j}(\mathbf{d})$ for $(i, j) \in \{(1, 2), (2, 1)\}$. Using the fact that $H_{z_0}^\dagger$ is a continuous bijection, we obtain a point $(P'_1, P'_2) \in \mathbb{H}^3 \times \mathbb{H}^3$ so that $H_{z_0}^\dagger(P'_1, P'_2) = \mathbf{d}_2$. Therefore, we conclude that the inequalities $\text{dist}(P_1, P'_1) < \epsilon$, $\text{dist}(P_2, P'_2) < \epsilon$ and $\text{dist}(P'_1, P'_2) < \text{dist}(P_1, P_2)$ where $\epsilon = \max\{\epsilon_1, \epsilon_2\}$ in the case $d_1 > d_2$.

Let us consider the case $d_2 > d_1$. We find a lower bound for the partial derivative $(f_{2,1})_2$. We see that

$$(23) \quad -\sinh(x_1 - x_2) = \frac{e^{x_1} + e^{-x_1}}{2} \cdot \frac{e^{x_2} - e^{-x_2}}{2} - \frac{e^{x_1} - e^{-x_1}}{2} \cdot \frac{e^{x_2} + e^{-x_2}}{2} \leq (f_{2,1})_2.$$

Since we assume $d_2 > d_1$, we have the inequality $-\sinh(d_1 - d_2) > 0$. Using the inequality in (23), we derive that $(f_{2,1})_2(\mathbf{d}) > 0$. We calculate the directional derivative $D_{\vec{v}_3} F_1(\mathbf{d}) = v_{i,j}^4$ of F_1 in the direction of $\vec{v}_3$ at the point $\mathbf{d}$ where

$$v_{i,j}^4 = \begin{cases} -(f_{i,j})_2(\mathbf{d}) & \text{if } (i,j) = (1,2), \\ -(f_{i,j})_2(\mathbf{d}) & \text{if } (i,j) = (2,1). \end{cases}$$

We derive from the directional derivative $D_{\vec{v}_3} F_1(\mathbf{d})$ that, for $\epsilon_1 > 0$ however small, there is a point $\mathbf{d}_1 = (d_1, d'_1)$ on the half line $\ell_4 = \{\vec{\mathbf{d}} + t\vec{v}_3 : t \in \mathbb{R}_+\} \subset \mathbb{R}_+^2$ such that $0 < d_2 - d'_2 < \epsilon_1$ and $f_{i,j}(\mathbf{d}_1) < f_{i,j}(\mathbf{d})$ for $(i,j) \in \{(1,2), (2,1)\}$ as the functions $f_{i,j}$ are increasing in the direction of the vector $\vec{v}_3$. Using the fact that $H_{z_0}^\dagger$ is a continuous bijection, we obtain a point $(P_1, P'_2) \in \mathbb{H}^3 \times \mathbb{H}^3$ such that $H_{z_0}^\dagger(P_1, P'_2) = \mathbf{d}_1$. In other words, we have $\text{dist}(P_1, P'_2) < \text{dist}(P_1, P_2)$ and $\text{dist}(P_2, P'_2) < \epsilon_1$. We can choose $\epsilon_1 > 0$ so that $d'_2 > d_1$.

Finally, we calculate the directional derivative $D_{\vec{v}_2} F_1(\mathbf{d}_1) = \langle v_{i,j}^3 \rangle$ of F_1 in the direction $\vec{v}_2$ at the point $\mathbf{d}_1$ where

$$v_{i,j}^5 = \begin{cases} -(f_{i,j})_1(\mathbf{d}_1) & \text{if } (i,j) = (1,2), \\ -(f_{i,j})_1(\mathbf{d}_1) & \text{if } (i,j) = (2,1). \end{cases}$$

Note that it may be the case that $(f_{i,j})_1(\mathbf{d}_1) < 0$. We know by the previous step that $f_{i,j}(\mathbf{d}_1) < f_{i,j}(\mathbf{d})$ for $(i,j) \in \{(1,2), (2,1)\}$. Since $f_{1,2}$ and $f_{2,1}$ are continuous functions, for $\epsilon_2 > 0$ however small, there is a point $\mathbf{d}_2 = (d'_1, d'_2)$ on the half line $\ell_5 = \{\vec{\mathbf{d}}_1 + t\vec{v}_2 : t \in \mathbb{R}_+\} \subset \mathbb{R}_+^2$ so that $0 < d_2 - d'_2 < \epsilon_2$ and $f_{i,j}(\mathbf{d}_2) < f_{i,j}(\mathbf{d}_1)$ for $(i,j) \in \{(1,2), (2,1)\}$. Using the fact that $H_{z_0}^\dagger$ is a continuous bijection, we obtain a point $(P'_1, P'_2) \in \mathbb{H}^3 \times \mathbb{H}^3$ so that $H_{z_0}^\dagger(P'_1, P'_2) = \mathbf{d}_2$. We conclude that the inequalities $\text{dist}(P_1, P'_1) < \epsilon$, $\text{dist}(P_2, P'_2) < \epsilon$ and $\text{dist}(P'_1, P'_2) < \text{dist}(P_1, P_2)$ where $\epsilon = \max\{\epsilon_1, \epsilon_2\}$ in the case $d_1 > d_2$. Hence, in any case $d_1 = d_2$, $d_1 > d_2$ or $d_2 > d_1$, there are points P'_1 and P'_2 which are arbitrarily close to points P_1 and P_2 on the geodesics C_1 and C_2 so that $\text{dist}(P'_1, P'_2) < \text{dist}(P_1, P_2)$.

Let ξ_1 and η_1 be two isometries of $\mathbb{H}^3$ so that $\xi_1^{-1} \cdot z_0 = P'_1$ and $\eta_1^{-1} \cdot z_0 = P'_2$. Since the points P'_1 and P'_2 are chosen arbitrarily close to the points P_1 and P_2 , (ξ_1, η_1) is arbitrarily close the point $(\xi_0, \eta_0) \in \mathfrak{GF}$. We know that $\mathfrak{GF}$ is an open subset of V . Hence, we get (ξ_1, η_1) is in $\mathfrak{GF}$. Then we derive that the expression $\text{dist}(z_0, \xi_1^{-1} \cdot z_0) = \text{dist}(P_0, P'_1) < \text{dist}(P_0, P_1) = \text{dist}(z_0, \xi_0 \cdot z_0)$ where we know that

$\text{dist}(z_0, \xi_1 \cdot z_0) = \text{dist}(z_0, \xi_1^{-1} \cdot z_0)$. Similarly, we have $\text{dist}(z_0, \eta_1 \cdot z_0) = \text{dist}(P_0, P'_2)$. Therefore, we find that the inequality $\text{dist}(P_0, P'_2) < \text{dist}(P_0, P_2) = \text{dist}(z_0, \eta_0 \cdot z_0)$ holds. We also have $\text{dist}(z_0, \xi_1 \eta_1 \cdot z_0) = \text{dist}(\xi_1^{-1} \cdot z_0, \eta_1 \cdot z_0)$. Then we see that

$$\text{dist}(\xi_1^{-1} \cdot z_0, \eta_1 \cdot z_0) = \text{dist}(P'_1, P'_2) < \text{dist}(P_1, P_2) = \text{dist}(\xi_0^{-1} \cdot z_0, \eta_0 \cdot z_0).$$

This means that we obtain a point $(\xi_1, \eta_1) \in \mathfrak{G}\mathfrak{F}$ so that $f_{z_0}^\dagger(\xi_1, \eta_1) < f_{z_0}^\dagger(\xi_0, \eta_0)$ which contradicts with the fact that $f_{z_0}^\dagger$ has a local minimum at the point (ξ_0, η_0) . We conclude that (ξ_0, η_0) is not in $\mathfrak{G}\mathfrak{F}$. This completes the proof. $\square$

A Proofs of Lemmas for the Decomposition $\Gamma_{\mathcal{D}^1}$

In this appendix we complete the proofs of Lemma 4.3, Lemma 4.4, Lemma 4.5, and Lemma 4.9 which show that Cases I, II, III and IV are not possible at a point where $F^\dagger: \Delta^7 \rightarrow \mathbb{R}$ takes its minimum value by proving Lemma 4.6 required for Case III, and Lemmas 4.7, 4.8, 4.10, 4.11, and 4.12 required for Case IV.

A.1 Proof of Lemma 4.6

We know that $x_2^* \in I = (0, 1/6)$. Remember that $g_\alpha(x) = (1-\alpha)^2 x^2 + \alpha$. First, we find an upper and a lower bound for $g_\alpha(x)$. Let $\tilde{F}(x, \alpha) = (1-\alpha)^2 x^2 + \alpha$ defined on $\mathbb{R} \times \mathbb{R}$ where $F(x, \alpha)$ is the restriction of $\tilde{F}$ on $I \times I_\alpha$. For any fixed $\alpha \in I_\alpha$, we calculate that $\lim_{x \rightarrow 0^+} F(x, \alpha) = \alpha$ and $\lim_{x \rightarrow 1/6^-} F(x, \alpha) = (1-\alpha)^2 (1/6)^2 + \alpha$. Therefore, $F(x, \alpha)$ is bounded over $I \times I_\alpha$. Note that we have $F_x(x, \alpha) = 2(1-\alpha)x > 0$ and $F_\alpha(x, \alpha) = -2(1-\alpha)x^2 + 1 > 0$ for every $(x, \alpha) \in I \times \text{int}(I_\alpha)$. Also, we determine that $\tilde{F}'(x, 9) = 128x > 0$ and $\tilde{F}'(x, 5+3\sqrt{2}) = 2(4+3\sqrt{2})^2 x > 0$ for every $x \in I$. Hence, we conclude that $\tilde{F}(0, 9) = 9 < g_\alpha(x) < (107/18) + (11/3)\sqrt{2} = \tilde{F}(1/6, 5+3\sqrt{2})$ over $I \times I_\alpha$.

Let $\tilde{G}(x, \alpha) = h_\alpha(x) + 2x$ defined on $\mathbb{R} \times \mathbb{R}$ where $G(x, \alpha)$ is the restriction of $\tilde{G}$ on $I \times I_\alpha$. For α in I_α , we calculate that $\lim_{x \rightarrow 0^+} G(x, \alpha) = (-1 + \sqrt{\alpha})/(\alpha - 1) > 0$ and $\lim_{x \rightarrow 1/6^-} G(x, \alpha) = (-5 - \alpha + 6(F(1/6, \alpha))^{1/2})(6\alpha - 6)^{-1} + 1/3 > 0$. Therefore, $G(x, \alpha)$ is bounded over $I \times I_\alpha$. We have $G_x(x, \alpha) = 1 + x(-1 + \alpha)F^{-1/2} > 0$ and $G_\alpha(x, \alpha) = (\alpha - 1)^{-2}((\alpha - 1)x^2 + 1/2)F^{-1/2} > 0$ for $(x, \alpha) \in I \times \text{int}(I_\alpha)$. For any $x \in I$, we also calculate the following derivatives $\tilde{G}'(x, 9) = 1 + 8x(F(x, 9))^{-1/2} > 0$ and $\tilde{G}'(x, 5+3\sqrt{2}) = 1 + (4+3\sqrt{2})x(F(x, 5+3\sqrt{2}))^{-1/2} > 0$. As a result, we determine that $\tilde{G}(0, 9) < G(x, \alpha) < \tilde{G}(1/6, 5+3\sqrt{2})$. In particular, we find that

$$0 < \frac{1}{4} < h_\alpha(x) + 2x < \frac{-2 + 3\sqrt{2} + \sqrt{214 + 132\sqrt{2}}}{6(4 + 3\sqrt{2})} < 1.$$

This means that $H_\alpha(x) > 0$ for every $(x, \alpha) \in I \times I_\alpha$.

We aim to show that for any given fixed $\alpha \in I_\alpha$, we have $H'_\alpha(x) < 0$ for every $x \in I$. Let us calculate the derivative of H_α using logarithmic differentiation:

$$\frac{H'_\alpha}{H_\alpha} = \frac{(h'_\alpha + 2)(1-x)x - (1-h_\alpha-2x)(h_\alpha+2x)}{(1-h_\alpha-2x)(h_\alpha+2x)(1-x)x}.$$

Note that we have $(1-h_\alpha-2x)(h_\alpha+2x)(1-x)x > 0$ for every $(x, \alpha) \in I \times I_\alpha$. We find that $\lim_{x \rightarrow 0^+} ((h'_\alpha(x) + 2)(1-x)x - (1-h_\alpha(x)-2x)(h_\alpha(x)+2x)) < 0$ because $\lim_{x \rightarrow 0^+} G_x(x, \alpha) = 1$. Since the numerator of H'_α/H_α is continuous, it is enough to show that the equation

$$(h'_\alpha(x) + 2)(1-x)x - (1-h_\alpha(x)-2x)(h_\alpha(x)+2x) = 0$$

has no solution on $I \times I_\alpha$. Let us assume on the contrary that there is a solution $(x_0, \alpha_0) \in I \times I_\alpha$. In particular, we see that

$$(1+x_0(-1+\alpha_0))(g_{\alpha_0})^{-1/2}(1-x_0)x_0 = (1-h_{\alpha_0}(x_0)-2x_0)(h_{\alpha_0}(x_0)+2x_0).$$

We simplify the equality above as follows: We expand both sides of the equation. We collect all the terms containing $g_{\alpha_0}(x_0)$ and $(g_{\alpha_0}(x_0))^{1/2}$ as factors on the left hand-side of the equation and the rest of the terms on the right hand-side. Then we substitute $(1-\alpha_0)^2x_0^2 + \alpha_0$ for the term $g_{\alpha_0}(x_0)$ but not for the term $(g_{\alpha_0}(x_0))^{1/2}$ on the left hand-side. Next, we collect all the similar terms together on the both sides of the equation. We see that the term $((\alpha_0-1)^2x_0^2 - 2(\alpha_0-1)x_0 + 2\alpha_0)$ is a factor on the left hand-side. We divide both sides by this factor. Then we obtain

$$\sqrt{g_{\alpha_0}(x_0)} = \frac{-(\alpha_0-1)^3x_0^3 + 2(\alpha_0-1)^2x_0^2 + 2\alpha_0(1-\alpha_0)x_0 + \alpha_0^2 + \alpha_0}{(\alpha_0-1)^2x_0^2 - 2(\alpha_0-1)x_0 + 2\alpha_0}.$$

Let us introduce the function $\tilde{Q}(x, \alpha) = r(x, \alpha)/s(x, \alpha)$ on $R = [0, \infty) \times [1, \infty)$ where we declare $r(x, \alpha) = -(\alpha-1)^3x^3 + 2(\alpha-1)^2x^2 + 2\alpha(1-\alpha)x + \alpha^2 + \alpha$ and $s(x, \alpha) = (\alpha-1)^2x^2 - 2(\alpha-1)x + 2\alpha$. Note that we have $s(x, \alpha) > 0$ for every $(x, \alpha) \in I \times I_\alpha \subset R$. Let $Q(x, \alpha)$ denote the restriction of $\tilde{Q}$ on $I \times I_\alpha$. For any fixed $\alpha \in I_\alpha$, we calculate that $\lim_{x \rightarrow 0^+} Q(x, \alpha) = (\alpha+1)/2$ and

$$\lim_{x \rightarrow \frac{1}{\alpha}^-} Q(x, \alpha) = -\frac{\alpha^3 - 159\alpha^2 - 261\alpha - 13}{6(\alpha^2 + 58\alpha + 13)}.$$

Therefore, we see that $Q(x, \alpha)$ is bounded on $I \times I_\alpha$. Let us determine the sign of the partial derivative $Q_x(x, \alpha)$ on $I \times I_\alpha$. We consider the expression below which is a factor of the numerator of $Q_x(x, \alpha)$:

$$\frac{r_x s - r s_x}{-(\alpha-1)^2} = (\alpha-1)^3x^4 - 4(\alpha-1)^2x^3 + 4(\alpha^2-1)x^2 + 2\alpha(\alpha-3)x + 2\alpha.$$

We have $x \in I$. We get $x^3 < x^2$ which implies that $4(\alpha^2 - 1)x^3 < 4(\alpha^2 - 1)x^2$. We consider the expressions $4(\alpha^2 - 1)$ and $4(\alpha - 1)^2$. We find the inequality $4(\alpha^2 - 1)x^3 > 4(\alpha - 1)^2x^3$ since $4(\alpha^2 - 1) - 4(\alpha - 1)^2 = 8(\alpha - 1) > 0$. In other words, we obtain that $4(\alpha^2 - 1)x^2 - 4(\alpha - 1)^2x^3 > 0$. Since we have the expression $(\alpha - 1)^3x^4 + 2\alpha(\alpha - 3)x + 2\alpha > 0$, we derive that $Q_x(x, \alpha) < 0$ for every $(x, \alpha) \in I \times \text{int}(I_\alpha)$. The argument above can be repeated to show that $\tilde{Q}'(x, 9) < 0$ and $\tilde{Q}'(x, 5 + 3\sqrt{2}) < 0$. As a result, we get the bounds

$$\left(\frac{107}{18} + \frac{11}{3}\sqrt{2}\right)^{1/2} < \frac{907}{231} = \tilde{Q}\left(\frac{1}{6}, 9\right) < Q(x, \alpha) < \tilde{Q}\left(0, 5 + 3\sqrt{2}\right) = \frac{6 + 3\sqrt{2}}{2}$$

for every $(x, \alpha) \in I \times I_\alpha$. In particular, the inequality above is true for the point (x_0, α_0) where we have $Q(x_0, \alpha_0) = (g_{\alpha_0}(x_0))^{1/2}$. In other words, we derive that $g_{\alpha_0}(x_0) > 107/18 + (11/3)\sqrt{2}$, a contradiction. Hence, we conclude that $H'_\alpha(x) < 0$ for every $(x, \alpha) \in I \times I_\alpha$ or H_α is a strictly decreasing function over the interval $(0, 1/6)$ for every $\alpha \in (9, 5 + 3\sqrt{2})$.

A.2 Proofs of Lemma 4.7 and 4.8

In this section, we will use the notation $(a_n^{q_n}, a_{n-1}^{q_{n-1}}, \dots, a_1^{q_1}, a_0^{q_0})$ to express the sum $\sum_{k=0}^n a_k \alpha^{q_k}$. For example, the notation $(2^3, -1^1, 2^{1/2}, 2^0)$ corresponds to the function $2\alpha^3 - \alpha + 2\alpha^{1/2} + 2$. If there is only one term in a sum, we will drop the parentheses, e.g., α^5 will be denoted by 1^5 . Multiplication of the sums $\sum_{k=0}^n a_k \alpha^{q_k}$ and $\sum_{k=0}^m b_k \alpha^{r_k}$ will be denoted by $(a_n^{q_n}, a_{n-1}^{q_{n-1}}, \dots, a_1^{q_1}, a_0^{q_0})(b_m^{r_m}, b_{m-1}^{r_{m-1}}, \dots, b_1^{r_1}, b_0^{r_0})$. The expression $K(2^3, -1^1, 2^{1/2}, 2^0)$ will mean $2K\alpha^3 - K\alpha + 2K\alpha^{1/2} + 2K$ where $K \neq 0$ is a constant. A power p of a sum $\sum_{k=0}^n a_k \alpha^{q_k}$ will be written as $(a_n^{q_n}, a_{n-1}^{q_{n-1}}, \dots, a_1^{q_1}, a_0^{q_0})^p$. All of the inequalities in this section will be considered over $I_\alpha = [9, 5 + 3\sqrt{2}]$. Generally, we will not preserve the order of the exponents. Let $\alpha_1 = 5 + 3\sqrt{2}$.

A.2.1 Proof of Lemma 4.7

Let us say $R_1 = (1^{7/2}, -1^{5/2}, 2^3, -4^2, -1^{3/2}, 2^1, 1^{1/2})$, $R_2 = (2^{5/2}, 4^2, -4^1, -2^{1/2})$, $R_4 = (-1^2, 1^{3/2}, 4^1, 1^{1/2}, -1^0)$. Let us introduce $R_5 = (-1^{3/2}, -4^1, -1^{1/2}, 2^0)$ and $R_6 = -1^0$. We calculate the derivative of G_+ as

$$G'_+(\alpha) = \frac{(g_+^2 R_1 + g_+ R_2 + R_3)g'_+ + (g_+^2 R_4 + g_+ R_5 + R_6)}{\sqrt{\alpha}(\sqrt{\alpha} + 1)^2(\alpha g_+ - g_+ + 1)^2}.$$

Define the functions $R_7 = (1^6, -2^{11/2}, -2^{9/2}, 9^4, 4^{7/2}, 8^{5/2}, -3^2, 2^{3/2}, 2^{1/2}, 1^0)$ and $R_8 = (-3^8, 8^7, -16^6, -24^5, 22^4, 8^3, 8^1, -3^0)$. We calculate the derivative of g_+ as

$$g'_+ = -\frac{R_7\sqrt{d} + R_8\sqrt{\alpha} + R_9}{4(\alpha^{\frac{7}{2}} + \alpha^2 - \alpha^{\frac{3}{2}} - 1)^2\sqrt{d}\sqrt{\alpha}}$$

where $R_9 = (1^9, -6^8, -12^7, 16^6, -14^5, -12^4, 28^3, -3^1, 2^0)$. Let $d = R_{10} - R_{11}$ so that $R_{10} = (1^6, 4^{9/2}, 6^3, 1^2, 6^{3/2}, 4^{1/2}, 4^0)$ and $R_{11} = (2^{11/2}, 3^5, 6^4, 4^{7/2}, 8^{5/2}, 3^1)$. Both R_{10} and R_{11} are increasing because $R'_{10} = (6^5, 18^{7/2}, 18^2, 2^1, 9^{1/2}, 2^{-1/2}) > 0$ and $R'_{11} = (11^{9/2}, 15^4, 24^3, 14^{5/2}, 20^{3/2}, 3^0) > 0$.

Define the functions $P_1 = (8^{11/2}, -22^5)$, $P_2 = (3^{11/2}, 36^4, 36^{5/2}, -30^{9/2}, -48^{7/2})$, and $P_3 = (1^{11/2}, 4^{3/2}, 18^1, 4^0, -28^3, -40^2, -6^{1/2})$ where $d'(\alpha) = (\alpha^{-1/2}/2) \sum_{i=1}^3 P_i$. We show that $P_i(\alpha) > 0$ for $i = 1, 2, 3$. We see that $P_1 = 1^5(8^{1/2}, -22^0) > 0$ since $\alpha \geq 9$. We calculate the derivative $P'_2 = ((33/2)^{9/2}, 144^3, 90^{3/2}, -135^{7/2}, -168^{5/2})$. We get $P'_2(\alpha) \geq \min((33/2)^{9/2}, 144^3, 90^{3/2}) - \max(135^{7/2}, 168^{5/2}) > 0$ since both of the functions $((33/2)^{9/2}, 144^3, 90^{3/2})$ and $(135^{7/2}, 168^{5/2})$ are clearly increasing. This means that P_2 is increasing. Therefore, we obtain that $P_2(\alpha) \geq P_2(9) > 0$.

We carry out a similar calculation for $P_3(\alpha)$. Note that we have the following inequality $P_3(\alpha) \geq \min_{\alpha \in I_\alpha} (1^{11/2}, 4^{3/2}, 18^1, 4^0) - \max_{\alpha \in I_\alpha} (28^3, 40^2, 6^{1/2}) > 0$ since both of the functions $(1^{11/2}, 4^{3/2}, 18^1, 4^0)$ and $(28^3, 40^2, 6^{1/2})$ are increasing. This means that P_3 is a positive function. Hence, we conclude that $d'(\alpha) > 0$ for every α . Consequently, $d(\alpha)$ is an increasing function.

We show that R_7 is a positive function. Let us write $R_7(\alpha) = r_1(\alpha) - r_2(\alpha)$ where $r_1 = (1^6, 9^4, 4^{7/2}, 8^{5/2}, 2^{1/2}, 1^0)$, and $r_2 = (2^{11/2}, 2^{9/2}, 3^2, 2^{3/2})$. Then we obtain that $R_7(\alpha) \geq \min_{\alpha \in I_\alpha} r_1(\alpha) - \max_{\alpha \in I_\alpha} r_2(\alpha) = r_1(9) - r_2(\alpha_1) > 0$ for every α . Next we examine the expression $R_7 d^{1/2} + R_8 \alpha^{1/2} + R_9$. Since $d(\alpha)$ is increasing, we obtain $R_7 d^{1/2} + R_8 \alpha^{1/2} + R_9 \leq R_7(\max_{\alpha \in I_\alpha} d^{1/2}) + R_8 \alpha^{1/2} + R_9 = KR_7 + R_8 \alpha^{1/2} + R_9$ where $K = 191040 + 135084\sqrt{2} - 54744\alpha_1^{1/2} - 38712\alpha_1^{1/2}\sqrt{2}$. We will show that $KR_7 + R_8 \alpha^{1/2} + R_9 < 0$.

Let us say $KR_7 + R_8 \alpha^{1/2} + R_9 = p_1 - p_2$ where $p_1 = K\tilde{p}_1 + \tilde{p}_2$ and $p_2 = K\bar{p}_1 + \bar{p}_2$ for $\tilde{p}_1 = (1^6, 9^4, 4^{7/2}, 8^{5/2}, 2^{1/2}, 1^0)$, $\tilde{p}_2 = (8^{15/2}, 22^{9/2}, 8^{7/2}, 8^{3/2}, 1^9, 16^6, 28^3, 2^0)$ and $\bar{p}_2 = (3^{17/2}, 16^{13/2}, 24^{11/2}, 3^{1/2}, 6^8, 12^7, 12^4, 14^5, 3^1)$. Then we obtain the inequality $KR_7 + R_8 \alpha^{1/2} + R_9 = p_1(\alpha) - p_2(\alpha) \leq \max_{\alpha \in I_\alpha} p_1(\alpha) - \min_{\alpha \in I_\alpha} p_2(\alpha) < 0$ because both p_1 and p_2 are clearly increasing functions.

The inequality we showed above implies that $g'_+(\alpha) > 0$ for every $\alpha \in \text{int}(I_\alpha)$. Then $g_+(\alpha)$ is an increasing, positive function since $g_+(9) > 0$. Thus, $g_+^2(\alpha)$ is an

increasing function. Then, it is straightforward to obtain the following inequality

$$\begin{aligned} g_+^2 R_1 + g_+ R_2 + R_3 &\leq (\max_{\alpha \in I_\alpha} g_+(\alpha))^2 R_1 + (\max_{\alpha \in I_\alpha} g_+(\alpha)) R_2 + R_3 \\ &= (g_+(\alpha_1))^2 R_1 + g_+(\alpha_1) R_2 + R_3. \end{aligned}$$

Let $K = g_+(\alpha_1) > 0$. We show that $K^2 R_1 + K R_2 + R_3 < 0$ for $\alpha \in I_\alpha$. Let us say $K^2 R_1 + K R_2 + R_3 = q_1 - q_2$ where $q_1 = K^2(1^{7/2}, 2^3, 2^1, 1^{1/2}) + K(2^{5/2}, 4^2) + (2^1, 1^{1/2})$ and $q_2 = K^2(1^{5/2}, 4^2, 1^{3/2}) + K(4^1, 2^{1/2}) + (1^{5/2}, 2^2)$. Then we obtain the inequality $K^2 R_1 + K R_2 + R_3 = q_1(\alpha) - q_2(\alpha) \leq \max_{\alpha \in I_\alpha} q_1(\alpha) - \min_{\alpha \in I_\alpha} q_2(\alpha) < 0$ since both of the functions $q_1(\alpha)$ and $q_2(\alpha)$ are clearly increasing. As a result, we find that $(g^2 R_1 + g R_2 + R_3)g' < 0$.

We show that $g^2 R_4 + g R_5 + R_6 < 0$. It is clear that $R_5 < 0$ and $R_6 < 0$. We have $R_4 = (1^{3/2}, 4^1) - (1^2, 1^0) \leq \max_{\alpha \in I_\alpha} (1^{3/2}, 4^1) - \min_{\alpha \in I_\alpha} (1^2, 1^0) < 0$ since both of the functions $(1^{3/2}, 4^1)$ and $(1^2, 1^0)$ are increasing. The last inequality implies that $g^2 R_4 + g R_5 + R_6 < 0$ because we have $g^2 > 0$ and $g > 0$ for every α . As a consequence, we conclude that the inequality $(g^2 R_1 + g R_2 + R_3)g' + (g^2 R_4 + g R_5 + R_6) < 0$ holds. In particular, we obtain that $(G_+)' < 0$ for every α . Thus, the following calculation gives the result of the first part of the lemma:

$$G_+(\alpha_1) = \frac{(\sqrt{\alpha_1} + 1)(\alpha_1 - 1)(g_+(\alpha_1))^2 + 2(\alpha_1 - 1)g_+(\alpha_1) + \sqrt{\alpha_1} + 3}{(\sqrt{\alpha_1} + 1)((\alpha_1 - 1)g_+(\alpha_1) + 1)} > 1$$

For the second part of the lemma, we study the derivative of the function G_- :

$$(G_-)'(\alpha) = \frac{(g_-^2 R_1 + g_- R_2 + R_3)g'_- + (g_-^2 R_4 + g_- R_5 + R_6)}{\sqrt{\alpha}(\sqrt{\alpha} + 1)^2(\alpha g_- - g_- + 1)^2}.$$

Note that $R_1, R_2, R_3, R_4, R_5, R_6$ are the same functions we used in the first part of the proof. Let us calculate the derivative of $g_-(\alpha)$:

$$g'_- = \frac{\hat{R}_7 \sqrt{d} + \hat{R}_8 \sqrt{\alpha} + \hat{R}_9}{4(\alpha^{\frac{7}{2}} + \alpha^2 - \alpha^{\frac{3}{2}} - 1)^2 \sqrt{d} \sqrt{\alpha}}$$

where $\hat{R}_7(\alpha) = -R_7(\alpha)$, $\hat{R}_8(\alpha) = R_8(\alpha)$, and $\hat{R}_9(\alpha) = R_9(\alpha)$. We already know that $(d(\alpha))^{1/2} > 0$ is an increasing function from the first part. Let us examine the expression $-R_7 \sqrt{d} + R_8 \alpha^{1/2} + R_9$. We know that $d(\alpha)$ is increasing. We obtain $-R_7 d^{1/2} + R_8 \alpha^{1/2} + R_9 \leq -R_7 (\min_{\alpha \in I_\alpha} d^{1/2}) + R_8 \alpha^{1/2} + R_9 = K R_7 + R_8 \alpha^{1/2} + R_9$ where $K = -33280$. We aim to show that $K R_7 + R_8 \alpha^{1/2} + R_9 > 0$.

Consider the equation $K R_7 + R_8 \alpha^{1/2} + R_9 = s_1 - s_2$ where we have the functions $s_1 = -K \tilde{s}_1 + \tilde{s}_2$ and $s_2 = -K \bar{s}_1 + \bar{s}_2$ so that $\bar{s}_1 = (1^6, 9^4, 4^{7/2}, 8^{5/2}, 2^{1/2}, 1^0)$, $\bar{s}_2 = (3^{17/2}, 16^{13/2}, 24^{11/2}, 3^{1/2}, 6^8, 12^7, 12^4, 14^5, 3^1)$, $\tilde{s}_1 = (2^{11/2}, 2^{9/2}, 3^2, 2^{3/2})$ and

$\tilde{s}_2 = (8^{15/2}, 22^{9/2}, 8^{7/2}, 8^{3/2}, 1^9, 16^6, 28^3, 2^0)$. Then we get the following inequality $KR_7 + R_8\alpha^{1/2} + R_9 = s_1 - s_2 \leq \max_{\alpha \in I_\alpha} s_1(\alpha) - \min_{\alpha \in I_\alpha} s_2(\alpha) < 0$ because both s_1 and s_2 are clearly increasing. The last inequality implies that $g'_-(\alpha) < 0$ for every α . We know that $g_-(\alpha)$ is a decreasing, positive function because $g_-(\alpha_1) > 0$. Therefore, $g_-^2(\alpha)$ is a decreasing function. In particular, we obtain

$$\begin{aligned} g_-^2 R_1 + g_- R_2 + R_3 &\geq (\min_{\alpha \in I_\alpha} g_-(\alpha))^2 R_1 + (\min_{\alpha \in I_\alpha} g_-(\alpha)) R_2 + R_3 \\ &= (g_-(\alpha_1))^2 R_1 + g_-(\alpha_1) R_2 + R_3. \end{aligned}$$

Let $K = g_-(\alpha_1) > 0$. We will show that $K^2 R_1 + KR_2 + R_3 < 0$ for every $\alpha \in I_\alpha$. Let $K^2 R_1 + KR_2 + R_3 = t_1 - t_2$ where $t_1 = K^2(1^{7/2}, 2^3, 2^1, 1^{1/2}) + K(2^{5/2}, 4^2) + (2^1, 1^{1/2})$ and $t_2 = K^2(1^{5/2}, 4^2, 1^{3/2}) + K(4^1, 2^{1/2}) + (1^{5/2}, 2^2)$. Then we derive the inequality

$$K^2 R_1 + KR_2 + R_3 = t_1(\alpha) - t_2(\alpha) \geq \min_{\alpha \in I_\alpha} t_1(\alpha) - \max_{\alpha \in I_\alpha} t_2(\alpha) > -381$$

because both of the functions $t_1(\alpha)$ and $t_2(\alpha)$ are clearly increasing functions. We aim to show that $g'_-(\alpha)$ is an increasing function. To this purpose, we calculate $g''_-(\alpha)$:

$$g''_- = \frac{d^{\frac{3}{2}}(Q_1 - Q_2) + Q_3 - Q_4}{8(\alpha^{\frac{7}{2}} + \alpha^2 - \alpha^{\frac{3}{2}} - 1)^3(d)^{\frac{3}{2}}(\alpha)^{\frac{3}{2}}}$$

Define the functions $\tilde{q}_1 = (96^6, 16^{9/2}, 60^4, 36^{7/2}, 8^3, 48^{5/2}) - (24^5, 12^2, 11^{3/2}, 1^0)$, $\tilde{q}_2 = (3^{19/2}, -8^9)$, $\tilde{q}_3 = (72^{15/2}, -15^8)$, and $\tilde{q}_4(\alpha) = 24^7 - (36^{11/2}, 24^5)$. Then we calculate that $Q_1 - Q_2 = \tilde{q}_1 + \tilde{q}_2 + \tilde{q}_3 + \tilde{q}_4$. We also define the functions q_1, q_2 and q_6 where

$$q_1 = (950^{\frac{27}{2}}, 810^{13}, 2194^{12}, 2034^{\frac{23}{2}}, 3336^{\frac{21}{2}}, 3624^{10}, 3090^9, 3072^{\frac{17}{2}}, 2166^{\frac{15}{2}}),$$

$q_2 = (2442^7, 2026^6, 1290^{11/2}, 974^{9/2}, 373^{3/2}, 12^{1/2}, 8^1)$ and $q_6 = 1^{17}(10^1, -3^{1/2})$. Let us say $q_3 = (4386^{11}, 4596^{19/2}, 5346^8, 4326^{13/2})$, $q_5 = 1^{16}(7^2, 15^{3/2}, 118^{1/2}, -132^1)$ and $q_4 = (2226^5, 90^4, 1098^{7/2}, 6^3, 213^{5/2}, 153^2, 6^1)$. Also let us define the function $q_7 = 1^{31/2}(162^{1/2}, -798^1) + 1^{14}(414^1, -2442^0) + 1^{25/2}(438^2, -4038^1)$. Then we see that $Q_3 - Q_4 = (q_1 + q_2) - (q_3 + q_4) + (q_5 + q_6 + q_7)$.

We aim to show that $Q_1(\alpha) - Q_2(\alpha) > 0$ for $\alpha \in I_\alpha$. It is clear that the inequalities $(3^{19/2}, -8^9) = 1^9(3^{1/2}, -8^1) > 1^9$, $(24^6, 12^{9/2}, 11^{7/2}, 1^0) > (24^5, 12^2, 11^{3/2}, 1^0)$, $(96^6, 16^{9/2}, 60^4, 36^{7/2}, 8^3, 48^{5/2}) > (24^6, 12^{9/2}, 11^{7/2}, 1^0)$ hold. Note that we have $(72^{15/2}, -15^8) = 1^{15/2}(72^0, -15^{1/2})$. Hence, we get $\tilde{q}_1 > 0$, $\tilde{q}_2 > 0$ and $\tilde{q}_3 > 0$ for every $\alpha \in I_\alpha$. As a result, for $\alpha \in I_\alpha$, we obtain the following inequality $\tilde{q}_4(\alpha) = 1^5(24^2 - (36^{1/2}, 24^0)) \geq 1^5(\min_{\alpha \in I_\alpha} (24^2) - \max_{\alpha \in I_\alpha} (36^{1/2}, 24^0)) \geq 1^5$ which implies that $\tilde{q}_4(\alpha)$ is also a positive function over I_α . Then we conclude that $Q_1 - Q_2 = \tilde{q}_1 + \tilde{q}_2 + \tilde{q}_3 + \tilde{q}_4 > 0$ for every $\alpha \in I_\alpha$.

We carry out similar calculations to show that the function $Q_3(\alpha) - Q_4(\alpha) > 0$ over I_α . By comparing the terms, we see that $q_1 + q_2 > q_3 + q_4$ for every $\alpha \in I_\alpha$. Since we have $\alpha \geq 3$, we obtain $q_5 > 0$ for every α . We consider the inequality $(7^2, 15^{3/2}, 118^{1/2}, -132^1) \geq (\min_{\alpha \in I_\alpha}(7^2, 15^{3/2}, 118^{1/2}) - \max_{\alpha \in I_\alpha}(132^1)) \geq 1$ which shows that $q_5 > 0$ over I_α . All the functions $(162^{1/2}, -798^1)$, $(414^1, -2442^0)$ and $(438^2, -4038^0)$ are positive functions since $\alpha \geq 3$. Therefore, we find that the inequality $Q_3 - Q_4 = q_1 + q_2 - (q_3 + q_4) + (q_5 + q_6 + q_7) > 0$ holds for every $\alpha \in I_\alpha$. Thus, we conclude that $g'_-(\alpha) > 0$ over the interval I_α which shows that $g'_-(\alpha)$ is an increasing function. Then we calculate that

$$g'_-(\alpha) \geq g'_-(9) = \frac{\hat{R}_7(9)\sqrt{d(9)} + \hat{R}_8(9)\sqrt{9} + \hat{R}_9(9)}{4(9^{\frac{7}{2}} + 9^2 - 9^{\frac{3}{2}} - 1)^2\sqrt{d(9)}\sqrt{9}} \geq -\frac{1}{40}.$$

Let us consider the inequality below

$$\begin{aligned} g_-^2 R_4 + g_- R_5 + R_6 &\geq (\min_{\alpha \in I_\alpha} g_-(\alpha))^2 R_4 + (\min_{\alpha \in I_\alpha} g_-(\alpha)) R_5 + R_6 \\ &= (g_-(\alpha_1))^2 R_4 + g_-(\alpha_1) R_5 + R_6. \end{aligned}$$

Let $K = g_-(\alpha_1) > 0$. We will show that $K^2 R_4 + K R_5 + R_6 > 0$ for every $\alpha \in I_\alpha$. Let us say $K^2 R_4 + K R_5 + R_6 = u_1 - u_2$ where $u_1 = K^2(1^{3/2}, 4^1, 1^{1/2}) + 2K$ and $u_2 = K^2(1^2, 1^0) + K(1^{3/2}, 4^1, 1^{1/2}) + (1^0)$. Note that both of the functions $u_1(\alpha)$ and $u_2(\alpha)$ are clearly increasing over I_α . Therefore, we obtain the following inequality $K^2 R_4 + K R_5 + R_6 = u_1(\alpha) - u_2(\alpha) \geq \min_{\alpha \in I_\alpha} u_1(\alpha) - \max_{\alpha \in I_\alpha} u_2(\alpha) > -6$ for every $\alpha \in I_\alpha$. Then we calculate that

$$(g_-^2 R_1 + g_- R_2 + R_3)g'_- + (g_-^2 R_4 + g_- R_5 + R_6) > (-381) \cdot \left(-\frac{1}{40}\right) + (-6) > 0$$

for every $\alpha \in I_\alpha$. Since we have $\alpha^{1/2}(\alpha^{1/2} + 1)^2(\alpha g_- - g_- + 1)^2 > 0$ for every $\alpha \in I_\alpha$, we obtain $(G_-)'(\alpha) > 0$ for every $\alpha \in I_\alpha$. Thus, the following calculation gives the result of the second part of the lemma:

$$G_-(9) = \frac{(\sqrt{9} + 1)(9 - 1)(h(9))^2 + 2(9 - 1)h(9) + \sqrt{9} + 3}{(\sqrt{9} + 1)((9 - 1)h(9) + 1)} > 1.$$

A.2.2 Proof of Lemma 4.8

Let $d = (1^1, -1^0)(1^3, -3^2, -5^1, -9^0)$, $R_1 = 2((1^1, -1^0))^2(1^4, -2^3, -2^2, -42^1, 5^0)$. Then we calculate $(H_+)'(\alpha)$:

$$(H_+)'(\alpha) = -\frac{R_1\sqrt{d} + R_2}{\sqrt{d}(\alpha^2 + 4\alpha - 1 + \sqrt{d})^2(\alpha + 1)^2(\alpha - 1)^2}$$

where $R_2 = 2((1^1, -1^0))^2(1^6, -1^5, -39^4, -38^3, 36^2, -27^1, 1^0)$. We first show that d is an increasing function. We find the derivative of d as $d' = (4^3, -12^2, -4^1, -4^0)$. We have $d'(\alpha) \geq \lim_{\alpha \rightarrow 9^+} (4^3) - \lim_{\alpha \rightarrow \alpha_1^-} (12^2, 4^1, 4^0) > 0$ since both of the functions (4^3) and $(12^2, 4^1, 4^0)$ are increasing over the interval I_α . Therefore, $d(\alpha)$ is an increasing function and $d(\alpha) \geq d(9) > 0$ for every $\alpha \in I_\alpha$.

Let us show that $R_1(\alpha)$ is a positive function over the interval I_α . It is enough to show that $(1^4, -2^3, -2^2, -42^1, 5^0) > 0$ for every $\alpha \in I_\alpha$ which follows from the inequality $(1^4, -2^3, -2^2, -42^1, 5^0) \geq (\min_{\alpha \in I_\alpha} (1^4, 5^0) - \max_{\alpha \in I_\alpha} (2^3, 2^2, 42^1)) > 0$. As a result, we derive that $R_1(\alpha)$ is a positive function over I_α . Then we obtain that $R_1 d^{1/2} + R_2 \geq (\min_{\alpha \in I_\alpha} d^{1/2}) R_1 + R_2 = R_1(3456)^{1/2} + R_2$.

Let $K = 3456^{1/2} > 0$. We show that $KR_1 + R_2 > 0$ for every $\alpha \in I_\alpha$. Note that we have $KR_1 + R_2 = 2(\alpha - 1)^2(u_1 - u_2)$ where we set $u_1 = (1^6, K^4, 19^2, 164^1, 5K^0)$ and $u_2 = (4^5, 47^4, 2K^3, 2K^2, 5^0)$. We see that $u_1(\alpha) - u_2(\alpha) > 0$ for every $\alpha \in I_\alpha$ which follows from $u_1(\alpha) - u_2(\alpha) \geq \min_{\alpha \in I_\alpha} u_1(\alpha) - \max_{\alpha \in I_\alpha} u_2(\alpha) = u_1(9) - u_2(\alpha_1) > 0$ for every $\alpha \in I_\alpha$ since both of the functions $u_1(\alpha)$ and $u_2(\alpha)$ are clearly increasing functions. Therefore, we find that $KR_1 + R_2 > 0$ for every $\alpha \in I_\alpha$. We already have that $d^{1/2}(\alpha^2 + 4\alpha - 1 + d^{1/2})^2(\alpha + 1)^2(\alpha - 1)^2 > 0$ over I_α . Thus, we get that $(H_+)' < 0$ for every $\alpha \in I_\alpha$. In other words, $H_+(\alpha)$ is a decreasing function over the interval I_α . The following calculation completes the first part of the lemma:

$$H_+(\alpha_1) = \frac{(\sqrt{\alpha_1} + 1)(\alpha_1 - 1)(g(\alpha_1))^2 + 2(\alpha_1 - 1)g(\alpha_1) + \sqrt{\alpha_1} + 3}{(\sqrt{\alpha_1} + 1)((\alpha_1 - 1)g(\alpha_1) + 1)} > \frac{1}{2}.$$

For the second part of the lemma, let us calculate the derivative of $H_-(\alpha)$ where

$$(H_-)'(\alpha) = \frac{R_1 \sqrt{d} + R_2}{\sqrt{d}(\alpha^2 + 4\alpha - 1 + \sqrt{d})^2(\alpha + 1)^2(\alpha - 1)^2}$$

for $d = (1^1, -1^0)(1^3, -3^2, -5^1, -9^0)$, $R_1 = -2((1^1, -1^0))^2(1^4, -2^3, -2^2, -42^1, 5^0)$, $R_2 = 2((1^1, -1^0))^2(1^6, -1^5, -39^4, -38^3, 36^2, -27^1, 1^0)$. We already showed that d is an increasing function and R_1 is a positive function over the interval I_α . Then we obtain that $R_1 d^{1/2} + R_2 \leq (\max_{\alpha \in I_\alpha} d^{1/2}) R_1 + R_2 = R_1(1972 + 1392\sqrt{2})^{1/2} + R_2$. Let us say $K = (1972 + 1392\sqrt{2})^{1/2} > 0$. We show that $KR_1 + R_2 < 0$ for every $\alpha \in I_\alpha$. We have $KR_1 + R_2 = -2(\alpha - 1)^2(t_1 - t_2)$ where $t_1(\alpha) = (4^5, (47 + K)^4, (5K + 5)^0)$ and $t_2(\alpha) = (1^6, (2K)^3, (19 + 2K)^2, (42K + 164)^1)$. We see that $t_1 - t_2 > 0$ over I_α since we have $t_1 - t_2 \geq \min_{\alpha \in I_\alpha} t_1(\alpha) - \max_{\alpha \in I_\alpha} t_2(\alpha) = t_1(9) - t_2(\alpha_1) > 0$. Because both of the functions $t_1(\alpha)$ and $t_2(\alpha)$ are clearly increasing functions. Therefore, we find that $KR_1 + R_2 < 0$ for every $\alpha \in I_\alpha$. We already have that the expression $d^{1/2}(\alpha^2 + 4\alpha - 1 + d^{1/2})^2(\alpha + 1)^2(\alpha - 1)^2 > 0$ for every $\alpha \in I_\alpha$. Therefore, we get

that $(H_-)' < 0$ for every $\alpha \in \text{int}(I_\alpha)$. In other words, $H_-(\alpha)$ is a decreasing function over the interval I_α . The following calculation completes the first part of the lemma:

$$H_-(\alpha_1) = \frac{(\sqrt{\alpha_1} + 1)(\alpha_1 - 1)(h(\alpha_1))^2 + 2(\alpha_1 - 1)h(\alpha_1) + \sqrt{\alpha_1} + 3}{(\sqrt{\alpha_1} + 1)((\alpha_1 - 1)h(\alpha_1) + 1)} > \frac{1}{2}.$$

A.3 Proof of Lemma 4.10

Remember that we have $x_1^* = x_7^*$ in Case IV and $K_2 = \Sigma_2^* + x_7^*$. From the expressions $\rho_1^* \sigma_8^* = \alpha_*$ and $\rho_1^* \sigma_4^* < \alpha_*$, we derive that $x_8^* < x_4^*$. By the assumption of (i), we know that $x_1^* + x_3^* < x_8^*$. Therefore, we get $0 < x_1^* + x_3^* < 1/3$. Let $I = (0, 1/3 - K_1)$ and $K = K_1 + K_2$ where $K_1 = x_3^*$ and $K_2 = \Sigma_4^*$. We need to show that $H'_{\alpha_*}(x_1^*) < 0$.

Let us calculate the derivative of $H_{\alpha_*}(x)$ using logarithmic differentiation:

$$\frac{H'_{\alpha_*}}{H_{\alpha_*}} = \frac{(1 + h'_{\alpha_*}) K_2(K_2 - 1 + 2x + 2h_{\alpha_*} + 2K_1)}{(1 - x - h_{\alpha_*} - K_1)(x + h_{\alpha_*} + K)(x + h_{\alpha_*} + K_1)(1 - x - h_{\alpha_*} - K)}.$$

We have $(\Sigma_4^* - 1 + 2\Sigma_1^*)\Sigma_4^* = (\Sigma_1^* - x_8^*)\Sigma_4^* < 0$ by the assumption of (i). Therefore, we obtain that $K_2(K_2 - 1 + 2x_1^* + 2K_1 + 2h_{\alpha_*}(x_1^*)) = (\Sigma_4^* - 1 + 2\Sigma_1^*)\Sigma_4^* < 0$. Remember that we have $H_{\alpha_*}(x_1^*) = \alpha_* > 0$ and $h_{\alpha_*}(x_1^*) = x_2^*$. Also, we see that

$$(1 - x - h_{\alpha_*} - K_1)(x + h_{\alpha_*} + K)(x + h_{\alpha_*} + K_1)(1 - x - h_{\alpha_*} - K) > 0$$

at $x = x_1^*$ because $x_1^* + x_3^* + h_{\alpha_*}(x_1^*) = \Sigma_1^* > 0$, $x_1^* + x_3^* + h_{\alpha_*}(x_1^*) + K_2 = \Sigma_{1,2}^* > 0$, $1 - (x_1^* + x_3^*) - h_{\alpha_*}(x_1^*) = 1 - \Sigma_1^* > 0$, and $1 - (x_1^* + x_3^*) - h_{\alpha_*}(x_1^*) - K_2 = x_7^* + x_8^* > 0$. Therefore, it is enough to show that $1 + h'_{\alpha_*}(x_1^*) > 0$. To this purpose, we show that $1 + h'_\alpha(x) > 0$ for every $(x, \alpha) \in I \times I_\alpha$. For any fixed $\alpha \in I_\alpha$, we obtain that

$$1 + h'_\alpha(x) = \frac{1}{2} + \frac{1}{2} \cdot \frac{(\alpha - 1)(1 - x - K_1) + 2}{\sqrt{(1 - \alpha)^2(1 - x - K_1)^2 + 4(1 - \alpha)(x + K_1)}}.$$

It is clear that we have $(\alpha - 1)(1 - x - K_1) + 2 > 0$ for every $(x, \alpha) \in I \times I_\alpha$. Remember that we set $g_\alpha(x) = (1 - \alpha)^2(1 - x - K_1)^2 + 4(1 - \alpha)(x + K_1)$. For $\alpha \in I_\alpha$, we calculate that $g'_\alpha(x) = -2(1 - \alpha)^2(1 - x - K_1) + 4(1 - \alpha) < 0$ for every $x \in I$. Therefore, $g_\alpha(x)$ is a strictly decreasing function over I for all $\alpha \in I_\alpha$. As a result, we see that $g_\alpha(x) > \lim_{x \rightarrow (1/3 - K_1)^+} g_\alpha(x) = (1 - \alpha)^2(4/9) + (1 - \alpha)(4/3) > 0$ for every $\alpha \in I_\alpha$. Then, we get $1 + h'_\alpha(x) > 0$ over $I \times I_\alpha$. In particular, this proves that $1 + h'_{\alpha_*}(x_1^*) > 0$ which in turn means that $H'_{\alpha_*}(x_1^*) < 0$. Finally, we conclude that $H_{\alpha_*}(x)$ is a strictly decreasing function around a neighborhood of x_1^* .

A.4 Proof of Lemma 4.11

We aim to show that $(F_{\alpha_*}^1)_y(x_1^*, x_3^*) \neq (F_{\alpha_*}^2)_y(x_1^*, x_3^*)$. Let us introduce the region $R = \{(x, y) \in \mathbb{R}^2 : 0 < x < 1/2, 0 < y < 1/4, y < x\}$. We know that $(x_1^*, x_3^*) \in R$. We calculate the partial derivatives $(F_{\alpha}^1)_y$ and $(F_{\alpha}^2)_y$. We get the functions below

$$(F_{\alpha}^1)_y = -1 - \left(\frac{1}{2} + \frac{(\alpha - 1)(1 - x - U) + 2}{\sqrt{d_2(x, y)}} \right) U_y,$$

$$(F_{\alpha}^2)_y = -\frac{1}{2} + \frac{(\alpha - 1)(1 - x - y) + 2}{\sqrt{D_1(x, y)}}.$$

It is clear that $(F_{\alpha}^2)_y(x, y) > -1/2$ since we have $(\alpha - 1)(1 - x - y) + 2 > 0$ on the region R for every $\alpha \in I_{\alpha}$. In particular, we find that $(F_{\alpha_*}^2)_y(x_1^*, x_3^*) > -1/2$. Note that the partial derivative of U with respect to y is given as

$$U_y = \frac{(\tau(y)/\sigma(x))}{2(\sqrt{\sigma(y)/\sigma(x)} + 1)^2 \sqrt{\sigma(y)/\sigma(x)}} > 0$$

for every $(x, y) \in R$. Since we have $U(x_1^*, x_3^*) = \Sigma_2^* < 1/2$ by the condition (v) and U is continuous on R , there is a neighborhood R' of (x_1^*, x_3^*) so that $U(x, y) < 1/2$ for every $(x, y) \in R'$. Therefore, we see that $(F_{\alpha_*}^1)_y(x, y) < -1$ for every $(x, y) \in R' \subset R$. In particular, we find that $(F_{\alpha_*}^1)_y(x_1^*, x_3^*) < -1$. By the Implicit Function Theorem, we conclude that there exists a neighborhood $R'' \subset R'$ of (x_1^*, x_3^*) such that y can be written as a function of x .

A.5 Proof of Lemma 4.12

Before we present the proof of Lemma 4.12, we need to prove the following statement

Lemma A.1 *Assume that Case IV and the inequalities $\Sigma_1^* > x_8^*$, $x_1^* < x_8^*$, and $\Sigma_2^* < 1/2$ hold at the point $\mathbf{x}^*$. Then we have*

$$\frac{x_2^*(1 - x_2^* - \Sigma_1^*)(1 - \Sigma_1^*)}{x_8^*(\Sigma_1^* - x_8^*)\Sigma_1^*} \neq \frac{x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*)}{(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*)}.$$

Proof Using the equalities $\rho_1^*\sigma_8^* = \alpha_*$, $\rho_1^*\sigma_2^* = \alpha_*$ and $\rho_2^*\sigma_3^* = \alpha_*$ in Case IV, we obtain expressions $(1 - \Sigma_1^*)(1 - x_8^*) = \alpha_*(\Sigma_1^*)x_8^*$, $(1 - x_2^*)\Sigma_1^* = \alpha_*(1 - \Sigma_1^*)x_2^*$ and $\Sigma_2^*(1 - x_3^*) = \alpha_*(1 - \Sigma_2^*)x_3^*$, respectively. Since we have $\Sigma_0^* = 1$ and $x_1^* = x_7^*$, we find that

$$(24) \quad (1 - \Sigma_1^*)(1 - x_8^*) = \alpha_* x_8^* \Sigma_1^*$$

$$(25) \quad (1 - x_2^*)\Sigma_1^* = \alpha_*(1 - \Sigma_1^*)x_2^*$$

$$(26) \quad (1 - x_1^* - \Sigma_1^* - x_8^*)(1 - x_3^*) = \alpha_*(x_1^* + \Sigma_1^* + x_8^*)x_3^*.$$

We expand each sides of the equalities in (24), (25) and (26). We multiply the expression in (24) by 1, in (25) by -1 and in (26) by -1 . Then we add all the equalities in (24), (25) and (26) side-by-side. We see that $K = \alpha_* K$ where we have $K = x_1^* x_8^* - x_2^* + x_2^* x_8^* + x_1^* x_2^* + (x_2^*)^2 - 2x_1^* x_3^* - (x_3^*)^2$. Since we have $\alpha_* \in I_\alpha$, we conclude that

$$(27) \quad x_1^* x_8^* - x_2^* + x_2^* x_8^* + x_1^* x_2^* + (x_2^*)^2 - 2x_1^* x_3^* - (x_3^*)^2 = 0.$$

We will use the equality in (27) in this proof.

Remember that we have $x_1^* = x_7^*$ in this case. Let us consider the quadratic polynomial $q_4(x) + q_5(x) = 2(\alpha_* - 1)x^2 + (1 - \alpha_*)x + 1$. We solve the equation $q_4(x) + q_5(x) = 0$. We obtain the following solutions

$$x_{+,-} = \frac{1}{4} \pm \frac{1}{4} \left(1 - \frac{8}{\alpha_* - 1} \right)^{1/2}.$$

Let us define the following two functions over the interval I_α :

$$x_+(\alpha) = \frac{1}{4} + \frac{1}{4} \left(1 - \frac{8}{\alpha - 1} \right)^{1/2} \text{ and } x_-(\alpha) = \frac{1}{4} - \frac{1}{4} \left(1 - \frac{8}{\alpha - 1} \right)^{1/2}.$$

Note that we have $x_+(\alpha_*) = x_+$ and $x_-(\alpha_*) = x_-$. We know that the inequalities $q_4(\Sigma_1^*) + q_5(\Sigma_1^*) = q_5(\Sigma_1^*) < 0$, $q_4(\Sigma_2^*) + q_5(\Sigma_2^*) = q_4(\Sigma_2^*) > 0$ hold together with $q_4(x_1^*) + q_5(x_1^*) = q_4(x_1^*) > 0$ and $q_4(x_8^*) + q_5(x_8^*) = q_5(x_8^*) < 0$ because we have $x_1^* < x_8^*$ and $x_8^* < \Sigma_1^*$ and $\Sigma_1^* < \Sigma_2^*$. Therefore, we derive that $x_- < x_8^* < \Sigma_1^* < x_+$ for any given $\alpha \in I_\alpha$. We divide the interval I_α into two intervals U and V so that $U = [\hat{\alpha}, \alpha_1]$ and $V = [\alpha_1, \alpha_2]$ where $\hat{\alpha} = 9.118994613$. First, we concentrate on the interval U .

Since $x_-(\alpha)$ is a decreasing function over the interval U , the smallest value of $x_-(\alpha)$ is obtained at $\alpha = \hat{\alpha}$. As a consequence, we see that

$$x_- \geq \frac{1}{4} - \frac{1}{4} \left(1 - \frac{8}{\hat{\alpha} - 1} \right)^{1/2} = a_1.$$

Therefore, we get that $a_1 < x_8^*$ for every $\alpha \in U$. It is clear that $x_8^* < 1/4$. Because if we have $1/4 < x_8^* < \Sigma_1^*$, we get $\alpha_* = \rho_1^* \sigma_8^* < 9$, a contradiction. If we have $x_8^* = 1/4$, then we get $\Sigma_1^* > 1/4$. We find that $\alpha_* = \rho_1^* \sigma_8^* \leq 9$. We know that $\alpha_* \geq 9$. Therefore, we find $\alpha_* = 9$. This implies that $\Sigma_1^* = 1/4$. This contradicts with hypothesis of Lemma A.1. Hence, we see that $a_1 < x_8^* < 1/4$. Let us call $a_2 = 1/4$.

It is possible to find bounds for the sum Σ_1^* using the equation $\rho_1^* \sigma_8^* = \alpha_*$. We know that $a_1 < x_8^* < a_2$. Then we derive the inequality $\rho_1^* \sigma(a_1) > \alpha_* > \rho_1^* \sigma(a_2)$. Since we

have $\alpha_* \geq 9$, we obtain $\Sigma_1^* < (1 - a_1)/(8a_1 + 1)$. Let us call $a_3 = (1 - a_1)/(8a_1 + 1)$. We know that $\alpha_* \leq \hat{\alpha}$. We find that $\Sigma_1^* > (1 - a_2)(\hat{\alpha}a_2 - a_2 + 1)$. Let us call $a_4 = (1 - a_2)/(\hat{\alpha}a_2 - a_2 + 1)$. Finally, we get $a_4 < \Sigma_1^* < a_3$.

Using the equation $\rho_*^1 \sigma_2^* = \alpha_*$, it is possible to find a lower and an upper bound for x_2^* . We know that $a_4 < \Sigma_1^* < a_3$. Then we derive the following inequalities $\sigma_2^*/\sigma(a_4) < \alpha_* < \sigma_2^*/\sigma(a_3)$. Since we know that $\alpha_* \leq \hat{\alpha}$, we find the expression $x_2^* > a_4/(\hat{\alpha}(1 - a_4) + a_4)$. Let us call $a_5 = a_4/(\hat{\alpha}(1 - a_4) + a_4)$. We know that $\alpha_* \geq 9$. Then we obtain that $x_2^* > a_3/(9 - 8a_3)$. Let us call $a_6 = a_3/(9 - 8a_3)$. We have $a_5 < x_2^* < a_6$.

We follow the operations below to find a lower and an upper bound for x_1^* . The equality in (27) implies that $x_1^*x_8^* - x_2^* + x_2^*x_8^* + x_1^*x_2^* + (x_2^*)^2 = 2x_1^*x_3^* + (x_3^*)^2$. Then we get $(x_1^*)^1 + 2x_1^*x_3^* + (x_3^*)^2 = (x_1^*)^2 + x_1^*x_8^* - x_2^* + x_2^*x_8^* + x_1^*x_2^* + (x_2^*)^2$ or equivalently $(x_1^*)^1 + 2x_1^*x_3^* + (x_3^*)^2 + x_2^* - (x_2^*)^2 - x_2^* - x_8^* = (x_1^*)^2 + x_1^*x_8^* + x_2^*x_8^*$. We add the term $((x_2^* + x_8^*)/2)^2$ to both sides of the last equality. We see that the righthand side is a complete square $(x_1^* + (x_2^* + x_8^*)/2)^2$. Then we take the square root of both side of the resulting equality and isolate the term x_1^* . We see that

$$x_1^* = \left((x_1^* + x_3^*)^2 + x_2^* - \frac{3}{4}(x_2^*)^2 + \frac{1}{4}(x_8^*)^2 \right)^{1/2} - \frac{x_2^* + x_8^*}{2}.$$

Let us define the function

$$f(x, y, z, t, u, v) = \left((x + y)^2 + z - \frac{3}{4}z^2 + \frac{1}{4}t^2 \right)^{1/2} - \frac{u + v}{2}.$$

Note that we have $f(x_1^*, x_3^*, x_2^*, x_8^*, x_2^*, x_8^*) = x_1^*$. We want to find the minimum lower bound for x_1^* using the bounds for x_8^* , Σ_1^* and x_2^* . We add side by side the inequalities $a_4 < \Sigma_1^* < a_3$ and $-a_6 < -x_2^* < -a_5$ to obtain that $a_4 - a_6 < x_1^* + x_3^* < a_3 - a_5$. We see that $f(a_4, -a_6, x_2^*, x_2^*, x_8^*, x_2^*, x_8^*) < x_1^*$. Since the function $z - (3/4)z^2$ is an increasing function over $(a_5, a_6) \subset (0, 1)$, we find that $f(a_4, -a_6, a_5, x_8^*, x_2^*, x_8^*) < x_1^*$. We conclude that $f(a_4, -a_6, a_5, a_1, a_6, a_2) < x_1^*$ because the square root function is an increasing function and f is the difference of two functions. Let us declare that $a_7 = f(a_4, -a_6, a_5, a_1, a_6, a_2)$.

We want to find the maximum upper bound for x_1^* using the bounds for x_8^* , Σ_1^* and x_2^* . Note that we can say $f(a_3, -a_5, x_2^*, x_2^*, x_8^*, x_2^*, x_8^*) > x_1^*$. Using the fact that $z - (3/4)z^2$ is an increasing function over $(a_5, a_6) \subset (0, 1)$ again, we derive $f(a_3, -a_5, a_6, x_8^*, x_2^*, x_8^*) > x_1^*$. We replace the values x_2^* and x_8^* with a_2 and a_5 because f is the difference of two functions and the square root function is increasing. In particular, we obtain that $f(a_3, -a_5, a_6, a_2, a_1, a_5) > x_1^*$. Let $a_8 = f(a_3, -a_5, a_6, a_2, a_1, a_5)$. Therefore, we have $a_7 < x_1^* < a_8$.

It is possible to find an upper and a lower bound for Σ_2^* using $\rho_2^* \sigma_1^* = \alpha_*$. We use the inequalities $a_7 < x_1^* < a_8$. We find that $\rho_2^* \sigma(a_7) > \alpha_* > \rho_2^* \sigma(a_8)$. Since we have $\alpha_* \geq 9$, we obtain that $\Sigma_2^* < (1 - a_7)/(8a_7 - 1)$. Let us declare that $a_9 = (1 - a_7)/(8a_7 - 1)$. Similarly, since we know that $\alpha_* \leq \hat{\alpha}$, we derive the inequality $\Sigma_2^* > (1 - a_8)/(\hat{\alpha}a_8 - a_8 + 1)$. Let us call $a_{10} = (1 - a_8)/(\hat{\alpha}a_8 - a_8 + 1)$. Then we have $a_{10} < \Sigma_2^* < a_9$.

Using the equality $\rho^2 \sigma_3^* = \alpha_*$ in Case IV, we find a lower bound and an upper bound for x_3^* . We use the inequality $a_{10} < \Sigma_2^* < a_9$. We derive the following inequality $\sigma_3^*/\sigma(a_{10}) < \alpha_* < \sigma_3^*/\sigma(a_9)$. We know that $\alpha_* \geq 9$. By the previous inequalities, we find that $x_3^* < a_9/(9 - 8a_9)$. Let $a_{11} = a_9/(9 - 8a_9)$. Since we know that $\alpha_* \leq \hat{\alpha}$, we obtain that $x_3^* > a_{10}/(\hat{\alpha}(1 - a_{10}) + a_{10})$. Let us call $a_{12} = a_{10}/(\hat{\alpha}(1 - a_{10}) + a_{10})$. Then we have $a_{12} < x_3^* < a_{11}$.

As a result, we calculate the following lower and upper bounds $a_1 < x_8^* < a_2$, $a_4 < \Sigma_1^* < a_3$, $a_5 < x_2^* < a_6$, $a_7 < x_1^* < a_8$, $a_{10} < \Sigma_2^* < a_9$, $a_{12} < x_3^* < a_{11}$ where

$$\begin{aligned}
a_1 &= \frac{1}{4} - \frac{1}{4} \left(1 - \frac{8}{\hat{\alpha} - 1} \right)^{1/2} && \approx 0.2197341744, \\
a_2 &= \frac{1}{4} && = 0.2500000000, \\
a_3 &= \frac{1 - a_1}{8a_1 + 1} && \approx 0.2829230040, \\
a_4 &= \frac{1 - a_2}{\hat{\alpha}a_2 - a_2 + 1} && \approx 0.2475452871, \\
a_5 &= \frac{\hat{\alpha}(1 - a_4) + a_4}{1 - a_4} && \approx 0.03482053272, \\
a_6 &= \frac{1 - a_4}{8a_4 + 1} && \approx 0.04199779316, \\
a_7 &= \left((a_4 - a_6)^2 + a_5 - \frac{3}{4}a_5^2 + \frac{1}{4}a_1^2 \right)^{1/2} - \frac{a_6 + a_2}{2} && \approx 0.1510393664, \\
a_8 &= \left((a_3 - a_5)^2 + a_6 - \frac{3}{4}a_6^2 + \frac{1}{4}a_2^2 \right)^{1/2} - \frac{a_1 + a_5}{2} && \approx 0.2160224693, \\
a_9 &= \frac{1 - a_7}{8a_7 + 1} && \approx 0.3844382075, \\
a_{10} &= \frac{1 - a_8}{\hat{\alpha}a_8 - a_8 + 1} && \approx 0.2846805352, \\
a_{11} &= \frac{a_9}{9 - 8a_9} && \approx 0.06488962356, \\
a_{12} &= \frac{a_{10}}{\hat{\alpha}(1 - a_{10}) + a_{10}} && \approx 0.04181758681.
\end{aligned}$$

Then we calculate that $x_2^*(1 - x_2^* - \Sigma_1^*) > a_5(1 - a_6 - a_3)$. Similarly, we find that $x_8^*(\Sigma_1^* - x_8^*) < a_2(a_3 - a_1)$, because we have $x_8^* < a_2$, $a_4 < \Sigma_1^* < a_3$ and $-x_8^* < -a_1$.

As a result, we calculate that the inequality below holds at the point $\mathbf{x}^*$

$$\frac{x_2^*(1 - x_2^* - \Sigma_1^*)(1 - \Sigma_1^*)}{x_8^*(\Sigma_1^* - x_8^*)\Sigma_1^*} > \frac{a_5(1 - a_6 - a_3)(1 - a_3)}{a_2(a_3 - a_1)a_3} \approx 3.771441275.$$

We know that $x_1^* < 1/2$, $x_3^* < 1/2$ and $\Sigma_2^* < 1/2$. Since the function $x - x^2$ is an increasing function over the interval $(0, 1/2)$, we derive the following inequality

$$x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*) < \max_{x_1^* \in (a_7, a_8)} x_1^*(1 - x_1^*) - \min_{x_3^* \in (a_{12}, a_{11})} x_3^*(1 - x_3^*)$$

or $x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*) < a_8(1 - a_8) - a_{12}(1 - a_{12})$. Similarly, we calculate that

$$(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*) > \min_{\Sigma_2^* \in (a_{10}, a_9)} \Sigma_2^*(1 - \Sigma_2^*) - \max_{x_1^* \in (a_7, a_8)} x_1^*(1 - x_1^*)$$

or $(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*) > a_{10}(1 - a_{10}) - a_8(1 - a_8)$. Then we find that

$$\frac{x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*)}{(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*)} < \frac{a_8(1 - a_8) - a_{12}(1 - a_{12})}{a_{10}(1 - a_{10}) - a_8(1 - a_8)} \approx 3.771440985.$$

Therefore, in the case $\alpha_* \in U$, we conclude that

$$\frac{x_2^*(1 - x_2^* - \Sigma_1^*)(1 - \Sigma_1^*)}{x_8^*(\Sigma_1^* - x_8^*)\Sigma_1^*} \neq \frac{x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*)}{(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*)}.$$

We follow the same calculations carried above for the interval V . Since $x_-(\alpha)$ is a decreasing function over the interval V , the smallest value of $x_-(\alpha)$ is obtained at $\alpha = \alpha_1$. We have

$$x_- \geq \frac{1}{4} - \frac{1}{4} \left(1 - \frac{8}{\alpha_1 - 1} \right)^{1/2}.$$

Let us call $a_1 = 1/4 - 1/4(1 - 8/(\alpha_1 - 1))^{1/2}$. Therefore, we get that $a_1 < x_8^*$ for every $\alpha \in V$. We showed in the previous case that $x_8^* < 1/4$. Let us call $a_2 = 1/4$. Then we have $a_1 < x_8^* < a_2$.

It is possible to find a lower and an upper bound for Σ_1^* . We use the inequality $a_1 < x_8^* < a_2$. We know that $\rho_1^*\sigma_8^* = \alpha_*$ from Case IV. Then we derive that $\rho_1^*\sigma(a_1) > \alpha_* > \rho_1^*\sigma(a_2)$. We know that $\alpha_* \geq \hat{\alpha}$. Then we obtain the bound $\Sigma_1^* < (1 - a_1)/(\hat{\alpha}a_1 + 1 - a_1)$. Let us call $a_3 = (1 - a_1)/(\hat{\alpha}a_1 + 1 - a_1)$. Since we know that $\alpha_* \leq \alpha_1$, we find that $\Sigma_1^* > (1 - a_2)/(\alpha_1a_2 - a_2 + 1)$. Let us call $a_4 = (1 - a_2)/(\alpha_1a_2 - a_2 + 1)$. We get $a_4 < \Sigma_1^* < a_3$.

Using the equality $\rho_*^1\sigma_2^* = \alpha_*$, we find a lower and an upper bound for x_2^* . We use the inequality $a_4 < t < a_3$. We obtain that $\sigma_2^*/\sigma(a_4) < \alpha_* < \sigma_2^*/\sigma(a_3)$. Since we know that $\alpha_* \leq \alpha_1$, we find that $x_2^* > a_4/(\alpha_1(1 - a_4) + a_4)$. Let us call

$a_5 = a_4/(\alpha_1(1 - a_4) + a_4)$. We know that $\alpha_* \geq \hat{\alpha}$. Then we obtain the bound $x_2^* < a_3/(\hat{\alpha}(1 - a_3) + a_3)$. Let us say $a_6 = a_3/(\hat{\alpha}(1 - a_3) + a_3)$. We get $a_5 < x_2^* < a_6$.

Remember that we have $f(x_1^*, x_3^*, x_2^*, x_8^*, x_2^*, x_8^*) = x_1^*$. We want to find the minimum lower bound for x_1^* using the bounds for x_8^* , Σ_1^* and x_2^* . Therefore, we add side by side the inequalities $a_4 < \Sigma_1^* < a_3$ and $-a_6 < -x_2^* < -a_5$ to obtain that $a_4 - a_6 < x_1^* + x_3^* < a_3 - a_5$. We see that $f(a_4, -a_6, x_2^*, x_2^*, x_8^*, x_2^*, x_8^*) < x_1^*$. Since the function $z - (3/4)z^2$ is an increasing function over $(a_5, a_6) \subset (0, 1)$, we find that $f(a_4, -a_6, a_5, x_8^*, x_2^*, x_8^*) < x_1^*$. We conclude that $f(a_4, -a_6, a_5, a_1, a_6, a_2) < x_1^*$ because the square root function is an increasing function and f is the difference of two functions. Let us say $a_7 = f(a_4, -a_6, a_5, a_1, a_6, a_2)$.

We want to find the maximum lower bound for x_1^* using the bounds for x_8^* , Σ_1^* and x_2^* . Note that we can say $f(a_3, -a_5, x_2^*, x_2^*, x_8^*, x_2^*, x_8^*) > x_1^*$. Using the fact that $z - (3/4)z^2$ is an increasing function over $(a_5, a_6) \subset (0, 1)$ again, we derive that $f(a_3, -a_5, a_6, x_8^*, x_2^*, x_8^*) > x_1^*$. We replace the values x_2^* and x_8^* with a_2 and a_5 because f is the difference of two functions and the square root function is increasing. In particular, we obtain that $f(a_3, -a_5, a_6, a_2, a_1, a_5) > x_1^*$. Therefore, we have the inequality $a_7 < x_1^* < a_8$.

We find an upper and a lower bound for Σ_2^* using the equality $\rho_2^* \sigma_1^* = \alpha_*$. We have the inequalities $a_7 < x_1^* < a_8$. We see that $\rho_2^* \sigma(a_7) > \alpha_* > \rho_2^* \sigma(a_8)$. Since we know that $\alpha_* \geq \hat{\alpha}$, we obtain $\Sigma_2^* < (1 - a_7)/(\hat{\alpha}a_7 + 1 - a_7)$. Let us call $a_9 = (1 - a_7)/(\hat{\alpha}a_7 + 1 - a_7)$. Similarly, we know that $\alpha_* \leq \alpha_1$. Therefore, we derive the inequality $\Sigma_2^* > (1 - a_8)/(\alpha_1 a_8 - a_8 + 1)$. Let us call this bound $a_{10} = (1 - a_8)/(\alpha_1 a_8 - a_8 + 1)$. Then we have $a_{10} < \Sigma_2^* < a_9$.

Using the equality $\rho^2 \sigma_3^* = \alpha_*$ in Case IV, we calculate a lower bound and an upper bound for x_3^* . We use the inequality $a_{10} < \Sigma_2^* < a_9$. We derive the expression $\sigma_3^*/\sigma(a_{10}) < \alpha_* < \sigma_3^*/\sigma(a_9)$. We know that $\alpha_* \geq \hat{\alpha}$. By the previous inequalities, we find that $x_3^* < a_9/(\hat{\alpha}(1 - a_9) + a_9)$. Let $a_{11} = a_9/(\hat{\alpha}(1 - a_9) + a_9)$. Since we know that $\alpha_* \leq \alpha_1$, we obtain $x_3^* > a_{10}/(\alpha_1(1 - a_{10}) + a_{10})$. Let us call $a_{12} = a_{10}/(\hat{\alpha}(1 - a_{10}) + a_{10})$. Then we have $a_{12} < x_3^* < a_{11}$.

Hence, we get the followings $a_1 < x_8^* < a_2$, $a_4 < \Sigma_1^* < a_3$, $a_5 < x_2^* < a_6$,

$a_7 < x_1^* < a_8$, $a_{10} < \Sigma_2^* < a_9$, and $a_{12} < x_3^* < a_{11}$ where we evaluate the bounds as

$$\begin{aligned}
a_1 &= \frac{1}{4} - \frac{1}{4} \left(1 - \frac{8}{\alpha_1 - 1} \right)^{1/2} && \approx 0.2197341744, \\
a_2 &= \frac{1}{4} && = 0.2500000000, \\
a_3 &= \frac{1 - a_1}{\hat{\alpha}a_1 + 1 - a_1} && \approx 0.2802658255, \\
a_4 &= \frac{1 - a_2}{\alpha_1 a_2 - a_4 + 1} && \approx 0.2450451726, \\
a_5 &= \frac{a_4}{\alpha_1(1 - a_4) + a_4} && \approx 0.03392651705, \\
a_6 &= \frac{a_3}{\hat{\alpha}(1 - a_3) + a_3} && \approx 0.04095347309, \\
a_7 &= \left((a_4 - a_6)^2 + a_5 - \frac{3}{4}a_5^2 + \frac{1}{4}a_1^2 \right)^{1/2} - \frac{a_6 + a_2}{2} && \approx 0.1491203770, \\
a_8 &= \left((a_3 - a_5)^2 + a_6 - \frac{3}{4}a_6^2 + \frac{1}{4}a_2^2 \right)^{1/2} - \frac{a_1 + a_5}{2} && \approx 0.2137627221, \\
a_9 &= \frac{1 - a_7}{\hat{\alpha}a_7 + 1 - a_7} && \approx 0.3817066009, \\
a_{10} &= \frac{1 - a_8}{\alpha_1 a_8 - a_8 + 1} && \approx 0.2846654650, \\
a_{11} &= \frac{a_9}{\hat{\alpha}(1 - a_9) + a_9} && \approx 0.06257148633, \\
a_{12} &= \frac{a_{10}}{\alpha_1(1 - a_{10}) + a_{10}} && \approx 0.04127832503.
\end{aligned}$$

Then we calculate that $x_2^*(1 - x_2^* - \Sigma_1^*) > a_5(1 - a_6 - a_3)$. Similarly, we find that $x_8^*(\Sigma_1^* - x_8^*) < a_2(a_3 - a_1)$, because we have $x_8^* < a_2$, $a_4 < \Sigma_1^* < a_3$ and $-x_8^* < -a_1$. Then we obtain that

$$\frac{x_2^*(1 - x_2^* - \Sigma_1^*)(1 - \Sigma_1^*)}{x_8^*(\Sigma_1^* - x_8^*)\Sigma_1^*} > \frac{a_5(1 - a_6 - a_3)(1 - a_3)}{a_2(a_3 - a_1)a_3} \approx 3.907942936.$$

We know that $x_1^* < 1/2$, $x_3^* < 1/2$ and $\Sigma_2^* < 1/2$. Since the function $x - x^2$ is an increasing function over the interval $(0, 1/2)$, we derive the following inequality

$$x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*) < \max_{x_1^* \in (a_7, a_8)} x_1^*(1 - x_1^*) - \min_{x_3^* \in (a_{12}, a_{11})} x_3^*(1 - x_3^*)$$

or $x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*) < a_8(1 - a_8) - a_{12}(1 - a_{12})$. Similarly, we calculate that

$$(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*) > \min_{\Sigma_2^* \in (a_{10}, a_9)} \Sigma_2^*(1 - \Sigma_2^*) - \max_{x_1^* \in (a_7, a_8)} x_1^*(1 - x_1^*)$$

or $(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*) > a_{10}(1 - a_{10}) - a_8(1 - a_8)$. Then we find that

$$\frac{x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*)}{(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*)} < \frac{a_8(1 - a_8) - a_{12}(1 - a_{12})}{a_{10}(1 - a_{10}) - a_8(1 - a_8)} \approx 3.613150070.$$

Therefore, in the case $\alpha_* \in V$, we conclude that the following expression holds at $\mathbf{x}^*$

$$\frac{x_2^*(1 - x_2^* - \Sigma_1^*)(1 - \Sigma_1^*)}{x_8^*(\Sigma_1^* - x_8^*)\Sigma_1^*} \neq \frac{x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*)}{(1 - \Sigma_2^*)\Sigma_2^* - x_1^*(1 - x_1^*)}.$$

□

We are ready to prove Lemma 4.12. We use logarithmic differentiation to calculate the following:

$$\frac{H'_{\alpha_*}}{H_{\alpha_*}} = -\frac{1}{2} \cdot \frac{h_{\alpha_*}(1 - h_{\alpha_*}) + x(1 - x)h'_{\alpha_*}}{xh_{\alpha_*}(1 - x)(1 - h_{\alpha_*})}.$$

We need to show that $H'_{\alpha_*}(x_1^*) \neq 0$. Let us assume otherwise. Then we get the equation $h_{\alpha_*}(x_1^*)(1 - h_{\alpha_*}(x_1^*)) + x_1^*(1 - x_1^*)h'_{\alpha_*}(x_1^*) = 0$. Solving the previous equality for $h'_{\alpha_*}(x_1^*)$, we find that $h'_{\alpha_*}(x_1^*) = -(x_3^*(1 - x_3^*)) / (x_1^*(1 - x_1^*))$. By Lemma 4.11, we know that $h_{\alpha}(x) = y(x)$ and $y(x_1^*) = x_3^*$. Using $F_{\alpha_*}^1(x, h_{\alpha_*}(x)) = F_{\alpha_*}^2(x, h_{\alpha_*}(x))$, we calculate $h'_{\alpha_*}(x)$. We obtain that

$$(28) \quad \frac{d'_2}{2\sqrt{d_2}} + \frac{D'_1}{2\sqrt{D_1}} = (\alpha_* - 1)(2 + h'_{\alpha_*} + U')$$

where

$$\frac{dU}{dx} = -\frac{\frac{1 - h_{\alpha_*}}{h_{\alpha_*}(1 - x)} - \frac{xh'_{\alpha_*}}{h_{\alpha_*}(1 - x)} - \frac{x(1 - h_{\alpha_*})h'_{\alpha_*}}{h_{\alpha_*}^2(1 - x)} + \frac{x(1 - h_{\alpha_*})}{h_{\alpha_*}(1 - x)^2}}{2 \left(1 + \sqrt{\sigma(h_{\alpha_*})/\sigma(x)}\right)^2 \sqrt{\sigma(h_{\alpha_*})/\sigma(x)}}.$$

We evaluate the derivative above at the point x_1^* . It is possible to simplify $U'(x_1^*)$ using the equalities given in $\rho_*^2\sigma_3^* = \alpha_*$ and $\rho_*^2\sigma_1^* = \alpha_*$ as follows:

$$\begin{aligned} U'(x_1^*) &= -\frac{\sigma_3^*\tau_*^1}{(1 + \sqrt{\sigma_3^*/\sigma_1^*})^2 \sqrt{\sigma_3^*/\sigma_1^*}} \\ &= -\rho_*^2\sigma_3^*\tau_*^1(U(x_1^*, x_3^*))^2 \\ &= -\alpha_*\tau_*^1(U(x_1^*, x_3^*))^2 \\ &= -\rho_*^2\sigma_1^*\tau_*^1(U(x_1^*, x_3^*))^2 \\ &= -\frac{\Sigma_2^*(1 - \Sigma_2^*)}{x_1^*(1 - x_1^*)}. \end{aligned}$$

Let $\delta_* = \alpha_* - 1$. We have $d'_2(x_1^*) = -2\delta_*(1 + U'(x_1^*))(\delta_*(1 - x_1^* - U(x_1^*, x_3^*)) + 2)$ and $D'_1(x_1^*) = -2\beta_*(1 + h'_{\alpha_*}(x_1^*))(\delta_*(1 - x_1^* - h_{\alpha_*}(x_1^*)) + 2)$. We simplify the equality in (28) by multiplying both sides with $4\sqrt{d_2}\sqrt{D_1}$. We find that

$$(29) \quad d'_2\sqrt{D_1} + D'_1\sqrt{d_2} = 2\delta_*\sqrt{d_2}\sqrt{D_1}(2 + h'_{\alpha_*} + U')$$

Note that we have $\sqrt{D_1(x_1^*, x_3^*)} = \delta_*(1 - x_2^* - \Sigma_1^*)$ and $\sqrt{d_2(x_1^*, x_3^*)} = \delta_*(\Sigma_1^* - x_8^*)$. We plug the two expressions above into the terms $d'_2\sqrt{D_1}$ and $D'_1\sqrt{d_2}$. We obtain

$$\begin{aligned} d'_2\sqrt{D_1} &= -2\delta_*^2(1 + U'(x_1^*)) (\delta_*(1 - x_1^* - U(x_1^*, x_3^*)) + 2) (1 - \Sigma_1^* - x_2^*), \\ D'_1\sqrt{d_2} &= -2\delta_*^2(1 + h'_{\alpha_*}(x_1^*)) (\delta_*(1 - x_1^* - h_{\alpha_*}(x_1^*)) + 2) (\Sigma_1^* - x_8^*), \end{aligned}$$

$$(30) \quad 2\sqrt{d_2}\sqrt{D_1}\delta_*(2 + h'_{\alpha_*} + U') = 2\delta_*^3(1 - \Sigma_1^* - x_2^*)(\Sigma_1^* - x_8^*)(2 + h'_{\alpha_*} + U').$$

Let $A_1 = (\delta_*(1 - x_1^* - U(x_1^*, x_3^*)) + 2)(1 - x_2^* - \Sigma_1^*)$, $A_3 = (1 - x_2^* - \Sigma_1^*)(\Sigma_1^* - x_8^*)$, and $A_2 = (\delta_*(1 - x_1^* - h_{\alpha_*}(x_1^*)) + 2)(\Sigma_1^* - x_8^*)$. The term $2(\alpha_* - 1)^2$ is a common factor for left-hand side of the equality in (29) and the right-hand sides of the equality in (30). We divide both sides with $2(\alpha_* - 1)^2$. We perform the operations

$$\begin{aligned} -(1 + U')A_1 - (1 + h'_{\alpha_*})A_2 &= \delta_*A_3(2 + h'_{\alpha_*} + U') \\ -(1 + U')A_1 - (1 + h'_{\alpha_*})A_2 &= \delta_*A_3(1 + h'_{\alpha_*} + 1 + U') \\ (31) \quad -(1 + U')(\delta_*A_3 + A_1) &= (1 + h'_{\alpha_*})(\delta_*A_3 + A_2). \end{aligned}$$

Let $(\alpha_* - 1)A_3 + A_1 = C_1$. Using the equality $\rho_1^*\sigma_8^* = \alpha_*$, we simplify C_1 as follows:

$$\begin{aligned} C_1 &= ((\alpha_* - 1)(1 - x_1^* - U(x_1^*, x_3^*) + \Sigma_1^* - x_8^*) + 2)(1 - \Sigma_1^* - x_2^*) \\ &= 2(\alpha_* - 1)\Sigma_1^*(1 - x_2^* - \Sigma_1^*) + 2(1 - x_2^* - \Sigma_1^*) \\ &= 2(1 - x_2^* - \Sigma_1^*)((\alpha_* - 1)\Sigma_1^* + 1) \\ &= 2(1 - x_2^* - \Sigma_1^*)((\rho_1^*\sigma_8^* - 1)\Sigma_1^* + 1) \\ (32) \quad &= 2(1 - x_2^* - \Sigma_1^*) \cdot \frac{1 - \Sigma_1^*}{x_8^*}. \end{aligned}$$

Let $(\alpha_* - 1)A_3 + A_2 = C_2$. Using the equality $\rho_1^*\sigma_2^* = \alpha_*$, we simplify C_2 as follows:

$$\begin{aligned} C_2 &= ((\alpha_* - 1)(2 - x_1^* - h_{\alpha_*}(x_1^*) - \Sigma_1^* - x_8^*) + 2)(\Sigma_1^* - x_8^*) \\ &= ((\alpha_* - 1)(2 - 2\Sigma_1^*) + 2)(\Sigma_1^* - x_8^*) \\ &= 2(\Sigma_1^* - x_8^*)((\rho_1^*\sigma_2^* - 1)(1 - \Sigma_1^*) + 1) \\ (33) \quad &= 2(\Sigma_1^* - x_8^*) \cdot \frac{\Sigma_1^*}{x_2^*}. \end{aligned}$$

We know that

$$(34) \quad -(1 + U'(x_1^*)) = \frac{\Sigma_2^*(1 - \Sigma_2^*) - x_1^*(1 - x_1^*)}{x_1^*(1 - x_1^*)},$$

$$(35) \quad (1 + h'_{\alpha_*}(x_1^*)) = \frac{x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*)}{x_1^*(1 - x_1^*)}.$$

We plug the expressions in (32), (33), (34) and (35) into the equality in (31). We simplify both sides of the equation in (31). We find that

$$\frac{x_2^*(1 - x_2^* - \Sigma_1^*)(1 - \Sigma_1^*)}{x_8^*(\Sigma_1^* - x_8^*)\Sigma_1^*} = \frac{x_1^*(1 - x_1^*) - x_3^*(1 - x_3^*)}{\Sigma_2^*(1 - \Sigma_2^*) - x_1^*(1 - x_1^*)}.$$

Lemma A.1 shows that this is not the case, therefore, a contradiction. As a result, we conclude that $H'_{\alpha_*}(x_1^*) \neq 0$ which means that $H_{\alpha_*}(x)$ is a strictly monotone function around a neighborhood of x_1^* .

References

- [1] Ian Agol, Tameness of hyperbolic 3-manifolds, <http://www.arXiv.org:math/0405568>.
- [2] Vicki Chuckrow, On Schottky groups with applications to Kleinian groups, in *Ann. of Math. (2)*, Volume 88, 1968, pp. 47–61.
- [3] Danny Calegari and David Gabai, Shrinkwrapping and the taming of hyperbolic 3-manifolds, in *J. Amer. Math. Soc.*, Volume 2, 2006, pp 385
- [4] Richard D. Canary, Ends of hyperbolic 3-manifolds, *J. Amer. Math. Soc.*, Volume 6, 1993, No 1, pp. 1–35
- [5] Albert Marden, The geometry of finitely generated Kleinian groups, *Ann. of Math. (2)*, Volume 99, 1974, pp. 383–462.
- [6] Colin Maclachlan and Alan W. Reid, *The arithmetic of hyperbolic 3-manifolds*, Graduate Texts in Mathematics, (Springer-Verlag, 219).
- [7] S. J. Patterson, Lectures on measures on limit sets of Kleinian groups, *Fundamentals of hyperbolic geometry: selected expositions*, London Math. Soc. Lecture Note Ser., Volume 328, pp. 291–335
- [8] John G. Ratcliffe, *Foundations of hyperbolic manifolds*, Graduate Texts in Mathematics, Springer 149, 2006
- [9] Marc Culler and Peter B. Shalen, Paradoxical decompositions, 2-generator Kleinian groups, and volumes of hyperbolic 3-manifolds, *J. Amer. Math. Soc.*, Volume 5, 1992, No 2, 231–288.
- [10] Marc Culler and Peter B. Shalen, Margulis Numbers For Haken Manifolds, <http://www.arXiv.org:math/1006.3467v1>.

Department of Mathematics, Boston College, Chestnut Hill, Massachusetts 02467-3806, USA

yucei@bc.edu

www2.bc.edu/~yucei